



## “Survival of the Fittest” Does Obesity Trait Evolved Over Time Confers Survival Advantage to Critically Ill Patients?

Noman Shahzad\*, Abdul Rehman Alvi, Romana Iqbal, Shaquat Rozi, Fatima Mannan and Delvene Soares

Department of Surgery, Aga Khan University Hospital Karachi, Pakistan

### Abstract

There is a steady increase in prevalence of obesity over last 2-3 decades to the extent of global epidemic. Overall 25% of world population is reported to be overweight and 10% are obese. The evolutionary origin of obesity points towards survival advantage of obese individuals, but in modern way of living, advantages of obesity are lost and hazardous effects have become more prominent including cardio-metabolic risk factors and some malignancies. There is very scanty information in medical literature about value of fat stores in critically ill patients. Available literature has pointed towards the potential benefits of fat store in combating the disease process in critically ill patients. Though obesity is known to adversely affect the overall health of individuals, its positive role in recovery of critically ill patients needs to be further evaluated.

### Introduction

Obesity refers to excess of body fat, also known as adiposity. Distribution of this excess of body fat varies from individual to individual and can be broadly classified into visceral obesity and body wall obesity. In visceral obesity, fat accumulates around abdominal viscera predominantly. It is known to correlate with adverse outcomes of obesity [1]. On the contrary, predominant accumulation of fat in abdominal wall is known to have relatively less hazardous effects [2]. There are various parameters to measure obesity, commonly used are anthropometric measures like Body Mass Index (BMI), Waist Circumference, Mid-arm Circumference and Triceps skin fold thickness [3,4]. Bioelectrical impedance is also used to measure composition of body during various physiological states [5].

### Burden and Health Hazards of Obesity

There is a steady increase in prevalence of obesity over last 2-3 decades to the extent of global epidemic [6,7]. Overall 25% of world population is overweight and 10% is obese [8]. Health hazards of obesity are well established. Obesity is known to lead to higher prevalence of cardio metabolic risk factors including diabetes mellitus, hypertension and hyperlipidemia [9,10] and certain malignancies [11,12]. It is known to be an independent risk factor for all-cause mortality [13]. Overweight and obesity are the fifth leading risk for global deaths. At least 2.8 million adults die each year as a result of being overweight or obese. In addition, 44% of the diabetes burden, 23% of the ischemic heart disease burden and between 7% and 41% of certain cancer burdens are attributable to overweight and obesity [14]. Once considered a high-income country problem, overweight and obesity are now on the rise in low- and middle-income countries, particularly in urban settings. Obesity is estimated to account for between 0.7% and 2.8% of a country's total healthcare expenditures. Furthermore, medical costs of obese individuals are approximately 30% greater than their normal weight peers [15].

### Obesity from Evolutionary Perspective

Survival of the fittest is one of the principles of most famous and accepted theory of Evolutionary origin of species by Darwin [16]. According to this principle, every individual has different set of genes that confer him/her a set of certain characteristics. New characteristic are acquired through random genetic mutations in nature. There are some characteristics that give an individual a survival advantage over others. As a result of this natural selection process, survival of these individuals is better and they get a chance to spread their genes to their offspring. According to this theory, every prevalent trait of human beings must have been acquired through this rigorous process of natural selection and must have given survival advantage at some point in the process of evolution. This is contrary to what we see about obesity in the modern age.

### OPEN ACCESS

#### \*Correspondence:

Noman Shahzad, Department of Surgery, Aga Khan University Hospital Karachi, Pakistan, Tel: +92-300-8833313;

E-mail: drns01@hotmail.com

Received Date: 24 Jun 2018

Accepted Date: 10 Jul 2018

Published Date: 17 Jul 2018

#### Citation:

Shahzad N, Alvi AR, Iqbal R, Rozi S, Mannan F, Soares D. “Survival of the Fittest” Does Obesity Trait Evolved Over Time Confers Survival Advantage to Critically Ill Patients?. *Ann Short Reports*. 2018; 1: 1013.

Copyright © 2018 Noman Shahzad.

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Looking at evolutionary origin of obesity, it is paramount to critically analyse the survival advantage that fat stores would have conferred to individuals. Life of evolutionary less advanced species was faced with frequent and often prolonged periods of famines. Along with that, extensive physical activity was required for daily activities to survive. Individuals having energy reserves were at survival advantage. These energy reserves evolved in the form of fat stores. These fat stores turned out to be valuable energy rich body reserves for them to be utilized to combat periods of famines and provide energy to meet metabolic demands to sail through periods of famines. These were also readily available sources of energy to be supplied for exertional activities. As a result, those individuals who had adequate fat stores had survival advantage and had natural selection to transfer their genes to their offspring's. These reservoirs also gave advantage to them in better coping with physical injuries and infections as fat stores were also utilized in process of repair and regeneration in recovering from physical injuries which were not uncommon in forest life.

With modern civilization, there is progressive change in environment and physical work demand. There is increased food security and as a result famines are rare. Advancement in technology, machines have taken over a lot of physical work reducing physical work load requirements for human beings. Physical injuries and combats are also far less in civilian life. So there is change in environment, but energy conserving and fat storing genes are the same. There is excessive accumulation of fat stores. These fat stores are rarely utilized for the purpose they were evolved for, and modern day human beings are prone to get obese [17,18]. In the absence of their use, counterproductive effects of these fat stores become evident in the form of cardiometabolic risk factors and some malignancies.

### **Fat Stores in Critically Ill Patients**

Patients are said to be critically ill if as a result of their disease they have laboratory and clinical manifestations of systemic involvement of disease, compromised hemodynamic status, and organ dysfunction. Scoring systems have been developed to grade the illness. One of these scoring systems is Acute Physiology and Chronic Health Evaluation (APACHE) II score [19].

Trauma and infections are the two most common causes of development of critical illness [20]. Critically ill patients have increased metabolic and immune requirements as compared to those of healthy individuals. A functional repair and regeneration mechanism is required for repair process and to counter physical insults. These pathways are energy dependent and hence metabolic rate of these individuals is higher as compared to their healthy counter-parts. To meet these energy requirements, fat stores can play an important role and can be an aide for them to sail through their illness. Body requires a strong immune system to fight with infection. Immunity is also energy dependent. Energy reserves in the form of fat stores can help support the immune system. It has been seen that increased levels of leptin in the body secondary to increased fat stores is known to strengthen the immune response and is associated with better sepsis outcome [21]. Thus critically ill obese patients are at least theoretically at an advantage of having energy rich fat reserves as compared to their non-obese counterparts.

### **Available Literature Evidence**

A few investigators have reported impact of obesity upon outcome of critically ill patients with varying results. Some of these

studies have shown that obese individuals are not at increased risk of morbidity or mortality [22,23]. Pieracci et al. [23] in retrospective review of their intensive care unit data have shown that BMI is not associated with mortality. Similar results have been shown by Saker Y et al. [22]. There are reports that have shown protective effect of obesity in critical illness [24,25]. For example Hutagalung R et al. [25] reviewed outcome of their 13000 intensive care unit admission patients over a period of five year and found that being overweight or obese was associated with decreased 60-days in hospital mortality. Similarly Piak SL et al. [24] analysed one year data of their intensive care unit patients. They found that BMI was determinant of short to medium term survival. They concluded that obesity was not associated with increased morbidity and could be protective for critically ill patients. Similarly Tafelski S et al. [26] demonstrated in their study that despite increase in mechanical ventilator requirements, obesity is not a risk factor for mortality in critically ill patients [26]. Recent study published by Shahzad N et al. [27] also supports the idea that being overweight puts critically ill patients in advantageous position [27]. It has been seen that fat utilization if obese patients is better when suffering from critical illness and this helps them conserve their muscle proteins [28].

On the other hand, there are some studies that have shown obesity to be a risk factor for increased morbidity or mortality [29]. Duchesne JC et al. [29] reviewed one hundred and four trauma patients who were admitted to intensive care unit. They found that severe obesity was significantly associated with adverse outcomes and increased resource utilization in trauma patients treated admitted to ICU. A meta-analysis of related studies was also conducted in 2008 by Akinnusi ME et al. [30]. They analysed fourteen studies having about sixty two thousand patients collectively. They found that obesity in critically ill patients is not associated with excess mortality but is significantly related to prolonged duration of mechanical ventilation and intensive care unit length of stay.

Problem with available studies evaluating impact of obesity upon outcome of critically ill patients is that most of these studies are either retrospective or are secondary analysis of data collected for some other purpose.

### **Future Perspective**

Meticulous knowledge of impact of obesity upon outcome of critically ill patients has important clinical implications in the management of these patients. Improved risk stratification of individual patients according to their obesity level will help in decision making especially those concerning surgical intervention. Critically ill patients are at high risk for development of intra-operative or post-operative complications, after any intervention irrespective of their level of obesity. This in turn has impact upon management of these patients.

### **Conclusion**

Though obesity is known to adversely affect the overall health of individuals, its positive role in recovery of critically ill patients needs to be further evaluated.

### **Conflicts of Interest**

The Authors declare that there is no conflict of interest.

### **References**

1. Zhang C, Rexrode KM, van Dam RM, Li TY, Hu FB. Abdominal obesity

- and the risk of all-cause, cardiovascular and cancer mortality: sixteen years of follow-up in US women. *Circulation*. 2008;117(13):1658-67.
2. Snijder MB, van Dam RM, Visser M, Seidell JC. What aspects of body fat are particularly hazardous and how do we measure them? *Int J Epidemiol*. 2006;35(1):83-92.
  3. Kartheuser AH, Leonard DF, Penninckx F, Paterson HM, Brandt D, Remue C, et al. Waist circumference and waist/hip ratio are better predictive risk factors for mortality and morbidity after colorectal surgery than body mass index and body surface area. *Ann Surg*. 2013;258(5):722-30.
  4. Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults--The Evidence Report. National Institutes of Health. *Obes Res*. 1998;6(Suppl 2):51S-209S.
  5. Schneider PL, Bassett Jr DR, Thompson DL, Crouter SE. Bioelectrical Impedance for Accuracy Detecting Body Composition Changes during an Activity Intervention. *Translational Journal of the American College of Sports Medicine*. 2017;2(19):122-8.
  6. Shen J, Goyal A, Sperling L. The emerging epidemic of obesity, diabetes, and the metabolic syndrome in china. *Cardiol Res Pract*. 2012;2012:178675.
  7. Wang Y. Cross-national comparison of childhood obesity: the epidemic and the relationship between obesity and socioeconomic status. *Int J Epidemiol*. 2001;30(5):1129-36.
  8. Kelly T, Yang W, Chen CS, Reynolds K, He J. Global burden of obesity in 2005 and projections to 2030. *Int J Obes (Lond)*. 2008;32(9):1431-7.
  9. Nguyen NT, Nguyen XM, Lane J, Wang P. Relationship between obesity and diabetes in a US adult population: findings from the National Health and Nutrition Examination Survey, 1999-2006. *Obes Surg*. 2011;21(3):351-5.
  10. Wang YC, McPherson K, Marsh T, Gortmaker SL, Brown M. Health and economic burden of the projected obesity trends in the USA and the UK. *Lancet*. 2011;378(9793):815-25.
  11. Kaneko R. A New Index of Abdominal Obesity which Effectively Predicts Risk of Colon Tumor Development in Female Japanese. *Asian Pac J Cancer Prev*. 15(2):1005-10.
  12. Dalamaga M. Obesity, insulin resistance, adipocytokines and breast cancer: New biomarkers and attractive therapeutic targets. *World J Exp Med*. 2013;3(3):34-42.
  13. Twig G, Afek A, Shamiss A, Derazne E, Landau Rabbi M, Tzur D, et al. Adolescence BMI and Trends in adulthood Mortality: A Study of 2.16 Million Adolescents. *J Clin Endocrinol Metab*. 2014;99(6):2095-103.
  14. Scheen AJ, Van Gaal LF. Combating the dual burden: therapeutic targeting of common pathways in obesity and type 2 diabetes. *Lancet Diabetes Endocrinol*. 2014;2(11):911-22.
  15. Withrow D, Alter DA. The economic burden of obesity worldwide: a systematic review of the direct costs of obesity. *Obes Rev*. 2011;12(2):131-41.
  16. Calabi L. On Darwin's 'metaphysical notebooks'. I: teleology and the project of a theory. *Riv Biol*. 2001;94(1):123-59.
  17. Bellisari A. Evolutionary origins of obesity. *Obes Rev*. 2008;9(2):165-80.
  18. Levaie Ran A. Human obesity: an evolutionary approach to understanding our bulging waistline. *Diabetes Metab Res Rev*. 2001;17(5):347-62.
  19. Fleig V, Brenck F, Wolff M, Weigand MA. [Scoring systems in intensive care medicine: principles, models, application and limits]. *Anaesthesist*. 2011;60(10):963-74.
  20. Niven DJ, Bastos JF, Stelfox HT. Critical care transition programs and the risk of readmission or death after discharge from an ICU: a systematic review and meta-analysis. *Crit Care Med*. 2014;42(1):179-87.
  21. Siegl D, Annecke T, Johnson BL, Schlag C, Martignoni A, Huber N, et al. Obesity-induced Hyperleptinemia Improves Survival and Immune Response in a Murine Model of Sepsis. *Anesthesiology*. 2014;121(1):98-114.
  22. Sakr Y, Madl C, Filipescu D, Moreno R, Groeneveld J, Artigas A, et al. Obesity was associated with increased morbidity but not mortality in critically ill patients. *Intensive Care Med*. 2008;34(11):1999-2009.
  23. Pieracci FM. The relationship between body mass index and postoperative mortality from critical illness. *Obesity surgery*. 2008;18(5):501-507.
  24. Peake SL, Moran JL, Ghelani DR, Lloyd AJ, Walker MJ. The effect of obesity on 12-month survival following admission to intensive care: A prospective study. *Critical Care Med*. 2006;34(12):2929-39.
  25. Hutagalung R, Marques J, Kobylka K, Zeidan M, Kabisch B, Brunkhorst F, et al. The obesity paradox in surgical intensive care unit patients. *Intensive Care Med*. 2011;37(11):1793-9.
  26. Tafelski S, Yi H, Ismael F, Krannich A, Spies C, Nachtigall I. Obesity in critically ill patients is associated with increased need of mechanical ventilation but not with mortality. *Journal of infection and public health*. 2016;9(5):577-85.
  27. Shahzad N, Alvi AR, Iqbal R, Rozi S, Mannan F, Khan MH, et al. Obesity and the Outcome of Surgical Intensive Care Unit Patients: A Prospective Cohort Study. *Clinics of Surgery*. 2018;1(1): 1-9.
  28. Goossens C, Marques MB, Derde S, Vander Perre S, Dufour T, Thiessen SE, et al. Premorbid Obesity Protects Against Critical Illness-Induced Muscle Wasting and Weakness. In *Energy Balance and Body Weight (posters) 2016 Apr (pp. SUN-619)*. Endocrine Society.
  29. Duchesne JC, Schmieg RE, Simmons JD, Islam T, McGinness CL, McSwain NE. Impact of obesity in damage control laparotomy patients. *J Trauma*. 2009;67(1):108-112.
  30. Hogue CW, Stearns JD, Colantuoni E, Robinson KA, Stierer T, Mitter N, et al. The impact of obesity on outcomes after critical illness: a meta-analysis. *Intensive Care Med*. 2009;35(7):1152-1170.