



Methicillin- and Vancomycin-Resistant *Staphylococcus Aureus* on Computer Keyboards Located in Different Wards of a Third-Level Hospital in Tehran, Iran

Ali Badamchi¹, Zahra Movahed², Shima Javidinia³, Mohammad Reza Shokrollah², Mehri Naghadalipoor⁴, Shirin Sayyahfar⁴ and Azardokht Tabatabaei^{1*}

¹Department of Lung disease, Tehran University of Medical Sciences, Iran

²Department of Lung disease, Qom University of Medical Sciences and Health services, Iran

³Department of Lung disease, Zabol University of Medical Sciences, Iran

⁴Department of Pediatric Infectious Disease, Iran University of Medical Sciences, Iran

Abstract

Introduction: Nosocomial infections are one of the health problems in all societies. The aim of this study was to evaluate contamination of computer keyboards located in different wards of Rasoul-e-Akram hospital in Tehran with methicillin-resistant *Staphylococcus aureus*.

Material and Methods: This cross-sectional study performed to identify the microorganisms in the Rasoul-e-Akram Hospital, 90 samples collected from different wards of this hospital by sterile swab for culture on EMB agar and blood agar media and used for differentiation test to identify the types of microorganisms. Data analyzed by using SPSS24 Software.

Results: The results of this study showed that *Staphylococcus* bacteria are the most common type of microorganisms in infectious wards (59.41%) includes *Staphylococcus coagulase negative* (42.02%) and *S. aureus* (17.39%). Also *S. aureus* isolated from PICU (50%), ICU(25%) and operating room (25%) were MRSA. one *S. aureus* was isolated with vancomycin MICs >2µg/ml.

Conclusion: Results showed the high incidence of microorganisms that causes nosocomial infections in different wards and there is need to use proper disinfectants to prevent the spread of these agents in a hospital environment.

Keywords: Nosocomial infections; Hospital; Hethicillin-resistant *Staphylococcus aureus*

Introduction

Every year, millions of people across the world suffer from Healthcare-associated Infections (HAI). Infections may also occur at surgery sites, known as surgical site infections [1]. HAIs may be caused by infectious agents from endogenous (body sites) or exogenous sources (patient care personnel, visitors, patient care equipment, medical devices or the health care environment) [2]. These infections of ten have little or nothing to do with the primary reason for the hospital visit but are consequence of deprived hygiene in the healthcare setting [3]. Healthcare equipment is commonly shared between hospital workers, who may have various hygiene practices. Computers and telephones are now found in all healthcare locations and their disinfection is often ignored more than medical devices. Therefore, the opportunity for the transmission of contaminating bacteria is potentially great [4-8]. There has been much debate over the infection risk to patients from contaminated health care surfaces [9]. It has been reported that, devices and different items in hospital environments, such as telephones and computers, are highly associated with HAIs transmission [3-7].

It is now recognized that the health care environment may facilitate transmission of several important health care-associated pathogens, including vancomycin-resistant *enterococci* (VRE) and methicillin-resistant *Staphylococcus aureus* (MRSA) [8,10]. These pathogens are frequently transmitted between patients and healthcare workers and consequently to the devices that are commonly used by other staff [10,11].

It is highly recommended that computer keyboards and mice used in patient care areas should

OPEN ACCESS

*Correspondence:

Hazrat-e-Rasoul, Department of Pediatric Infectious Disease, Iran University of Medical, Sciences and Health Services, Iran, Tel: +98- 21- 66516049; E-mail: azardokht_tabatabaei@yahoo.com

Received Date: 12 Feb 2018

Accepted Date: 20 Mar 2018

Published Date: 27 Mar 2018

Citation:

Badamchi A, Movahedi Z, Javidinia S, Shokrollahi MR, Naghadalipoor M, Sayyahfar S, et al. Methicillin- and Vancomycin-Resistant *Staphylococcus Aureus* on Computer Keyboards Located in Different Wards of a Third-Level Hospital in Tehran, Iran. *Ann Infect Dis Epidemiol.* 2018; 3(1): 1028.

ISSN: 2475-5664

Copyright © 2018 Azardokht

Tabatabaei. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1: Frequency of positive microbial culture according to type of microbial Contamination.

Bacterial isolates	No. of bacterialisolates from		
	Keyboards	mice	Total (%)
<i>Staphylococcus coagulase negative</i>	35	23	58(42.02)
<i>S. aureus</i>	18	6	24(17.39)
<i>Bacillus spp</i>	10	10	20(14.49)
<i>Diphtheroidspp</i>	10	7	17(12.31)
<i>Gram negative bacteria (fermentative)</i>	0	1	1(0.74)
<i>Pseudomonas spp</i>	0	1	1(0.74)
<i>Fungal</i>	17	0	17(12.31)
Total	90	48	138(100)

Table 2: The frequency of *Staphylococcus* species of isolated from variety of wards hospital.

Wards Hospital	Number of computers	Number of sample	<i>Staphylococcus species(n)</i>	
	In wards		CoNS	S.aureus
Emergency	4	8	8	0
PICU	3	12	5	6
ICU	2	8	4	4
Operating Room	2	8	4	4
Internal	3	6	5	0
Blood	3	6	5	0
Skin	3	6	5	0
Obstetrics and gynecology	3	6	5	0
Ophthalmology	2	8	5	2
SportClinic, heart	3	6	4	0
ClinicofNeurology	1	2	1	0
Psychiatry	2	8	4	4
lungandENT	3	6	1	4
Neurosurgery	2	4	1	0
Pediatric	2	4	1	0
Total€	38	98	58	24

be disinfected each day however it is neglected in almost all sectors of the public health system. In this study the contamination of computer keyboards located in different wards of Rasoul-e- Akram hos0pital with methicillin-resistant *Staphylococcus aureus* was investigated.

Materials and Methods

Sampling

A cross-sectional study conducted at Rasoul-e-Akram Hospital, Tehran, Iran, since April 1 t\o August 31/2015. Samples were taken from 98 keyboards and mice of multiple-user computers in 15 departments of the Hospital using pre-sterilized swabs (Table 1). The sterile swabs immersed in Trypticase Soy Broth (TSB) and used to obtain samples from the computer keyboards and mice by slowly, but firmly, rotating the swab on the different buttons and surfaces of the computers. The swabs were then re-immersed in the TSB broth to prevent from drying and were taken immediately to the microbiology laboratory for further processing (Table 2).

Isolation and Characterization

Swabs were cut in sterile condition and the head swabs were inoculated into TSB media in 5 ml tubes. The tubes were incubated at 37°C for 24 hours and aliquots were inoculated on blood agar,

McConkey agar and Sabouraud dextrose agar plates. The preliminary characterization of the isolates performed based on colony morphology, Gram-staining, catalase and oxidase tests, bacterial isolates. Also, a slide coagulase test (Microgen Staph, UK) used to differentiate isolates of staphylococci into *S.aureus* and coagulase-negative *staphylococci*. A tube coagulase test used for confirmation (Figure 1).

Detection of Methicillin-Resistance

Methicillin-resistant *S. aureus* (MRSA) strains were confirmed on Mueller Hinton Agar (MHA) containing 4% NaCl (Oxoid) and supplemented with 6 µg/ml Oxacillin (Oxoid). All *S. aureus* strains were tested for methicillin-resistance using the disk diffusion method (Cefoxitin 30 µg, B D, Germany) in 33 to 35°C ambient air for 24 hours to identify MRSA (Table 3).

Antimicrobial susceptibility testing

Susceptibility of *S. aureus* isolates to Cefoxitin (Fox), Penicillin (P), Erythromycin (E), Clindamycin (Cd), sulfamethoxazole trimethoprim (sxt), Ciprofloxacin (Cip), Gentamicin (GM), Tetracyclin (Te) and Vancomycin (V) (MAST Group, United Kingdom) was determined by disc diffusion method according to the guidelines of Clinical and Laboratory Standards Institute (CLSI) [12]. All methicillin resistant

Table 3: Frequency of MRSA of isolated from variety of wards hospital.

Source sample	Noof <i>S.aureus</i>	No. of MRSA (%)	No. of MSSA (%)
ICU	4	1(25)	3(75)
PICU	6	3(50)	3(50)
Operating Room	4	1(25)	3(75)
Ophthalmology ward	2	0	2(100)
Psychiatry Ward	4	0	4(100)
Lung and ENT ward	4	0	4(100)
Total	24	5(20.8)	19(79.2)

strains were collected and MICs of vancomycin among MRSA isolates were determined by Etest (AB, Bio merieux, France) according to the manufacturer's instructions and were repeated by CLSI guidelines [12].

Detection of Coding Genes for Methicillin Resistance

DNA extraction

The genomic DNA was extracted from *S. aureus* isolate using QIA amp genomic DNA kit (Qiagen, Germany) as per manufacturer's protocol [13,14].

PCR-Amplification and Detection of *mecA* Genes

Detection of *mecA* gene by polymerase chain reaction (PCR) was considered the gold standard. A *mecA* negative strain of *S. aureus* (ATCC[®] 25923 (Cefoxitin zone 23–29 mm)) and A *mecA* positive strain of *S. aureus* (ATCC[®] 43300 (Cefoxitin zone \leq 21 mm)) were used as negative and positive controls, respectively. The primers used for PCR-amplifications were forward 5'-TCGAGGTAAGGTTGGCC-3', Reverse 5'-AGTTCTGCAGTACCGGATTGTC-3' [15]. The PCR mixture was prepared in a final volume of 25 μ L. The amplification mixture consisted of a 2.5 μ L template DNA, 2 μ L primers, 2 μ L of a 10-fold concentrate PCR buffer, 2 μ L dNTP, 0.5 mM MgCl₂, 15 μ L D.W. and 1 U of Taq DNA (QiaGen, Germany). The PCR program for the detection of the *mecA* gene was: initial denaturation at 94°C: 3 min, followed by 30 cycles of amplification with 94°C: 30 seconds, annealing at 55°C: 30 seconds, and extension at 72°C: 30 seconds, except for the final cycle, which had an extension step of 4 minutes. PCR products were then visualized on a 1% Agarose gel *MecA* positive isolates detected as 480 bp long amplicons observed on the agarose gel (Figure 1).


Statistical Analysis

Data obtained in the study were descriptively and statistically analyzed using SPSS₂₄ software. The means were separated using one sample t-test ($P < 0.05$) is significant and ($P > 0.05$) is not significant.

Results

We investigated the bacterial contamination of two objects used daily: computer keyboards, computer mice. A total of 98 samples were collected from the two different objects. 87% percent of the total samples collected were contaminated with mixed bacterial growth. Coagulase-negative staphylococci dominated the isolates. The second most common bacterial growth in all specimens was Gram-positive bacilli. Potential pathogens isolated from all specimens were: *Staphylococcus aureus*, *Pseudomonas spp.* and *Gram negative bacilli*.

All computer keyboards, mice were cultured and analyzed for



pathogenic organisms. 1 (1.0%) of the 98 samples were colonized with six different bacteria. 58samples (59.18%) grew gram-positive bacteria, and 2sample (2.0%) grew gram-negative bacteria. 17 sample (17.3 %) grew fungi. The fungal isolates are as follows *Aspergillus sp.*, *Mucor sp.*, *Penicillium sp.* The most commonly isolated species was *Aspergillus*.

In all coagulase-positive isolates, the *mecA* gene was detected by PCR.

In this study, *Staphylococcus aureus* was also isolated 24(17.39 %), of which 19(79.2%) were sensitive to methicillin and 5(20.8%) were methicillin resistant. *S. aureus* isolated from PICU (50%) ,ICU(25%) and operating room (25%) were MRSA. Vancomycin minimum inhibitory concentration (MIC) of resistant isolates was also determined using Etest. one *S. aureus* was isolated with vancomycin MICs $> 2\mu\text{g/ml}$.

Discussion

The emergence of nosocomial infections, especially types of antibiotic resistance, is one of themaj or problems in the hospitals and environmental pollution has fundamental role in the development of nosocomial infections. Surfaces and different equipment in different hospital wards are suitable sites for colonization of microorganisms. Different bacteria and fungi are able to survive on inanimate surfaces for a long time. For instance *Staphylococci* can survive in dried blood for 3 months. Computer devices are widely used in hospital wards. Various studies in different parts of the world had assessed the microbial contamination on computer keyboards. In the present study, all the collected samples from mice and keyboards of computers were contaminated with human pathogenic bacteria. Gram positive bacteria were more frequently isolated from all surfaces compared to

Gram negative. This could be due to the fact that survival of Gram positive species on laminate surfaces is greater than that of Gram negative organisms [13].

Coagulase-Negative *Staphylococcus* is found on skin or in the nasal environment and only survives on dry skin. Its emergence on keyboards is due to the frequent use of these devices. Methylamine resistant *Staphylococcus aureus* (MRSA) which was isolated from the ICU key board, can cause infections in patients [14-16]. Distributions of MRSA isolates were varied in different wards which partly reflected the fact that some patients, e.g., critically ill patients in ICUs, had a greater chance of becoming colonized or infected. *Enterococci* are part of the normal flora in human gut and are adapted to extreme conditions.

These results are somewhat similar with the findings of Alemu. study at the Vali-e-Asr Hospital of Birjand, the bacterial contamination of computer keyboards were included *Enterobacteriaceae*, bacilligram-positive spore-bearing, *CoNS*, *diphtheroid* and *S. aureus* [17]. Bacterial contamination of computer keyboards in hospitals of Isfahan province were also included *Bacillus* species, *CoNS*, *S. aureus*, *Enterococcus* species, *Micrococcus* species, gram negative bacilli, methicillin-resistant *S. aureus* and vancomycin-resistant *Enterococci* [18-21].

According to guidelines proposed by the Centers for Disease Control and Prevention (CDC) in the USA, difficult-to-clean hospital equipment (eg. computer keyboards) should be protected from potential contamination by means of special protective covers [22-26].

This study has demonstrated that microbial contamination of multiple-user computer keyboards may be a common mechanism of transfer of potentially pathogenic bacteria among users.

Conclusion

The isolation of the bacteria from computer keyboards and mice is a clear indication that the sterilization system of health care centers is not effective in reducing the level of the organism on these surfaces to an acceptable level. Microbes are everywhere, therefore, it is highly recommended that hand-washing hygiene should be adopted before and after using the computers to reduce the microbial transmission. Computer keyboards and mice should also be cleaned with alcohol or other disinfectants on a regular basis.

References

- Types of Healthcare-associated Infections. Healthcare-associated Infections.
- Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control*. 2008;36(5):309-32.
- Messina G, Ceriale E, Lenzi D, Burgassi S, Azzolini E, Manzi P. Environmental contaminants in hospital settings and progress in disinfecting techniques. *BioMed research international*. 2013;2013(2013):8.
- Messina G, Ceriale E, Burgassi S, Russo C, Defranceschi C, Mariani L, et al. Impact of a disinfecting technique on microbial contamination of computer keyboards and telephone handsets. *Journal of Hospital Administration*. 2013;2(4):p1.
- Messina G, Quercioli C, Burgassi S, Nisticò F, Lupoli A, Nante N. How many bacteria live on the keyboard of your computer? *Am J Infect Control*. 2011 Sep;39(7):616-8.
- Rutala WA, White MS, Gergen MF, Weber DJ. Bacterial contamination of keyboards: efficacy and functional impact of disinfectants. *Infect Control Hosp Epidemiol*. 2006;27(4):372-7.
- Singh A, Purohit B. Mobile phones in hospital settings: a serious threat to infection. *Occupational health & safety*. 2012;81(3):42.
- Dancer SJ. Importance of the environment in methicillin-resistant *Staphylococcus aureus* acquisition: the case for hospital cleaning. *Lancet Infect Dis*. 2008;8(2):101-13.
- Bures S, Fishbain JT, Uyehara CF, Parker JM, Berg BW. Computer keyboards and faucet handles as reservoirs of nosocomial pathogens in the intensive care unit. *Am J Infect Control*. 2000;28(6):465-71.
- Kramer A, Schwebke I, Kampf G. How long do nosocomial pathogens persist on inanimate surfaces? A systematic review. *BMC Infect Dis*. 2006;6:130.
- Wagenvoort J, Sluijmsmans W, Penders R. Better environmental survival of outbreak vs. sporadic MRSA isolates. *Journal of Hospital Infection*. *J Hosp Infect*. 2000;45(3):231-4.
- Clinical and Laboratory Standards Institute 2015. Performance standards for antimicrobial susceptibility testing; 25th informational supplement. CLSI M100-S25. Clinical and Laboratory Standards Institute, Wayne, PA.
- Saba F, Papizadeh M, Khansha J, Sedghi M, Rasooli M, Amoozgar M A, et al. A rapid and reproducible genomic dna extraction protocol for sequence-based identification of archaea, bacteria, cyanobacteria, diatoms, fungi, and green algae. *Journal of Medical Bacteriology*. 2016;5(3-4):22-28.
- Badamchi A, Javadinia S, Soboti B, Tabatabaee A. Detection of *tstH* gene in *Staphylococcus aureus* isolates from hospitalized burn children. *Journal of Medical Bacteriology*. 2017;6(1-2):1-7.
- Iyer A, Kumosani T, Azhar E, Barbour E, Harakeh S. High incidence rate of methicillin-resistant *Staphylococcus aureus* (MRSA) among healthcare workers in Saudi Arabia. *J Infect Dev Ctries*. 2014;8(3):372-8.
- Enemuor S, Apeh T, Oguntibeju O. Microorganisms associated with computer keyboards and mice in a university environment. *African Journal of Microbiology Research*. 2012;6(20):4424-6.
- Esfandiari A, Salari H, Rashidian A, Masoumi Asl H, Rahimi Foroushani A, Akbari Sari A. Eliminating healthcare-associated infections in Iran: a qualitative study to explore stakeholders' views. *Int J Health Policy Manag*. 2018;7(1):27-34.
- Schultz M, Gill J, Zubairi S, Huber R, Gordin F. Bacterial contamination of computer keyboards in a teaching hospital. *Infect Control Hosp Epidemiol*. 2003;24(4):302-3.
- Karbasizade V, Mohammadi Sichani M, Parsafar S. Bacterial contamination of computer keyboards in hospitals in Isfahan in Iran. *International Journal of Biosciences (IJB)*. 2014;4(1):320-4.
- Hasan NM, Sima S, Hoda K, Nahid A, Alireza JS. Contamination of computer keyboards in various wards of Vali-e-Asr teaching hospital in Birjand, Iran. *Payavard*. 2012;5(5):10-17.
- Karbasizade V, Mohammadi Sichani M, Parsafar S. Bacterial contamination of computer keyboards in hospitals in Isfahan in Iran. *International Journal of Biosciences (IJB)*. 2014;4(1):320-4.
- Kampf G, Kramer A. Epidemiologic background of hand hygiene and evaluation of the most important agents for scrubs and rubs. *Clin Microbiol Rev*. 2004;17(4):863-93.
- Brunetti L, Santoro E, De Caro F, Cavallo P, Boccia A, Capunzo M, et al. Surveillance of nosocomial infections: a preliminary study on hand hygiene compliance of healthcare workers. *Journal of preventive medicine and hygiene*. 2006;47(2).
- Moradi-Tabriz H, Hadadi A, Sotoudeh-Anvari M, Rahimi-Foroushani A, Soleimani T, Mehdipour-Aghabagher B, et al. Prevalence and

- antimicrobial susceptibility of staphylococcus aureus isolated from nasal carriers. *Journal of Medical Bacteriology*. 2014;3(3-4):14-9.
25. Rutala WA, Weber DJ. The Healthcare Infection Control Practices Advisory Committee (HICPAC). Guideline for disinfection and sterilization in healthcare facilities 2008. 2010.
26. Neely AN, Maley MP, Warden GD. Computer keyboards as reservoirs for *Acinetobacter baumannii* in a burn hospital. *Clinical infectious diseases*. 1999;29(5):1358-9.