



Human Coronavirus NL63 and Acute Encephalitis

Ami Schattner*, Shoshana Hadar and Ina Dubin

Departments of Medicine and Neurology, Hebrew University and Hadassah Medical School, Israel

Short Communication

An active, working 70-years-old woman was admitted in December with cough, low-grade fever and altered behavior over one week during which she stopped eating and speaking and became poorly responsive. She had mild, well-controlled diabetes and hypertension but no prior admissions.

Fever (38.6°C) and tachycardia were noted, with apathy, slow speech and disorientation. Neither meningeal irritation signs nor other abnormality were found on examination. On the next day, somnolence with fluctuations, four-limb symmetric spasticity and later, catatonia with incontinence developed. Psychosis was ruled out.

Intravenous ceftriaxone and acyclovir were started, and nasogastric tube inserted for feeding and oseltamivir treatment. Valproate was added when EEG demonstrated temporal epileptiform activity (Lt.>Rt.) and generalized slowing. Laboratory tests showed leukocytosis (15.6X10⁹/L), thrombocytosis (594 X10⁹/L), and CRP 250 mg/dL without metabolic abnormality or autoantibodies. The cerebrospinal fluid was twice normal, including PCR for multiple infectious agents. Chest X-ray showed interstitial changes. Neuroimaging revealed only pansinusitis. Extensive cultures and serology tests remained negative. Repeated nasopharyngeal swabs PCR revealed coronavirus NL63 whereas other coronaviruses, rhinovirus, influenza, para-influenza, RSV, adenovirus and herpesviruses were not detected (BioFire Diagnostics).

Fever and cough abated. On the second week the patient made dramatic improvement gradually regaining her previous functional status. She was discharged home on the 15th hospital day.

Our patient had an acute, self-limited CNS infection manifesting primarily by altered consciousness, confusion, refusal to feed, catatonia and epileptiform activity which were fortunately, completely reversible.

The occurrence in the wintertime, associated respiratory infection and sinusitis, and spontaneous recovery are all consistent with human coronavirus NL63 infection [1,2] - the single pathogen identified.

OPEN ACCESS

*Correspondence:

Ami Schattner, Hebrew University-
Hadassah Medical School, Jerusalem,
Israel, Tel: 972 8 939 0330;
E-mail: amischatt@gmail.com

Received Date: 23 Jul 2018

Accepted Date: 17 Aug 2018

Published Date: 24 Aug 2018

Citation:

Schattner A, Hadar S, Dubin I.
Human Coronavirus NL63 and Acute
Encephalitis. *Ann Infect Dis Epidemiol.*
2018; 3(2): 1032.

ISSN: 2475-5664

Copyright © 2018 Ami Schattner. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Coronaviruses predominantly infect the upper (usually mild illness) and lower (potentially severe disease) respiratory tract. No previous cases of human coronavirus NL63-associated encephalitis have been reported. However, similar coronaviruses have strong CNS tropism, as both clinical and experimental studies have demonstrated [3-5]. The etiology of neurologic disease remains poorly understood and could be due to misdirected host immune responses [5]. Thus, in vulnerable populations such as the elderly, spread from the respiratory tract to the CNS must be considered, and as our patient suggests, coronavirus NL63 should be considered among viral agents capable of causing acute encephalitis in humans.

References

1. Greenberg SB. Update on rhinovirus and coronavirus infections. *Semin Respir Crit Care Med.* 2011;32:433-46.
2. Pitkaranta A, Arruda E, Malmberg H, Hayden FG. Detection of rhinovirus in sinus brushings of patients with acute community acquired sinusitis by reverse transcription-PCR. *J Clin Microbiol.* 1997;35:1791-3.
3. Bergmann CC, Lane TE, Stohman SA. Coronavirus infection of the central nervous system: host-virus stand-off. *Nat Rev Microbiol.* 2006;4:121-32.
4. Arabi YM, Harthi A, Hussein J, Bouchama A, Johani S, Hajeer AH, et al. Severe neurologic syndrome associated with Middle East respiratory syndrome corona virus (MERS-CoV). *Infection.* 2015; 43:495-501.
5. Desforges M, Le Coupanec A, Stodola JK, Meessen-Pinard M, Talbot PJ. Human coronaviruses: viral and cellular factors involved in neuroinvasiveness and neuropathogenesis. *Virus Res.* 2014;194:145-58.