



# Exercise and Enrichment Recruit Positive Factors for Hippocampus Plasticity and Cognitive Enhancement

Naijuan Xu, Lei Zhang and Xinhua Zhang\*

Department of Anatomy, Co-innovative Center of Neuroregeneration, Nantong University, Nantong 226001, China

## Short Communication

The hippocampus plays critical roles in consolidation of information from short- to long-term memory, as well as spatial navigation, which are defined as cognitive function. A number of lifestyle interventions are proven to be cognitive enhancers, such as physical exercise, Environmental Enrichments (EE) including mental stimulation, social stimulation, such as reading a favorite book, playing games, participation in group activities or a targeted cognitive training. These non-pharmacological cognitive enhancers circumvent the ethical considerations associated with pharmaceutical or technological cognitive enhancement. Meanwhile, they are low in cost and low risk to health and well-being. The effects of exercise and EE on enhancing cognitive function have been attributed to the hippocampal plastic changes including *neurogenesis*, *synaptogenesis* and *angiogenesis*. Some of the cellular mechanisms, such as the increased expression of growth factors, signal pathway, likely underlie each of these changes.

Neural Stem/Progenitor Cells (NSPCs), the major role in *neurogenesis*, are present throughout life in the Subgranular Zone (SGZ) of Dentate Gyrus (DG) and Subventricular Zone (SVZ). Approximately 700 newborn granular neurons are formed every day in the adult human DG [1]. NSPCs in the SGZ differentiate into the granular cells which anchor within the granular layer of the DG establishing synaptic connections with the neighbor neurons to maintain the hippocampal functions. It has been reported that exercise exerts positive effect the adult hippocampal *neurogenesis* in all aspects, including cell proliferation, survival, differentiation and recruitment in the DG [2-5]. Researchers discovered that exposure to EE also showed significantly high differentiation of newborn cells into granular neurons in hippocampus [6,7]. More recently, several reports suggested that the notable EE-induced increase in adult *neurogenesis* was attributed to physical activity associated with exercise [8,9].

## OPEN ACCESS

### \*Correspondence:

Xinhua Zhang, Department of Anatomy,  
Nantong University, Nantong, Jiangsu  
226000, China, Tel: +86-513-8505  
1717;

E-mail: zhangxinhua@ntu.edu.cn

Received Date: 27 Jun 2018

Accepted Date: 11 Jul 2018

Published Date: 18 Jul 2018

### Citation:

Xu N, Zhang L, Zhang X. Exercise and  
Enrichment Recruit Positive Factors for  
Hippocampus Plasticity and Cognitive  
Enhancement. *J Neurosci Cogn Stud.*  
2018; 2(1): 1008.


Copyright © 2018 Xinhua Zhang. This  
is an open access article distributed  
under the Creative Commons Attribution  
License, which permits unrestricted  
use, distribution, and reproduction in  
any medium, provided the original work  
is properly cited.

Formation of new synapses is much critical to store new information in the brain. Granular cells in the DG sprout mossy fibers targeting and forming synaptic connect with neurons in the Cornu Ammonis 3 (CA3) area of the hippocampus, which is an important part of the neural trisynaptic circuit in maintaining cognitive function. It has been reported that exercise significantly increased the levels of synaptic vesicle proteins synaptophysin and synapsin-I indicating the exercise-induced *synaptogenesis* [10-12]. As well, the common exercise running increases the density of dendritic spines and mossy fibre sprouting in granule cells in DG and CA1 pyramidal neurons in hippocampus and layer III pyramidal neurons in entorhinal cortex [13,14].

There are evidences to support the exercise or EE-induced *angiogenesis* in the hippocampus, motor cortex and cerebellum [2,15-18]. It has also been suggested that *angiogenesis* increases circulation which deliver more nutrient metabolites, hormones, growth factors and oxygen to the cognition-related brain regions, and also facilitate metabolic waste disposal, leading to increased cell survival and enhanced *neurogenesis* [19]. Thus, *angiogenesis* could be another key mechanism mediating exercise or EE-induced cognitive enhancement (Figure1).

According to the studies, it is speculated that the molecules such as nutrient metabolites, hormones, growth factors and oxygen in circulation or the local brain region may attribute to the exercise or EE-induced hippocampal plasticity. The following potential trophic factors whose expression can be induced by exercise and EE likely show the critical effects during the plastic processes.

**Brain-Derived Neurotrophic Factor:** Studies have shown that exercise significantly increases peripheral and central levels of Brain-Derived Neurotrophic Factor (BDNF) [20-22], which have been reported to be involved in several functions such as enhancing neuronal survival, *neurogenesis*



**Figure:** The trophic factors mediate exercise and EE-induced cognitive enhancement.

and cognition [21,23]. BDNF administration can induce a form of LTP and improve hippocampus-dependent learning [24-26]. BDNF is able to regulate neuronal growth and survival, which was reported to be involved in the exercise-increased brain's resistance to damage and degeneration [27]. BDNF was found to be robustly involved in *neurogenesis* because the *neurogenesis* was attenuated by BDNF knockdown in the adult DG, but increased in response to exogenous BDNF injection [28,29]. Dendritic growth in adult hippocampal neurons was also decreased after BDNF deletion and increased after BDNF overexpression [30]. The evidences indicate that exercise-induced *synaptogenesis* may be a BDNF-dependent process [10-12].

**Basic Fibroblast Growth Factor:** In the adult CNS, the basic Fibroblast Growth Factor (FGF-2) and its receptors (FGFR) are expressed by astrocytes and neurons located in the SVZ and SGZ, as well other brain regions [31,32]. After birth, FGF-2 is concentrated primarily in the hippocampal subfields CA1-3, and in neurons of the medial septum and the vertical limb of the diagonal band nuclei. Within the mature hippocampus, the CA2 region is the primary area of neuron-derived FGF-2 expression [33], suggesting that FGF-2 may play a role in the development and function of the adult hippocampus. Studies suggested that exercise and EE both increased peripheral and central level of FGF-2, which showed the positive effect on hippocampal *neurogenesis*. For example, loss of FGF-2 in animals caused decrease in adult hippocampal *neurogenesis*. However, these defects could not be rescued by exogenous FGF-2 [34]. Yoshimura et al. reported that hippocampal *neurogenesis* increased in normal adult mice after brain injury, but this phenomenon did not appear in FGF-2 knockout adult mice [35]. These results indicated that endogenous FGF-2 is necessary to stimulate *neurogenesis* in the adult hippocampus. Genetic deletion of FGFR1 resulted in reduced proliferation of hippocampal NSPCs and reduced hippocampal volume during embryonic and postnatal development [36]. These studies suggested that the FGF-2/FGFR system mediates *neurogenesis* in the adult hippocampus. FGF-2 is also a strong pro-angiogenic factor acting as a stimulator of endothelial cell migration, proliferation, sprouting, and tube formation [37,38]. FGF-2 was also proven to be able to enhance hippocampal *synaptogenesis*, including increase of excitatory synapses and synaptic cycling vesicles on hippocampal neurons [39].

**Nerve Growth Factor:** Early studies confirmed that Nerve Growth Factor (NGF) is crucial for neuronal survival and growth, especially the cholinergic neurons, and neurotransmission in both the CNS and peripheral nervous system [40,41]. Recent reports indicated that

continuous NGF infusion promotes proliferation and *synaptogenesis* in the hippocampus and enhanced survival of new neurons in the DG granule cell layer of young adult rats [42,43]. Neurogenic conditions in the hippocampus may be enhanced by the synergistic interactions of NGF and its receptor, TrkA, as well as by NGF-mediated cholinergic regulation. Finally, intracerebroventricular NGF infusion rescued hippocampal *neurogenesis* deficiencies in a transgenic mouse model of Huntington's disease [44], suggesting that the effects of NGF on neuroplasticity likely support the treatment of this disease through exercises or EE.

**Insulin-like Growth Factor 1:** IGF-1, primarily produced in the liver, plays a major role in brain development. It may link the systemic and the central changes induced by exercise. IGF-1 deletion or blockage of its receptors markedly impaired exercise-induced cognitive enhancement that would be ameliorated by exogenous IGF-1 administration [11,45]. Reports showed that IGF-1 directly or indirectly improves the proliferation, survival, and neuronal differentiation of NSPCs, as well prevents their apoptosis in the SGZ of the adult mammalian [46-48]. Zhu and colleagues not only demonstrated its neurotrophic property, but also showed that IGF-1 could enhance neurovascular regeneration in a mouse model of permanent focal cerebral ischemia [49]. The evidences that IGF-1 stimulates an increase in the density of spines in the basal dendrites of CA1 pyramidal neurons while a decrease in the serum IGF-I levels causes a reduction of glutamatergic boutons in the hippocampus [50,51]. This finding suggests that IGF-1 can promote hippocampal synapse formation and/or maintenance.

**Vascular Endothelial Growth Factor:** Peripheral Vascular Endothelial Growth Factor (VEGF) is mainly produced by skeletal muscles. Exercises can increase level of VEGF in skeletal muscle and hippocampus [52]. However, effect of exercise on VEGF expression in the hippocampus is very little. Peripheral VEGF must cross the Blood-Brain Barrier (BBB) to show the functions in the brain. The BBB permeability may increase in response to exercise, providing a potential route for signaling proteins to enter the brain parenchyma from the circulation. VEGF is a strong *angiogenesis* factor, as well shows neurotrophic and neuroprotective effects [53-56].

As the most important mitogen in the process of *angiogenesis*, VEGF plays a role in the angiogenic effects of exercise [57,58]. Pharmacological blockade of *angiogenesis* in the hippocampus impairs spatial learning [59]. The *angiogenesis* is mediated by the binding of VEGF to its receptors on the surface of endothelial cells

through activation of intracellular tyrosine kinases and the multiple downstream signals.

In concert with the *neurogenesis*, VEGF also play an important role in exercise-induced adult hippocampal *neurogenesis* because the increased number of newborn neuronal precursor cells in the hippocampus were not present in adult conditional skeletal myofiber-specific VEGF gene-ablated mice [60,61], suggesting that VEGF expressed by skeletal myofibers may directly or indirectly regulate hippocampal *neurogenesis*. In addition, VEGF secreted in the adult hippocampal NSPCs are known to functionally maintain the neurogenic niche [62]. Specific loss of VEGF in NSPC resulted in impairment of stem cell maintenance although VEGF produced from other cell types was still present [62]. Evidence from knockout mice indicated that hippocampal *neurogenesis* was impaired in VEGF B-KO mice, whereas intraventricular administration of VEGF B restored *neurogenesis* to control levels [63]. These findings suggested that VEGF is involved in *neurogenesis* in the adult hippocampus. Indeed, increasing evidence has shown that VEGF acts as a molecular mediator for adult hippocampal *neurogenesis* and is upregulated by antidepressant treatments including drugs, electroconvulsive seizure [64,65], exercise, and enriched environments [66,67], indicating that VEGF is a promising target for treatment of neural disorders.

In conclusion, the processes of *neurogenesis*, *synaptogenesis* and *angiogenesis* are known as the main mechanisms mediating the exercise or EE-induced cognitive enhancement. The upregulation of these factors may be the critical molecules promoting the above processes. Certainly, it appears that exercise or EE simultaneously stimulates upregulation of several growth factors who would exert the angiogenic, neurogenic and synaptogenic promotion effects at the same time. Keeping exercises and enrichments keep youth.

## Acknowledgment

Grants: 1. The fifth JiangSu "333 program" research project (BRA2016450). 2. Nantong Science and Technology Project.

## References

- Spalding KL, Bergmann O, Alkass K, Bernard S, Salehpour M, Huttner HB, et al. Dynamics of hippocampal neurogenesis in adult humans. *Cell*. 2013;153(6):1219-27.
- Clark PJ, Brzezinska WJ, Puchalski EK, Krone DA, Rhodes JS. Functional analysis of neurovascular adaptations to exercise in the dentate gyrus of young adult mice associated with cognitive gain. *Hippocampus*. 2009;19(10):937-50.
- Speisman RB, Kumar A, Rani A, Foster TC, Ormerod BK. Daily exercise improves memory, stimulates hippocampal neurogenesis and modulates immune and neuroimmune cytokines in aging rats. *Brain Behav Immun*. 2013;28:25-43.
- van Praag H, Christie BR, Sejnowski TJ, Gage FH. Running enhances neurogenesis, learning, and long-term potentiation in mice. *Proc Natl Acad Sci USA*. 1999;96(23):13427-31.
- Praag VH, Kempermann G, Gage FH. Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus. *Nat Neurosci*. 1999;2(3):266-70.
- Kempermann G, Kuhn HG, Gage FH. More hippocampal neurons in adult mice living in an enriched environment. *Nature*. 1997;386(6624):493-5.
- Kempermann G, Gast D, Gage FH. Neuroplasticity in old age: sustained fivefold induction of hippocampal neurogenesis by long-term environmental enrichment. *Ann Neurol*. 2002;52(2):135-43.
- Kobilo T, Liu QR, Gandhi K, Mughal M, Shaham Y, van Praag H. Running is the neurogenic and neurotrophic stimulus in environmental enrichment. *Learn Mem*. 2011;18(9):605-9.
- Mustroph ML, Chen S, Desai SC, Cay EB, DeYoung EK, Rhodes JS. Aerobic exercise is the critical variable in an enriched environment that increases hippocampal neurogenesis and water maze learning in male C57BL/6J mice. *Neuroscience*. 2012;219:62-71.
- Vaynman S, Ying Z, Gómez-Pinilla F. Exercise induces BDNF and synapsin I to specific hippocampal subfields. *J Neurosci Res*. 2004;76(3):356-62.
- Ding Q, Vaynman S, Akhavan M, Ying Z, Gomez-Pinilla F. Insulin-like growth factor I interfaces with brain-derived neurotrophic factor-mediated synaptic plasticity to modulate aspects of exercise-induced cognitive function. *Neuroscience*. 2006;140(3):823-33.
- Quirié A, Hervieu M, Garnier P, Demougeot C, Mossiat C, Bertrand N. Comparative effect of treadmill exercise on mature BDNF production in control versus stroke rats. *PLoS One*. 2012;7(9):e44218.
- Stranahan AM, Khalil D, Gould E. Running induces widespread structural alterations in the hippocampus and entorhinal cortex. *Hippocampus*. 2007;17(11):1017-22.
- Toscano-Silva M, Gomes da Silva S, Scorza FA, Bonvent JJ, Cavalheiro EA, Arida RM. Hippocampal mossy fiber sprouting induced by forced and voluntary physical exercise. *Physiol Behav*. 2010;101(2):302-8.
- Isaacs KR, Anderson BJ, Alcantara AA, Black JE, Greenough WT. Exercise and the brain: angiogenesis in the adult rat cerebellum after vigorous physical activity and motor skill learning. *J Cereb Blood Flow Metab*. 1992;12(1):110-9.
- Swain RA, Harris AB, Wiener EC, Dutka MV, Morris HD, Theien BE, et al. Prolonged exercise induces angiogenesis and increases cerebral blood volume in primary motor cortex of the rat. *Neuroscience*. 2003;117(4):1037-46.
- Van der Borght K, Kóbor-Nyakas DE, Klauke K, Eggen BJ, Nyakas C, Van der Zee EA, et al. Physical exercise leads to rapid adaptations in hippocampal vasculature: temporal dynamics and relationship to cell proliferation and neurogenesis. *Hippocampus*. 2009;19(10):928-36.
- Ekstrand J, Hellsten J, Tingström A. Environmental enrichment, exercise and corticosterone affect endothelial cell proliferation in adult rat hippocampus and prefrontal cortex. *Neurosci Lett*. 2008;442(3):203-7.
- Olson AK, Eadie BD, Ernst C, Christie BR. Environmental enrichment and voluntary exercise massively increase neurogenesis in the adult hippocampus via dissociable pathways. *Hippocampus*. 2006;16(3):250-60.
- Russo-Neustadt A, Beard RC, Cotman CW. Exercise, antidepressant medications, and enhanced brain derived neurotrophic factor expression. *Neuropsychopharmacology*. 1999;21(5):679-82.
- Ding Q, Ying Z, Gómez-Pinilla F. Exercise influences hippocampal plasticity by modulating brain-derived neurotrophic factor processing. *Neuroscience*. 2011;192:773-80.
- Griffin ÉW, Mullally S, Foley C, Warmington SA, O'Mara SM, Kelly AM. Aerobic exercise improves hippocampal function and increases BDNF in the serum of young adult males. *Physiol Behav*. 2011;104(5):934-41.
- Zigova T, Pencea V, Wiegand SJ, Luskin MB. Intraventricular administration of BDNF increases the number of newly generated neurons in the adult olfactory bulb. *Mol Cell Neurosci*. 1998;11(4):234-45.
- Bramham CR, Panja D. BDNF regulation of synaptic structure, function, and plasticity. *Neuropharmacology*. 2014;76:601-2.
- Griffin EW, Bechara RG, Birch AM, Kelly AM. Exercise enhances hippocampal-dependent learning in the rat: evidence for a BDNF-related mechanism. *Hippocampus*. 2009;19(10):973-80.
- Bechara RG, Lyne R, Kelly ÁM. BDNF-stimulated intracellular signalling mechanisms underlie exercise-induced improvement in spatial memory in

- the male Wistar rat. *Behav Brain Res.* 2014;275:297-306.
27. Neeper SA, Gómez-Pinilla F, Choi J, Cotman CW. Physical activity increases mRNA for brain-derived neurotrophic factor and nerve growth factor in rat brain. *Brain Res.* 1996;726(1-2):49-56.
  28. Taliat D, Stall N, Dar DE, Zangen A. Knockdown of brain-derived neurotrophic factor in specific brain sites precipitates behaviors associated with depression and reduces neurogenesis. *Mol Psychiatry.* 2010;15(1):80-92.
  29. Scharfman H, Goodman J, Macleod A, Phani S, Antonelli C, Croll S. Increased neurogenesis and the ectopic granule cells after intrahippocampal BDNF infusion in adult rats. *Exp Neurol.* 2005;192(2):348-56.
  30. Wang L, Chang X, She L, Xu D, Huang W, Poo MM. Autocrine action of BDNF on dendrite development of adult-born hippocampal neurons. *J Neurosci.* 2015;35(22):8384-93.
  31. Chadashvili T, Peterson DA. Cytoarchitecture of fibroblast growth factor receptor 2 (FGFR-2) immunoreactivity in astrocytes of neurogenic and non-neurogenic regions of the young adult and aged rat brain. *J Comp Neurol.* 2006;498(1):1-15.
  32. Weickert CS, Kittell DA, Saunders RC, Herman MM, Horlick RA, Kleinman JE, et al. Basic fibroblast growth factor and fibroblast growth factor receptor-1 in the human hippocampal formation. *Neuroscience.* 2005; 131(1):219-33.
  33. Gómez-Pinilla F, Lee JW, Cotman CW. Distribution of basic fibroblast growth factor in the developing rat brain. *Neuroscience.* 1994;61(4):911-23.
  34. Werner S, Unsicker K, von Bohlen und Halbach O. Fibroblast growth factor-2 deficiency causes defects in adult hippocampal neurogenesis, which are not rescued by exogenous fibroblast growth factor-2. *J Neurosci Res.* 2011;89(10):1605-17.
  35. Yoshimura S, Takagi Y, Harada J, Teramoto T, Thomas SS, Waeber C, et al. FGF-2 regulation of neurogenesis in adult hippocampus after brain injury. *Proc Natl Acad Sci USA.* 2001;98(10):5874-9.
  36. Ohkubo Y, Uchida AO, Shin D, Partanen J, Vaccarino FM. Fibroblast growth factor receptor 1 is required for the proliferation of hippocampal progenitor cells and for hippocampal growth in mouse. *J Neurosci.* 2004; 24(27):6057-69.
  37. Folkman J, Klagsbrun M. Angiogenic factors. *Science.* 1987;235(4787):442-7.
  38. Kent KC, Mii S, Harrington EO, Chang JD, Mallette S, Ware JA. Requirement for protein kinase C activation in basic fibroblast growth factor-induced human endothelial cell proliferation. *Circ Res.* 1995;77(2):231-8.
  39. Li AJ, Suzuki S, Suzuki M, Mizukoshi E, Imamura T. Fibroblast growth factor-2 increases functional excitatory synapses on hippocampal neurons. *Eur J Neurosci.* 2002;16(7):1313-24.
  40. Moises HC, Womble MD, Washburn MS, Williams LR. Nerve growth factor facilitates cholinergic neurotransmission between nucleus basalis and the amygdala in rat: an electrophysiological analysis. *J Neurosci.* 1995;15(12):8131-42.
  41. Cooper JD, Salehi A, Delcroix JD, Howe CL, Belichenko PV, Chua-Couzens J, et al. Failed retrograde transport of NGF in a mouse model of Down's syndrome: reversal of cholinergic neurodegenerative phenotypes following NGF infusion. *Proc Natl Acad Sci USA.* 2001;98(18):10439-44.
  42. Frielingsdorf H, Simpson DR, Thal LJ, Pizzo DP. Nerve growth factor promotes survival of new neurons in the adult hippocampus. *Neurobiol Dis.* 2007;26(1):47-55.
  43. Birch AM, Kelly ÁM. Chronic intracerebroventricular infusion of nerve growth factor improves recognition memory in the rat. *Neuropharmacology.* 2013;75:255-61.
  44. Zhang H, Petit GH, Gaughwin PM, Hansen C, Ranganathan S, Zuo X, et al. NGF rescues hippocampal cholinergic neuronal markers, restores neurogenesis, and improves the spatial working memory in a mouse model of Huntington's Disease. *J Huntingtons Dis.* 2013;2(1):69-82.
  45. Trejo JL, Llorens-Martín MV, Torres-Alemán I. Torres-Aleman, The effects of exercise on spatial learning and anxiety-like behavior are mediated by an IGF-I-dependent mechanism related to hippocampal neurogenesis. *Mol Cell Neurosci.* 2008;37(2):402-11.
  46. Grote HE, Hannan AJ. Regulators of adult neurogenesis in the healthy and diseased brain. *Clin Exp Pharmacol Physiol.* 2007;34(5-6):533-45.
  47. Zygar CA, Colbert S, Yang D, Fernald RD. IGF-1 produced by cone photoreceptors regulates rod progenitor proliferation in the teleost retina. *Brain Res Dev Brain Res.* 2005;154(1):91-100.
  48. Kouroupi G, Lavdas AA, Gaitanou M, Thomaidou D, Stylianopoulou F, Matsas R. Lentivirus-mediated expression of insulin-like growth factor-I promotes neural stem/precursor cell proliferation and enhances their potential to generate neurons. *J Neurochem.* 2010;115(2):460-74.
  49. Zhu W, Fan Y, Hao Q, Shen F, Hashimoto T, Yang GY, et al. Postischemic IGF-1 gene transfer promotes neurovascular regeneration after experimental stroke. *J Cereb Blood Flow Metab.* 2009;29(9):1528-37.
  50. Glasper ER, Llorens-Martin MV, Leuner B, Gould E, Trejo JL. Blockade of insulin-like growth factor-I has complex effects on structural plasticity in the hippocampus. *Hippocampus.* 2010;20(6):706-12.
  51. Trejo JL, Piriz J, Llorens-Martin MV, Fernandez AM, Bolós M, LeRoith D, et al. Central actions of liver-derived insulin-like growth factor I underlying its pro-cognitive effects. *Mol Psychiatry.* 2007;12(12):1118-28.
  52. Tang K, Xia FC, Wagner PD, Breen EC. Exercise-induced VEGF transcriptional activation in brain, lung and skeletal muscle. *Respir Physiol Neurobiol.* 2010;170(1):16-22.
  53. Sondell M, Lundborg G, Kanje M. Vascular endothelial growth factor has neurotrophic activity and stimulates axonal outgrowth, enhancing cell survival and Schwann cell proliferation in the peripheral nervous system. *J Neurosci.* 1999;19(14):5731-40.
  54. Jin KL, Mao XO, Greenberg DA. Vascular endothelial growth factor: direct neuroprotective effect in in vitro ischemia. *Proc Natl Acad Sci USA.* 2000;97(18):10242-7.
  55. Jin K, Zhu Y, Sun Y, Mao XO, Xie L, Greenberg DA. Vascular endothelial growth factor (VEGF) stimulates neurogenesis in vitro and in vivo. *Proc Natl Acad Sci USA.* 2002;99(18):11946-50.
  56. Matsuzaki H, Tamatani M, Yamaguchi A, Namikawa K, Kiyama H, Vitek MP, et al. Vascular endothelial growth factor rescues hippocampal neurons from glutamate-induced toxicity: signal transduction cascades. *FASEB J.* 2001;15(7):1218-20.
  57. Ferrara N. VEGF-A: a critical regulator of blood vessel growth. *Eur Cytokine Netw.* 2009;20(4):158-63.
  58. van Praag H. Exercise and the brain: something to chew on. *Trends Neurosci.* 2009;32(5):283-90.
  59. Kerr AL, Steuer EL, Pochtarev V, Swain RA. Angiogenesis but not neurogenesis is critical for normal learning and memory acquisition. *Neuroscience.* 2010;171(1):214-26.
  60. Rich B, Scadeng M, Yamaguchi M, Wagner PD, Breen EC. Skeletal myofiber vascular endothelial growth factor is required for the exercise training-induced increase in dentate gyrus neuronal precursor cells. *J Physiol.* 2017;595(17):5931-43.
  61. Ballard HJ. Exercise makes your brain bigger: skeletal muscle VEGF and hippocampal neurogenesis. *J Physiol.* 2017;595(17):5721-2.
  62. Kirby ED, Kuwahara AA, Messer RL, Wyss-Coray T. Adult hippocampal

- neural stem and progenitor cells regulate the neurogenic niche by secreting VEGF. *Proc Natl Acad Sci USA*. 2015;112(13):4128-33.
63. Sun Y, Jin K, Childs JT, Xie L, Mao XO, Greenberg DA. Vascular endothelial growth factor-B (VEGFB) stimulates neurogenesis: evidence from knockout mice and growth factor administration. *Dev Biol*. 2006;289(2): 329-35.
64. Segi-Nishida E, Warner-Schmidt JL, Duman RS. Electroconvulsive seizure and VEGF increase the proliferation of neural stem-like cells in rat hippocampus. *Proc Natl Acad Sci USA*. 2008;105(32):11352-7.
65. Fournier NM, Duman RS. Role of vascular endothelial growth factor in adult hippocampal neurogenesis: implications for the pathophysiology and treatment of depression. *Behav Brain Res*. 2012;227(2):440-9.
66. Fabel K, Fabel K, Tam B, Kaufer D, Baiker A, Simmons N, et al. VEGF is necessary for exercise-induced adult hippocampal neurogenesis. *Eur J Neurosci*. 2003;18(10):2803-12.
67. Cao L, Jiao X, Zuzga DS, Liu Y, Fong DM, Young D, et al. VEGF links hippocampal activity with neurogenesis, learning and memory. *Nat Genet*. 2004;36(8):827-35.