



Fennel (*Foeniculum vulgare* Mill.): A Common Spice with Unique Medicinal Properties

Shahid Akbar*

Department of Pharmacology, Qassim University, Saudi Arabia

Abstract

Introduction: *Foeniculum vulgare* is a perennial, aromatic plant belonging to Apiaceae (Umbelliferae) family. The fruits, commonly referred to as seeds, are ridged, aromatic and oblong or ellipsoid shaped. Originally indigenous to the Mediterranean region, it has now been naturalized and cultivated throughout the world and is universally known as fennel and by more than 100 other names. The dried fruits and aerial parts have widely been used since ancient times in culinary and traditional folk medicines in many cultures around the world. It was well-known to the ancient Egyptians, Greeks, Romans and Chinese. Hippocrates and Dioscorides described it as a diuretic and Emmenagogue and to strengthen eyesight. The fruits are reputed as carminative, stomachic, diuretic, Emmenagogue and Galactagogue and to promote menstruation and facilitate birth.

Photochemical and Pharmacological Studies: Fennel fruits contain carbohydrates, alkaloids, Phytosterols, phenols, tannins and flavonoids. They are also a rich source of dietary fiber, protein, calcium, iron, magnesium and manganese. Pharmacologically, the fruits possess antioxidant, anti-inflammatory, antispasmodic, diuretic, antihypertensive, antimicrobial, Gastro protective, estrogenic, Hepatoprotective and antithrombotic activities.

Discussion and Conclusion: This article reviews the use of fennel in medical practices in various parts of the world and reflects upon the scientific studies conducted and how they reconcile with its traditional uses. The clinical studies in small number of patients revealed several clinical effects but are not sufficiently wide ranging and have not been reproduced in numbers that would make them conclusive. Randomized, double-blinded, placebo-controlled clinical trials would establish the clinical usefulness of fennel in a scientific manner.

Keywords: Fennel; *Foeniculum Vulgare*; Anethole; Fenchone; Estragole

OPEN ACCESS

*Correspondence:

Shahid Akbar, Department of
Pharmacology, Qassim University,
Saudi Arabia,
E-mail: Drakbarmdp@d@gmail.Com,
Shahidakbar@sbcglobal.net

Received Date: 22 Sep 2018

Accepted Date: 05 Nov 2018

Published Date: 07 Nov 2018

Citation:

Akbar S. Fennel (*Foeniculum vulgare*
Mill.): A Common Spice with Unique
Medicinal Properties. *Ann Complement*
Altern Med. 2018; 1(1): 1001.

Copyright © 2018 Shahid Akbar. This
is an open access article distributed
under the Creative Commons Attribution
License, which permits unrestricted
use, distribution, and reproduction in
any medium, provided the original work
is properly cited.

Introduction

Historical perspective and medicinal uses

Foeniculum Vulgare is usually a perennial, aromatic plant belonging to Apiaceae (Umbelliferae) family with many subspecies and varieties. *F. vulgare* subsp. *vulgare* var. *Dulce* is called sweet fennel, while *F. vulgare* mill. Subsp. *vulgare* var. *vulgare* is referred to as bitter fennel. Mostly these two varieties are commercially available and are used medicinally. The plant grows wild in Mediterranean region and in European countries of temperate climate but is now cultivated in most parts of the world for commercial purposes. The fruits commonly referred to as seeds are ridged, oblong or ellipsoid shaped, aromatic and are universally known as fennel and by more than 100 other names throughout the world and have been used medicinally since ancient times as one of the ancient Saxon people's nine sacred herbs, fennel was credited with the power to cure and was valued as a magic herb [1,2]. In the middle ages, it was draped over doorways on midsummer's eve to protect the household from evil spirits [3]. Fennel is also one of the most frequently quoted plants in the chilandar medical codex, the best preserved medieval Serbian manuscript on European medical science from the 12th to 15th centuries [4]. Famous Greek physicians, Hippocrates and Dioscorides mentioned fennel as a diuretic and Emmenagogue and its juice was supposed to sharpen the eyesight [5]. The fruits are especially reputed to increase milk secretion, promote menstruation, facilitate birth, alleviate the symptoms of the male climacteric and increase libido [6]. In Indian traditional medicines, it is considered a stimulant, carminative, aromatic, stomachic, Emmenagogue and Galactagogue; the fennel fruit water is traditionally used to relieve flatulent colic and also as a diuretic and diaphoretic in children and infants. A hot infusion is used to treat amenorrhea and to improve lactation, [5,7-12] to improve eyesight and to open liver and spleen obstructions [7]. In

Europe and Mediterranean countries, fennel is traditionally used as antispasmodic, diuretic, anti-inflammatory, analgesic, secret motor, Galactagogue, eye lotion and antioxidant remedy; topically, fennel powder is used as a poultice for snake bites and fennel infusions are used in Europe for nursing babies to prevent flatulence and colic spasm [3]. It is approved in Europe for use in GI disorders in children above the age of 4 years since November, 2005 by the committee on herbal medicinal products (hmpc) of the European medicines agency, but is not recommended for use for more than two weeks [13]. It is one of the most commonly consumed herbs by more than a quarter of Italian pregnant women every day for at least 3 months during pregnancy [14]. A subspecies of fennel, *F. vulgare* subsp. *piperitum*, is used for mouth ulcers in the Basilicata region of southern Italy in Portugal, it is highly recommended for treatment of diabetes, bronchitis and chronic coughs, and for kidney stones [15,16]. In Mexican traditional medicine, a decoction is used as a galactagogue and to treat tuberculosis and other respiratory diseases [17,18]. In Palestine, wildy grown fennel is traditionally used as a regular part of diet and as a popular digestive stimulant in north-east Lebanon [19,20]. In Chinese medicine, it is regarded carminative, stomachic and stimulant, and is used to relieve chills, abdominal distension, vomiting and diarrhea [21,22]. Fennel leaves infusion is also used to treat infant's stomach-ache and the seeds are used by adults to dispel gases in Guyana and Surinam and boiled or roasted roots are used for the treatment of gonorrhoea in east Africa [23,24]. Fennel essential oil is also widely used as a flavoring agent in products such as liqueurs, bread, cheese and as an ingredient of cosmetics and pharmaceutical products [3,25-29].

Photo constituents

Fennel fruits contain alkaloids, carbohydrates, Phytosterols, phenols, tannins, coumarins and flavonoids as nonvolatile substances; the acetone extract is richer in phenols, while methanol extract contains higher amounts of flavonoids [30]. Kaur et al. [31] reported the presence of alkaloids, flavonoids, tannins, saponins and trace amounts of cardiac glycosides in hot water fruit extract and methanol extract of fruit sample from Egypt was reported to contain flavonoids, terpenoids, alkaloids, phenols and sterols; which had estragole (methyl chavicol) (71.1%) as the predominant alcohol, gallic acid (18.9%) as the major phenolic compound and l-limonene (11.9%) as the most prevalent monoterpene hydrocarbon [32]. Major phenolic compounds identified in fennel include 3-o-caffeoylquinic acid, chlorogenic acid, 4-o-caffeoylquinic acid, eriocitrin, rutin, miquelianin, 1,3-o-dicaffeoylquinic acid, 1,5-o-dicaffeoylquinic acid, 1,4-o-dicaffeoylquinic acid and rosmarinic acid; [33] identified flavone (OL)-o-glycosides are quercetin 3-glucuronide, isoquercitrin, rutin, and quercetin 3-arabinoside; other phenols reported are kaempferol 3-glucuronide and kaempferol 3-arabinoside [34]. Bergapten, columbianetin, osthonol, psoralen, scoparone, seselin, vanillin, beta-sitosterol and stigmasterol have also been identified in fruits [35]. Two diglucoside stilbene trimers and a benzoisofuranone derivative have also been isolated from the fruits [36]. The elements present in fennel fruits from Ethiopia were reported as calcium, magnesium, iron, manganese, copper, chromium, cobalt, zinc, nickel and cadmium [2]. Fennel is reported as one of the plant sources with highest amounts of calcium, potassium, sodium and phosphorus [1].

The method of distillation significantly affects the yield and qualitative composition of the essential oil [37]. Essential oil composition also varies depending on the maturation stages of the

plant [38]. The yield of Turkish essential oil (5.0 ml/kg) and content of trans-anethole are very low (34.8%), whereas yield of essential oil is maximum in fennel from Norway and Austria (50.7 ml/kg and 50.5 ml/kg), respectively; these samples are richer in Fenchone (21.2% and 22.8%, respectively), but contain less trans-anethole (64.6% to 63.7%) than samples from Estonia and Moldova (82.0% and 80.9%) [39]. In fennel samples collected from the wild population in the center and south of Portugal, the yields of essential oils varied greatly from 1.1% to 2.9%, and the main constituents, trans-anethole (7.9% to 77.7%), Fenchone (16.9% to 34.7%) and estragole (2.5% to 66.0%) also showed great variations [40]. In general, fennel oil extracted by either distillation-extraction or supercritical fluid extraction shows similar compositions, with trans-anethole, estragole and Fenchone as the main components [37,41-42]. Trans-anethole (85.63%) is generally the predominant constituent of the oil [43-47], while estragole is found in small amounts (2.87%), and the quantity of Fenchone is <1% [45]. Trans-anethole (69.8%) and limonene (22.5%), though, were identified as the major constituents of essential oil in fennel samples cultivated in southeastern Brazil [48] and Miguel et al. [49] reported estragole as the dominant constituent in the fruit essential oil of samples from Portugal, and trans-anethole, alpha-pinene and limonene being the main components of dried aerial parts essential oil. In samples of essential oil of fennel grown under different climatic conditions in Romania, major compounds identified in all samples were trans-anethole, estragole, fenchone, limonene, alpha-pinene and gamma-terpinene [50] and fennel oil samples from Egypt also showed trans-anethole, estragole, fenchone and limonene as the major constituents [51]. Essential oils obtained from various wild Italian varieties contained five chemical groups characterized by (i) alpha-phell andrene, methyl chavicol (estragole) and trans-anethole; (ii) alpha-pinene, limonene and trans-anethole; (iii) methyl chavicol and alpha-phell andrene; (iv) methyl chavicol and alpha-pinene; and (v) alpha-phell andrene [52].

In Chinese medicine various frying methods are used before the fruits are incorporated in poly herbal preparations. After different frying methods, contents of all twenty-four ingredients of the volatile oil from these fruits were changed, and eighteen new compounds, including Linalyl acetate, farnesene, p-allylphenyl aromatic oxide, and Menthone and hexyl octanoate were created; however, trans-anethole remained the largest of the effective ingredients in the fried samples [53].

Estragole, an isomer of anethole, is regarded carcinogenic and genotoxic by the hmpc of European medicine agency. Estragole is rapidly metabolized in humans and excreted in urine as 1'-hydroxyestragole [54] and the hmpc recommends that the total human daily intake of estragole from all food sources should not exceed 0.07 mg/kg [3]. Nevertheless, contents of estragole in commercial teabags and loose fruits vary widely. For example, estragole contents in teas made from teabags ranged from 241 to 2,058 µg/l, while in diluted instant teas they were from 9 to 912 µg/l, estragole contents ranged from 251 to 1718 µg/l in unpackaged fruits (seeds) [55] a study among finish consumers, though, found daily consumption of estragole due to fennel containing plant food supplements to be moderate and not detrimental to health [56].

Results

Experimental studies

Antioxidant effects: Aqueous fruit extract (AE) demonstrates

direct and highly significant *in vitro* nitric oxide scavenging [57] and antioxidant activities [58-59] and improves activities of antioxidant enzymes in trichloroacetic acid-exposed rats [60]. The quantity of AE required for 50% inhibition of hydroxyl radicals was 700 microg compared to 4500 microg for ascorbic acid; whereas, scavenging of superoxide radicals by 50% required 205 microg fennels vs. 260 microg ascorbic acid [59]. Although aqueous, methanol acetone and other extracts of fruits have all been reported to possess antioxidant activity, the methanol and acetone extracts are comparable to activity of butyrate hydroxyl toluene because acetone fruit extract is reported to have comparatively high amount of total phenolics, whereas methanol extract had highest amount of total flavonoids [30,32,61]. Moderate antioxidant activity of ethanol fruit extract was, however, independent of the total flavonoid content present in the extract [62]. Four coumarins, isolated from the methanol extract, have shown excellent *in vitro* antioxidant and anti-inflammatory activities [63]. Both methanol extract and essential oil also significantly improved activities of antioxidant enzymes and inhibited lipid peroxidation [64-66]. The essential oil with predominant trans-anethole content showed a significantly higher antioxidant activity with low IC50 value [67] and demonstrated *in vitro* antioxidant capacities comparable to that of alpha-tocopherol and butyrate hydroxyl toluene [68]. Essential oil is, however, rapidly auto-oxidized by light, and the rate of its oxidation is reported to be slower in the dark [69]. The shoots have the highest radical-scavenging activity and lipid peroxidation inhibition capacity, due to highest content of phenolics and ascorbic acid, and high concentration of tocopherol and allegedly are the only part to contain flavonoids, but Goswami et al. [30] reported the presence of flavonoids in fruits too [30]. In early fruiting stage, aerial parts contain lowest flavonoid contents and higher phenolic acid content and aerial parts infusion rather than decoction shows better *in vitro* antioxidant activity [70,71]. The length of storage period of aerial parts, however, significantly affects the antioxidant content and antioxidant activity [72].

Antimicrobial effects: Significant *in vitro* antibacterial activity of hot aqueous and acetone fruit extracts against *Enterococcus faecalis*, *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Salmonella typhimurium* and *Shigella flexneri* has been reported. The ambient temperature water extract showed weaker antibacterial activity, whereas the activity was completely lost in the boiling water extract and the acetone extract was active with significantly lower Minimum Inhibitory Concentration (MIC) values than the water extract [31]. While a number of reports indicated that the essential oil exhibits insignificant or moderate antibacterial activity against standard strains of *E. coli*, *S. aureus*, *P. aeruginosa*, *Bacillus subtilis*, *Bacillus megaterium*, and *Bacillus cereus* [51,73-75]; significant *in vitro* activity of essential oil against various strains of *S. aureus* and *E. coli* [40,50,76], against *Salmonella enterica*, *Listeria monocytogenes*, *B. cereus*, *Candida albicans*, *Aspergillus niger*, and *Penicillium spp* and synergistic activity with amoxicillin or tetracycline against *E. coli*, *Sarcina lutea* and *B. subtilis* strains have been reported [40,50,76]. The essential oil also exhibited *in vitro* antifungal activity against *C. albicans*, [50-51,77] aspergillus species [51,78] and potent activity against dermatophytes *Trichophyton rubrum*, *T. tonsurans*, *T. mentagrophytes* and *Micro sporum gypseum*, better than the standard antifungal agents, fluconazole and amphotericin b[79]. Jazani et al. [80] reported the essential oil bactericidal against isolates of multi-drug resistant *Acinetobacter baumannii*, a gram negative coccobacillus, increasingly responsible for nosocomial infections.

Essential oil of Turkish fennel was also reported effective against *E. coli*, *L. monocytogenes*, *S. typhimurium*, and *S. aureus*, [43] *E. faecalis*, *Staphylococcusepidermidis* and *S. aureus*; against *Morganella morganii*, *Proteus mirabilis*, *Salmonella enteritidis*, and *P. aeruginosa* [40], Strains of klebsiella Pneumonia producing Extended-Spectrum Beta-Lactamase (ESBL) enzyme [81]. Moderate activity of methanol extract against 15 strains of *Helicobacter pylori* with an MIC of 50 mcg/ml [82], significant activity of hydro-ethanol extract against *campylobacter jejuni* [83] and against the multidrug resistant *Mycobacterium tuberculosis* has also been reported [18]. Fennel oil, used at sub-inhibitory concentrations, also decreased the expression of *S. aureus* exotoxins, including α -toxin, *Staphylococcal* enterotoxins and toxic shock syndrome toxin 1 [47].

Gastro protective effect: Pretreatment with fennel water extract significantly reduced ethanol-induced gastric damage in rats, which was suggested to be due to reduction in lipid peroxidation and augmentation in the antioxidant activity [84]. The water extract is reported to increase gastric acid secretion by more than three times of basal secretion in rats that was not blocked by atropine; but significantly reduced basal acid secretion in aspirin-induced gastric mucosal damage [85]. Oral administration of essential oil and anethole is also significantly protective against ethanol-induced gastric lesions in rats [86]. The antioxidant activity of phenol contents is partly credited for the anti-ulcer effects [84].

Anti-inflammatory effect: Methanol fruit extract exhibited inhibitory effects against both acute and sub-acute inflammation and a central analgesic effect, and significantly increased plasma antioxidant enzymes activities and the HDL-cholesterol level [64]. Four coumarins isolated from the fruits showed excellent *in vitro* antioxidant activity and imperatorin had the greatest anti-inflammatory and antioxidant activities, inhibiting pro-inflammatory cytokines production, including interleukin-6 and tumor necrosis factor- α (TNF- α) in Lipopolysaccharides (LPS)-stimulated raw 264.7 cells [62]. Essential oils from the aerial parts and fruits are also reported to inhibit 5-lipoxygenase enzyme [49].

Estrogenic effects: A number of experimental studies point to the estrogenic effects of fennel. Water extract shows renoprotective effects in experimental polycystic ovary syndrome in rats [87] and prevents ovariectomy-induced bone loss, reducing both osteoclast differentiation and function [88]. Administration of acetone fruit extract also exhibited oestrogenic activity [89] increasing weight of mammary glands [89] oviducts of ovariectomized rats [90] vaginal cornification and oestrus cycle in female rats, and significantly reducing protein contents of testes and vas deferens in male rats [89].

Cardiovascular effects: It is one of the commonly used plants to lower blood pressure that acts by causing diuresis, increasing excretion of sodium and water from the body [91]. Oral fruit water extract lowered systolic blood pressure of spontaneously hypertensive rats and increased water, sodium and potassium excretion [92]. Intravenous administration of boiled water extract of leaves also significantly lowered arterial blood pressure, whereas the non-boiled water extract showed insignificant hypotensive activity [93]. An increase in both RBC and WBC counts of rats was observed after alternate day oral administration of the hydro alcohol extract for 30 days [94]. Oral administration of essential oil and anethole to mice produced a significant antithrombotic activity and both inhibited platelet aggregation, and prevented thrombin-induced clot retraction [86,95]. Co-administration of fennel essential oil with Emamectin

Table: Selected Pharmacological and Clinical Observations of Fennel (*Foeniculum vulgare*).

S. No.	Study Type	Study Year	Part Used	Extract Type/Essential oil	Study Subjects	Reference No.
Experimental						
1	Antioxidant	2004, 2008, 2009, 2011, 2014	Fruits	Water, Methanol, Acetone, and Ethanol Extracts	<i>In Vitro</i>	[30,32,58,59,61,69]
2	Antioxidant	2008	Fruits	Water Infusion	Rats	[60]
3	Antioxidant	2000, 2011	Fruits	Essential oil	<i>In Vitro</i>	[66,67]
4	Antioxidant	2009	Aerial Parts		<i>In Vitro</i>	[70]
5	Antioxidant	2009	Fruits	Coumarins	<i>In Vitro</i>	[63]
6	Antioxidant	2015	Fruits	Ethanol Extract	<i>In Vitro</i>	[62]
7	Nitric Oxide Scavenging	2003	Fruits	Water Extract	<i>In Vitro</i>	[57]
8	Antimicrobial	2009	Fruits	Hot Water and Acetone Extracts	<i>E. faecalis</i> , <i>S. aureus</i> , <i>E. coli</i> , <i>P. aeruginosa</i> , <i>S. typhi</i> , <i>S. typhimurium</i> , <i>S. flexneri</i>	[31]
9	Antimicrobial	2015, 2004, 2008, 2011, 2002, 2002, 2007, 2010, 2010, 2015, 2009, 2011	Fruits	Essential oil	<i>E. coli</i> , <i>S. aureus</i> , <i>P. aeruginosa</i> , <i>B. subtilis</i> , <i>B. megaterium</i> , <i>B. cereus</i> , <i>S. enterica</i> , <i>L. monocytogenes</i> , <i>C. albicans</i> , <i>A. niger</i> , <i>T. rubrum</i> , <i>T. tonsurans</i> , <i>T. mentagrophytes</i> <i>M. gypseum</i> , <i>A. baumannii</i> , <i>S. typhimurium</i> , <i>E. faecalis</i> , <i>S. epidermidis</i> <i>Morganella morganii</i> , <i>P. mirabilis</i> , <i>S. enteritidis</i> , strains of <i>K. pneumonia</i>	[40,43,50,51,73-76,78-81]
10	Antimicrobial	2005	Fruits	Methanol Extract	15 strains of <i>H. pylori</i>	[82]
11	Antimicrobial	2008, 2010	Fruits	Hydro-ethanol extract	<i>C. jejuni</i> , MDR <i>M. tuberculosis</i>	[18,83]
12	Antispasmodic	2003	Fruits	Oil emulsion	Infants	[108]
13	Gastric Antiulcer	2000, 2007	Fruits	Water Extract	Rats	[84,85]
14	Gastric Antiulcer	2007	Fruits	Essential oil	Rats	[86]
15	Anti inflammatory	2004	Fruits	Methanol Extract		[64]
16	Antihypertensive	2001	Fruits	Water Extract	SHR	[92]
17	Antihypertensive	1998	Leaves	Water Extract (I.V.)	Rats	[93]
18	Antithrombotic	2007, 2006	Fruits	Essential oil	Mice, Rats	[86,95]
Clinical						
19	Analgesic	2013, 2006, 2003, 2012	Fruits	Extract or Essence	Young Women with Primary Dysmenorrhea	[108-111]
20	Antispasmodic	2003	Fruits	Fennel Oil Emulsion	Infants (2 to 12 weeks old)	[106]
21	Anorexogenic	2015	Fruits	Tea	Overweight Women	[107]
22	Vaginal Symptoms	2016	Fruits	Vaginal Cream	Postmenopausal Women	[112]
23	Idiopathic Hirsutism	2003, 2014	Fruits	Cream and Gel	Women	[113,114]

Abbreviations: *A. baumannii*: *Acinetobacter baumannii*; *A. niger*: *Aspergillus niger*; *B. cereus*: *Bacillus cereus*; *B. megaterium*: *Bacillus megaterium*; *epidermidis*; *B. subtilis*: *Bacillus subtilis*; *C. jejuni*: *Campylobacter jejuni*; *C. albicans*: *Candida albicans*; *E. coli*: *Escherichia coli*; *E. faecalis*: *Enterococcus faecalis*; *H. pylori*: *Helicobacter pylori*; *K. pneumonia*: *Klebsiella pneumonia*; *L. monocytogenes*: *Listeria monocytogenes*; *M. gypseum*: *Microsporium gypseum*; *M. morganii*: *Morganella morganii*; *M. tuberculosis*: *Mycobacterium tuberculosis*; *P. aeruginosa*: *Pseudomonas aeruginosa*; *P. mirabilis*: *Proteus mirabilis*; *S. aureus*: *Staphylococcus aureus*; *S. epidermidis*: *Staphylococcus epidermidis*; *S. typhimurium*: *Salmonella typhimurium*; *S. enterica*: *Salmonella enterica*; *S. enteritidis*: *Salmonella enteritidis*; *S. typhi*: *Salmonella typhi*; *S. flexneri*: *Shigella flexneri*; *T. mentagrophytes*: *Trichophyton mentagrophytes*; *T. tonsurans*: *Trichophyton tonsurans*; *T. r* *Trichophyton mentagrophytes ubrum*: *Trichophyton rubrum*

Benzoate (EB), an avermectin insecticide used extensively in pest control on vegetable and field crops, to rats ameliorated EB-induced coagulative necrosis and blood vessels congestion of the liver and necrosis of the white pulp of the spleen [96]. Topical application of water fruit extract also significantly reduced intraocular pressure in both normotensive and experimental glaucoma model in rabbits [97].

Miscellaneous effects: While Joshi et al. [98] reported that administration of methanol extract of the whole plant for eight successive days ameliorated scopolamine-induced amnesic effect, aging-induced memory deficits, and significantly inhibited Acetylcholinesterase (AChE) in Mice et al. [99] reported significant ache and Butyrylcholinesterase inhibitory activity of the essential oil, more than any single active component, Mata et al. [100] and Aazza et al. [66] reported poor to moderate *in vitro* ache inhibitory activity

of fennel oil. The essential oil of aerial parts also exhibited potential anxiolytic activity in mice [101]. Ethanol extract, and essential oil also significantly produce *in vitro* relaxant effect on methacholine-induced contraction of tracheal chains; calcium channels inhibition was not a contributing mechanism of the relaxant effect [102]. Fennel supplementation in diet protected rats against cyclosporine-nephrotoxicity [103] and the essential oil potently protected against ccl4-hepatotoxicity in rats [104]. The fruit extract was also reported to exhibit *in vitro* immuno-modulatory Nf kappaB activities [105].

Clinical studies: Results of relevant clinical studies on fennel have been presented in table 1. A number of clinical studies have shown significant antispasmodic, analgesic, and putative estrogenic activities of fennel. In a Randomized, Placebo-Controlled Trial (RCT), use of fennel oil emulsion relieved colic in 65% of 2 to 12 weeks

old infants, compared to 23.7% infants treated with placebo [106]. Drinking fennel tea followed by buffet lunch in overweight Korean women significantly decreased appetite and food consumption, and increased feeling of fullness in a placebo-controlled, single-blinded, randomized, crossover study [107]. Several clinical studies in young high-school Iranian girls with moderate to severe dysmenorrhea showed that treatment with fennel extract or essence of fennel's fruit effectively relieved menstrual pain, comparable to mefenamic acid [108-111]. However, patients complained of the odor and unpleasant fennel taste and 16% patients withdrew from one study for this reason [110]. In a double-blind RCT, once daily intravaginal application of fennel 5% vaginal cream in postmenopausal women for 8 weeks, significantly improved postmenopausal vaginal symptoms, increasing number of superficial cells and significantly decreasing the vaginal pH [112]. In another double-blind RCT in women with mild to moderate idiopathic hirsutism, topical application of 3% fennel gel [113] or application of cream containing 2% ethanol fruit extract significantly reduced hair thickness after 24 weeks of treatment compared to placebo [114]. Topical application of 5% and 8% fennel oil-containing aerosol and cream produced 84% and 70% repellency against mosquitoes after 90 min of exposure, respectively [115]. An interesting observation reported was that inhalation of fennel essential oil caused a 1.5 to 2.5 fold increase in relative sympathetic activity in normal healthy subjects [116] human adverse effects and toxicity: the fruits used under guidance of an herbalist and even without supervision but in moderate use, are generally very safe. The European medicines agency lists allergic reactions affecting skin or the respiratory system as the most common side effect. Intoxication in infants resulting in methemoglobinemia after eating homemade fennel purée has been reported, though all patients recovered after treatment with methylene blue [117]. Long-term use to relieve gas and to regulate intestinal function in children is also suggested to cause premature thelarche [118]. An epileptic patient, with well-controlled epilepsy, developed typical generalized tonic-clonic seizures after consuming a number of cakes containing an unknown quantity of fennel essential oil. Therefore, caution must be exercised while using fennel essential oil in patients with epilepsy [119].

Animal toxicity: Ethanol extract is reported nonlethal and nontoxic to mice up to a single oral dose of 3 g/kg [120-121] and chronic administration of the extract in a dose of 100 mg/kg/day for 90 days caused only significant weight gain in male mice [120]. The oral LD₅₀ of the essential oil in rats is reported to be 1.32 g/kg [122].

CYP450 and potential for drug-drug interactions: water fennel extract is reported to potently inhibit CYP2D6 and CYP3A4 potentially causing a clinically relevant inhibition of intestinal CYP3A4 that may influence the pharmacokinetics of drugs metabolized by this isozyme [123,124]. Warfarin is a drug with a narrow therapeutic window, and one of its main metabolites, 10-hydroxywarfarin is formed by metabolism mediated by CYP3A4, use of fennel with warfarin may thus result in an increased effect of warfarin, and increased chances of bleeding [125]. Concomitant administration of water extract with ciprofloxacin to rats was also reported to affect absorption, distribution and elimination, significantly lowering maximum plasma concentration (C_{max}), area under the curve (AUC) and urinary recovery of ciprofloxacin [126] methanol fruit extract also time-dependently inhibited CYP3A4 in human liver microsomes [127] 5-methoxypsoralen (5-MOP) has been suggested as responsible for the CYP3A4 inhibitory activity [128].

Discussion

From the foregoing description, it is apparent that there are very limited randomized, double-blind, placebo-controlled clinical studies on fennel, except its anti-flatulent and anti-colic effects, to draw any conclusions about various medicinal uses in an objective scientific manner. Nevertheless, its use since ancient times is a testament for its usefulness in certain conditions. It is now established that a number of diseases have cellular damage due to oxidative stress as a component of their pathology. Fennel contains significant amounts of antioxidant substances, such as phenolic compounds and flavonoids that display a broad spectrum of physiological activities and protect against various disease processes. Various extracts of fennel, its essential oil and coumarins isolated from the methanol extract have all shown significant *in vitro* free radicals scavenging capacity, which might be a factor for its usefulness in certain diseases. Qualitative and quantitative differences in chemical constituents of plant-derived drugs are however responsible for the inconsistencies and variations in the often observed pharmacological or clinical effects. Despite the fact that physicians and botanists from the earlier centuries laid down the rules for collecting (harvesting) the plants for medicinal use from a particular location, in a specific season, month or even time of the day, this aspect was ignored and even ridiculed by the scientific community for a long time and was only relatively lately recognized when quantitative evaluation techniques of chemical constituents became available. Variation in chemical constituents of fennel from different parts of the world is but one such example. The pharmacologically active estrogenic agents are polymers of anethole, such as dianethole and photo anethole. Six several components in fennel fruits, working together, are responsible for the observed effects, as the actions of individual components do not wholly explain these effects [8]. Nevertheless, the clinical studies reported are very few, largely not well organized, and in small number of patients. Therefore, full potentials of fennel could be ascertained only after more blinded RCTS in larger number of patients that could supplement the results obtained in experimental studies.

References

1. Badgujar SB, Patel VV, Bandivdekar AH. *Foeniculum vulgare* Mill: a review of its botany, phytochemistry, pharmacology, contemporary application, and toxicology. *Biomed Res Int*. 2014;2014:842674.
2. Endalamaw FD, Chandravanshi BS. Levels of major and trace elements in fennel (*Foeniculum vulgare* Mill.) fruits cultivated in Ethiopia. *Springerplus*. 2015;4:5.
3. Gori L, Gallo E, Mascherini V, Mugelli A, Vannacci A, Firenzuoli F. Can estragole in fennel seed decoctions really be considered a danger for human health? A fennel safety update. *Evid Based Complement Alternat Med*. 2012;860542:2012.
4. Jarić S, Mitrović M, Djurdjević L, Kostić O, Gajić G, Pavlović D, et al. Phytotherapy in medieval Serbian medicine according to the pharmacological manuscripts of the Chilandar Medical Codex (15th, 16th centuries). *J Ethnopharmacol*. 2011;137(1):601-19.
5. Dymock W, Warden CJH, Hooper D. *Pharmacographia Indica*. 1890. Karachi, Pakistan: Reprinted by The Institute of Health and Tibbi Research. Hamdard National Foundation. 1972;2:124-6.
6. Albert-Puleo M. Fennel and anise as estrogenic agents. *J Ethnopharmacol*. 1980;2(4):337-44.
7. Kabeeruddin M. *Kitabul-Advia*, Vol. II Makhzan-al-Mufradat, Delhi:Aligarh Barqi Press. 1937:57-8.

8. Keys John D. Chinese Herbs, Their Botany, Chemistry and Pharmacodynamics. Rutland, Vermont, Tokyo, Japan: Charles E. Tuttle Company. 1985:199.
9. Khory RN, Katrak NN. Materia Medica of India and Their Therapeutics. Delhi-52: Neeraj Publishing House. 1984:292.
10. Nadkarni AK. Indian Materia Medica. 3rd edn. Bombay: Popular Book Depot and Panval Dhootapapeshwar Parakashan Ltd. 1954.
11. Saha JC, Kasinathan S. Ecobolic properties of Indian medicinal plants II. Indian J Med Res. 1961;49:1094-8.
12. Watt JM, Breyer-Brandwijk MG. Medicinal and Poisonous Plants of Southern and Eastern Africa. 2nd edn. Edinburgh: E & S Livingston Ltd. 1962.
13. Anonymous. *Foeniculum vulgare* Miller subsp. *vulgare* var. *dulce* (Miller) Thellung, Fructus. European Medicine Agency, Committee on Herbal Medicinal Products (HMPC).
14. Facchinetti F, Pedrielli G, Benoni G, Joppi M, Verlato G, Dante G, et al. Herbal supplements in pregnancy: unexpected results from a multicentre study. Hum Reprod. 2012;27(11):3161-7.
15. Guarrera PM, Salerno G, Caneva G. Folk phytotherapeutical plants from Maratea area (Basilicata, Italy). J Ethnopharmacol. 2005;99(3):367-78.
16. Barros L, Heleno SA, Carvalho AM, Ferreira IC. Systematic evaluation of the antioxidant potential of different parts of *Foeniculum vulgare* Mill. from Portugal. Food Chem Toxicol. 2009;47(10):2458-64.
17. Martinez M: Las Plantas Medicinale de Mexico. Mexico. 1933.
18. Camacho-Corona Mdel R, Ramírez-Cabrera MA, Santiago OG, Garza-González E, Palacios Ide P, Luna-Herrera J. Activity against drug resistant-tuberculosis strains of plants used in Mexican traditional medicine to treat tuberculosis and other respiratory diseases. Phytoter Res. 2008;22(1):82-5.
19. Ali-Shtayah MS, Jamous RM, Al-Shafie' JH, Elgharabah WA, Kherfan FA, Qarariah KH, et al. Traditional knowledge of wild edible plants used in Palestine (Northern West Bank): a comparative study. J Ethnobiol Ethnomed. 2008;4:13.
20. Jeambey Z, Johns T, Talhouk S, Batal M. Perceived health and medicinal properties of six species of wild edible plants in north-east Lebanon. Public Health Nutr. 2009;12(10):1902-11.
21. Hsu Hong-Yen. How to treat yourself with Chinese Herbs. Oriental Healing Art Institute. 1980:225.
22. Hsu Hong-Yen, Peacher WG. Chinese Herb Medicine and Therapy. Oriental Healing Art Institute. 1976:172.
23. DeFilipps Robert A, Maina Shirley L, Crepin Juliette. Medicinal Plants of the Guianas. (Guyana, Surinam, French Guiana). 2004:17.
24. Kokwaro JO. Medicinal Plants of East Africa. Kampala, Nairobi, Dar es Salam: East African Literature Bureau. 1976.
25. Claus EP, Tyler VE, Brady LR. Pharmacognosy 6th edn., Philadelphia, PA: Lea & Febiger; 1970.
26. Osol A, Pratt R, Altschule MD. The United States Dispensatory and Physicians Pharmacology. 26th edn. Philadelphia. J B Lippincott Co. 1960.
27. Ramstad E. Modern Pharmacognosy. New York: Mc Graw Hill Book Co. Inc. 1959;480.
28. Stecher PG. (Editor). The Merck Index; an Encyclopedia of Chemicals and Drugs. 8th edn. Rahway, N J Merck & Co. 1968:1713.
29. Trease GE, Evans WC. Pharmacognosy. 10th Ed. Baltimore, MD: Williams and Wilkins Co. 1972.
30. Goswami N, Chatterjee S. Assessment of free radical scavenging potential and oxidative DNA damage preventive activity of *Trachyspermum ammi* L. (carom) and *Foeniculum vulgare* Mill. (fennel) seed extracts. Biomed Res Int. 2014;2014:582767.
31. Kaur GJ, Arora DS. Antibacterial and phytochemical screening of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi*. BMC Complement Altern Med. 2009;9:30.
32. Mohamad RH, El-Bastawesy AM, Abdel-Monem MG, Noor AM, Al-Mehdar HA, Sharawy SM, et al. Antioxidant and anti-carcinogenic effects of methanolic extract and volatile oil of fennel seeds (*Foeniculum vulgare*). J Med Food. 2011;14(9):986-1001.
33. Krizman M, Baricevic D, Prosek M. Determination of phenolic compounds in fennel by HPLC and HPLC-MS using a monolithic reversed-phase column. J Pharm Biomed Anal. 2007;43(2):481-5.
34. Kunzemann J, Herrmann K. [Isolation and identification of flavon(ol)-O-glycosides in caraway (*Carum carvi* L.), fennel (*Foeniculum vulgare* Mill.), anise (*Pimpinella anisum* L.), and coriander (*Coriandrum sativum* L.), and of flavon-C-glycosides in anise. I. Phenolics of spices (author's transl)]. Z Lebensm Unters Forsch. 1977;164(3):194-200.
35. Menez J, Castro-Poceiro J. Coumarins in *Foeniculum vulgare* fruits. Rev Latinoam Quim. 1981;12(2):91-2.
36. De Marino S, Gala F, Borbone N, Zollo F, Vitalini S, Visioli F, et al. Phenolic glycosides from *Foeniculum vulgare* fruit and evaluation of antioxidative activity. Phytochemistry. 2007;68(13):1805-12.
37. Mimica-Dukic N, Kujundzic S, Sokovic M, Couladis M. Essential oil composition and antifungal activity of *Foeniculum vulgare* Mill obtained by different distillation conditions. Phytoter Res. 2003;17(4):368-71.
38. Msaada K, Hosni K, Taarit MB, Chahed T, Kchouk ME, Marzouk M. Changes on essential oil composition of coriander (*Coriandrum sativum* L.) fruits during three stages of maturity. Food Chem. 2007;102:1131-4.
39. Raal A, Orav A, Arak E. Essential oil composition of *Foeniculum vulgare* Mill. Fruits from pharmacies in different countries. Nat Prod Res. 2012;26(13):1173-8.
40. Mota AS, Martins MR, Arantes S, Lopes VR, Bettencourt E, Pombal S, et al. Antimicrobial activity and chemical composition of the essential oils of Portuguese *Foeniculum vulgare* fruits. Nat Prod Commun. 2015;10(4):673-6.
41. Díaz-Maroto MC, Díaz-Maroto Hidalgo IJ, Sánchez-Palomo E, Pérez-Coello MS. Volatile components and key odorants of fennel (*Foeniculum vulgare* Mill.) and thyme (*Thymus vulgaris* L.) oil extracts obtained by simultaneous distillation-extraction and supercritical fluid extraction. J Agric Food Chem. 2005;53(13):5385-9.
42. Bajan M, Aprotosoae AC, Spac A, Stanescu U. [Chemical composition of essential oil obtained from Romanian fennel fruits]. Rev Med Chir Soc Med Nat Iasi. 2011;115(2):590-4.
43. Dadalioglu I, Evrendilek GA. Chemical compositions and antibacterial effects of essential oils of Turkish oregano (*Origanum minutiflorum*), bay laurel (*Laurus nobilis*), Spanish lavender (*Lavandula stoechas* L.), and fennel (*Foeniculum vulgare*) on common foodborne pathogens. J Agric Food Chem. 2004;52(26):8255-60.
44. Díaz-Maroto MC, Pérez-Coello MS, Esteban J, Sanz J. Comparison of the volatile composition of wild fennel samples (*Foeniculum vulgare* Mill.) from central Spain. J Agric Food Chem. 2006;54(18):6814-8.
45. Cosge B, Kiralan M, Gürbüz B. Characteristics of fatty acids and essential oil from sweet fennel (*Foeniculum vulgare* Mill. var. *dulce*) and bitter fennel fruits (*F. vulgare* Mill. var. *vulgare*) growing in Turkey. Nat Prod Res. 2008;22(12):1011-6.
46. Abdollahi A, Hassani A, Ghosta Y, Bernousi I, Meshkatsadat MH, Shabani R, et al. Evaluation of essential oils for maintaining postharvest quality of Thompson seedless table grape. Nat Prod Res. 2012;26(1):77-83.
47. Qiu J, Li H, Su H, Dong J, Luo M, Wang J, et al. Chemical composition

- of fennel essential oil and its impact on *Staphylococcus aureus* exotoxin production. *World J Microbiol Biotechnol*. 2012;28(4):1399-405.
48. Wakabayashi KA, de Melo NI, Aguiar DP, de Oliveira PF, Groppo M, da Silva Filho AA, et al. Anthelmintic effects of the essential oil of fennel (*Foeniculum vulgare* Mill., Apiaceae) against *Schistosoma mansoni*. *Chem Biodivers*. 2015;12(7):1105-14.
49. Miguel MG, Cruz C, Faleiro L, Simões MT, Figueiredo AC, Barroso JG, et al. *Foeniculum vulgare* essential oils: chemical composition, antioxidant and antimicrobial activities. *Nat Prod Commun*. 2010;5(2):319-28.
50. Aprotosoia AC, Hâncianu M, Poiată A, Tuchiluş C, Spac A, Cioană O, et al. *In vitro* antimicrobial activity and chemical composition of the essential oil of *Foeniculum vulgare* Mill. *Rev Med Chir Soc Med Nat Iasi*. 2008;112(3):832-6.
51. Shahat AA, Ibrahim AY, Hendawy SF, Omer EA, Hammouda FM, Abdel-Rahman FH, et al. Chemical composition, antimicrobial and antioxidant activities of essential oils from organically cultivated fennel cultivars. *Molecules*. 2011;16(2):1366-77.
52. Piccaglia R, Marotti M. Characterization of some Italian types of wild fennel (*Foeniculum vulgare* Mill.). *J Agric Food Chem*. 2001;49(1):239-44.
53. Zhang F, Li Z, Tian S, Ma L. [Analysis on changes of chemical compounds in different samples of fried *Foeniculum vulgare*]. *Zhongguo Zhong Yao Za Zhi*. 2009;34(7):829-32.
54. Zeller A, Horst K, Rychlik M. Study of the metabolism of estragole in humans consuming fennel tea. *Chem Res Toxicol*. 2009;22(12):1929-37.
55. Raffo A, Nicoli S, Leclercq C. Quantification of estragole in fennel herbal teas: implications on the assessment of dietary exposure to estragole. *Food Chem Toxicol*. 2011;49(2):370-5.
56. Uusitalo L, Salmenhaara M, Isoniemi M, Garcia-Alvarez A, Serra-Majem L, Ribas-Barba L, et al. Intake of selected bioactive compounds from plant food supplements containing fennel (*Foeniculum vulgare*) among Finnish consumers. *Food Chem*. 2016;194:619-25.
57. Baliga MS, Jagetia GC, Rao SK, Babu K. Evaluation of nitric oxide scavenging activity of certain spices *in vitro*: a preliminary study. *Nahrung*. 2003;47(4):261-4.
58. Faudale M, Viladomat F, Bastida J, Poli F, Codina C. Antioxidant activity and phenolic composition of wild, edible, and medicinal fennel from different Mediterranean countries. *J Agric Food Chem*. 2008;56(6):1912-20.
59. Satyanarayana S, Sushruta K, Sarma GS, Srinivas N, Subba Raju GV. Antioxidant activity of the aqueous extracts of spicy food additives--evaluation and comparison with ascorbic acid in *in-vitro* systems. *J Herb Pharmacother*. 2004;4(2):1-10.
60. Celik I, Isik I. Determination of chemo preventive role of *Foeniculum vulgare* and *Salvia officinalis* infusion on trichloroacetic acid-induced increased serum marker enzymes lipid peroxidation and antioxidative defense systems in rats. *Nat Prod Res*. 2008;22(1):66-75.
61. Parejo I, Viladomat F, Bastida J, Rosas-Romero A, Flerlage N, Burillo J, et al. Comparison between the radical scavenging activity and antioxidant activity of six distilled and nondistilled mediterranean herbs and aromatic plants. *J Agric Food Chem*. 2002;50(23):6882-90.
62. Nickavar B, Abolhasani FA. Screening of antioxidant properties of seven Umbelliferae fruits from Iran. *Pak J Pharm Sci*. 2009;22(1):30-5.
63. Yang JJ, Lee DU, Shin HM. Anti-inflammatory and antioxidant effects of coumarins isolated from *Foeniculum vulgare* in lipopolysaccharide-stimulated macrophages and 12-O-tetradecanoylphorbol-13-acetate-stimulated mice. *Immunopharmacol Immunotoxicol*. 2015;37(3):308-17.
64. Choi EM, Hwang JK. Antiinflammatory, analgesic and antioxidant activities of the fruit of *Foeniculum vulgare*. *Fitoterapia*. 2004;75(6):557-65.
65. Toda S. Inhibitory effects of aromatic herbs on lipid peroxidation and protein oxidative modification by copper. *Phytother Res*. 2003;17(5):546-8.
66. Tripathi P, Tripathi R, Patel RK, Pancholi SS. Investigation of antimutagenic potential of *Foeniculum vulgare* essential oil on cyclophosphamide induced genotoxicity and oxidative stress in mice. *Drug Chem Toxicol*. 2013;36(1):35-41.
67. Aazza S, Lyoussi B, Miguel MG. Antioxidant and antiacetylcholinesterase activities of some commercial essential oils and their major compounds. *Molecules*. 2011;16(9):7672-90.
68. Ruberto G, Baratta MT, Deans SG, Dorman HJ. Antioxidant and antimicrobial activity of *Foeniculum vulgare* and *Crithmum maritimum* essential oils. *Planta Med*. 2000;66(8):687-93.
69. Misharina TA, Polshkov AN. [Antioxidant properties of essential oils: autoxidation of essential oils from laurel and fennel and effects of mixing with essential oil from coriander]. *Prikl Biokhim Mikrobiol*. 2005;41(6):693-702.
70. Barros L, Heleno SA, Carvalho AM, Ferreira IC. Systematic evaluation of the antioxidant potential of different parts of *Foeniculum vulgare* Mill. From Portugal. *Food Chem Toxicol*. 2009;47(10):2458-64.
71. Papageorgiou V, Mallouchos A, Komaitis M. Investigation of the antioxidant behavior of air- and freeze-dried aromatic plant materials in relation to their phenolic content and vegetative cycle. *J Agric Food Chem*. 2008;56(14):5743-52.
72. Guimarães R, Barreira JC, Barros L, Carvalho AM, Ferreira IC. Effects of oral dosage form and storage period on the antioxidant properties of four species used in traditional herbal medicine. *Phytother Res*. 2011;25(4):484-92.
73. Aridoğan BC, Baydar H, Kaya S, Demirci M, Özbaşar D, Mumcu E. Antimicrobial activity and chemical composition of some essential oils. *Arch Pharm Res*. 2002;25(6):860-4.
74. Singh G, Kapoor IP, Pandey SK, Singh UK, Singh RK. Studies on essential oils: part 10; antibacterial activity of volatile oils of some spices. *Phytother Res*. 2002;16(7):680-2.
75. Mohsenzadeh M. Evaluation of antibacterial activity of selected Iranian essential oils against *Staphylococcus aureus* and *Escherichia coli* in nutrient broth medium. *Pak J Biol Sci*. 2007;10(20):3693-7.
76. Lixandru BE, Drăcea NO, Dragomirescu CC, Drăgulescu EC, Coldea IL, Anton L, et al. Antimicrobial activity of plant essential oils against bacterial and fungal species involved in food poisoning and/or food decay. *Roum Arch Microbiol Immunol*. 2010;69(4):224-30.
77. Pai MB, Prashant GM, Murlikrishna KS, Shivakumar KM, Chandu GN. Antifungal efficacy of *Punicagranatum*, *Acacia nilotica*, *Cuminumcyminum* and *Foeniculum vulgare* on *Candida albicans*: an *in vitro* study. *Indian J Dent Res*. 2010;21(3):334-6.
78. Alizadeh A, Zamani E, Sharaifi R, Javan-Nikkhah M, Nazari S. Antifungal activity of some essential oils against toxigenic *Aspergillus* species. *Commun Agric Appl Biol Sci*. 2010;75(4):761-7.
79. Zeng H, Chen X, Liang J. *In vitro* antifungal activity and mechanism of essential oil from fennel (*Foeniculum vulgare* L.) on dermatophyte species. *J Med Microbiol*. 2015;64(Pt 1):93-103.
80. Jazani NH, Zartoshti M, Babazadeh H, Ali-daiee N, Zarrin S, Hosseini S. Antibacterial effects of Iranian fennel essential oil on isolates of *Acinetobacter baumannii*. *Pak J Biol Sci*. 2009;12(9):738-41.
81. Orhan IE, Özcelik B, Kan Y, Kartal M. Inhibitory effects of various essential oils and individual components against extended-spectrum beta-lactamase (ESBL) produced by *Klebsiella pneumoniae* and their chemical compositions. *J Food Sci*. 2011;76(8):M538-46.
82. Mahady GB, Pendland SL, Stoia A, Hamill FA, Fabricant D, Dietz BM, et

- al. *In vitro* susceptibility of *Helicobacter pylori* to botanical extracts used traditionally for the treatment of gastrointestinal disorders. *Phytother Res.* 2005;19(11):988-91.
83. Cwikla C, Schmidt K, Matthias A, Bone KM, Lehmann R, Tiralongo E. Investigations into the antibacterial activities of phytotherapeutics against *Helicobacter pylori* and *Campylobacter jejuni*. *Phytother Res.* 2010;24(5):649-56.
84. Birdane FM, Cemek M, Birdane YO, Gülçin I, Büyükkuroğlu ME. Beneficial effects of *Foeniculum vulgare* on ethanol-induced acute gastric mucosal injury in rats. *World J Gastroenterol.* 2007;13(4):607-11.
85. Vasudevan K, Vembar S, Veeraraghavan K, Haranath PS. Influence of intragastric perfusion of aqueous spice extracts on acid secretion in anesthetized albino rats. *Indian J Gastroenterol.* 2000;19(2):53-6.
86. Tognolini M, Ballabeni V, Bertoni S, Bruni R, Impicciatore M, Barocelli E. Protective effect of *Foeniculum vulgare* essential oil and anethole in an experimental model of thrombosis. *Pharmacol Res.* 2007;56(3):254-60.
87. Sadrefozalayi S, Farokhi F. Effect of the aqueous extract of *Foeniculum vulgare* (fennel) on the kidney in experimental PCOS female rats. *Avicenna J Phytomed.* 2014;4(2):110-7.
88. Kim TH, Kim HJ, Lee SH, Kim SY. Potent inhibitory effect of *Foeniculum vulgare* Miller extract on osteoclast differentiation and ovariectomy-induced bone loss. *Int J Mol Med.* 2012;29(6):1053-9.
89. Malini T, Vanithakumari G, Megala N, Anusya S, Devi K, Elango V. Effect of *Foeniculum vulgare* Mill. seed extract on the genital organs of male and female rats. *Indian J Physiol Pharmacol.* 1985;29(1):21-6.
90. Devi K, Vanithakumari G, Anusya S, Mekala N, Malini T, Elango V. Effect of *Foeniculum vulgare* seed extract on mammary glands and oviducts of ovariectomised rats. *Anc Sci Life.* 1985;5(2):129-32.
91. Wright CI, Van-Buren L, Kroner CI, Koning MM. Herbal medicines as diuretics: a review of the scientific evidence. *J Ethnopharmacol.* 2007;114(1):1-31.
92. El Bardai S, Lyoussi B, Wibo M, Morel N. Pharmacological evidence of hypotensive activity of *Marrubium vulgare* and *Foeniculum vulgare* in spontaneously hypertensive rat. *Clin Exp Hypertens.* 2001;23(4):329-43.
93. Abdul-Ghani AS, Amin R. The vascular action of aqueous extracts of *Foeniculum vulgare* leaves. *J Ethnopharmacol.* 1988;24(2-3):213-8.
94. Mansouri E, Kooti W, Bazvand M, Ghasemi Boroon M, Amirzargar A, Afrisham R, et al. The Effect of Hydro-Alcoholic Extract of *Foeniculum vulgare* Mill on Leukocytes and Hematological Tests in Male Rats. *Jundishapur J Nat Pharm Prod.* 2015;10(1):e18396.
95. Tognolini M, Barocelli E, Ballabeni V, Bruni R, Bianchi A, Chiavarini M, et al. Comparative screening of plant essential oils: phenylpropanoid moiety as basic core for antiplatelet activity. *Life Sci.* 2006;78(13):1419-32.
96. El-Sheikh el-SA, Galal AA. Toxic effects of sub-chronic exposure of male albino rats to emamectin benzoate and possible ameliorative role of *Foeniculum vulgare* essential oil. *Environ Toxicol Pharmacol.* 2015;39(3):1177-88.
97. Agarwal R, Gupta SK, Agrawal SS, Srivastava S, Saxena R. Oculohypotensive effects of *Foeniculum vulgare* in experimental models of glaucoma. *Indian J Physiol Pharmacol.* 2008;52(1):77-83.
98. Joshi H, Parle M. Cholinergic basis of memory-strengthening effect of *Foeniculum vulgare* Linn. *J Med Food.* 2006;9(3):413-7.
99. Orhan I, Kartal M, Kan Y, Sener B. Activity of essential oils and individual components against acetyl- and butyrylcholinesterase. *Z Naturforsch C.* 2008;63(7-8):547-53.
100. Mata AT, Proença C, Ferreira AR, Serralheiro MLM, Nogueira JMF, Araújo MEM. Antioxidant and antiacetylcholinesterase activities of five plants used as Portuguese food spices. *Food Chem.* 2007;103(3):778-86.
101. Mesfin M, Asres K, Shibeshi W. Evaluation of anxiolytic activity of the essential oil of the aerial part of *Foeniculum vulgare* Miller in mice. *BMC Complement Altern Med.* 2014;14:310.
102. Boskabady MH, Khatami A, Nazari A. Possible mechanism(s) for relaxant effects of *Foeniculum vulgare* on guinea pig tracheal chains. *Pharmazie.* 2004;59(7):561-4.
103. Shalby AB, Hamza AH, Ahmed HH. New insights on the anti-inflammatory effect of some Egyptian plants against renal dysfunction induced by cyclosporine. *Eur Rev Med Pharmacol Sci.* 2012;16(4):455-61.
104. Ozbek H, Uğraş S, Dülger H, Bayram I, Tuncer I, Öztürk G, et al. Hepatoprotective effect of *Foeniculum vulgare* essential oil. *Fitoterapia.* 2003;74(3):317-9.
105. Kaileh M, Berghe WV, Boone E, Essawi T, Haegeman G. Screening of indigenous Palestinian medicinal plants for potential anti-inflammatory and cytotoxic activity. *J Ethnopharmacol.* 2007;113(3):510-6.
106. Alexandrovich I, Rakovitskaya O, Kolmo E, Sidorova T, Shushunov S. The effect of fennel (*Foeniculum vulgare*) seed oil emulsion in infantile colic: a randomized, placebo-controlled study. *Altern Ther Health Med.* 2003;9(4):58-61.
107. Bae J, Kim J, Choue R, Lim H. Fennel (*Foeniculum vulgare*) and fenugreek (*Trigonella foenum-graecum*) tea drinking suppresses subjective short-term appetite in overweight women. *Clin Nutr Res.* 2015;4(3):168-74.
108. Bokaie M, Farajkhoda T, Enjezab B, Khoshbin A, Karimi-Zarchi M. Oral fennel (*Foeniculum vulgare*) drop effect on primary dysmenorrhea: Effectiveness of herbal drug. *Iran J Nurs Midwifery Res.* 2013;18(2):128-32.
109. ModaresNejad V, Asadipour M. Comparison of the effectiveness of fennel and mefenamic acid on pain intensity in dysmenorrhoea. *East Mediterr Health J.* 2006;12(3-4):423-7.
110. NamavarJahromi B, Tartifzadeh A, Khabnadideh S. Comparison of fennel and mefenamic acid for the treatment of primary dysmenorrhea. *Int J Gynaecol Obstet.* 2003;80(2):153-7.
111. Omidvar S, Esmailzadeh S, Baradaran M, Basirat Z. Effect of fennel on pain intensity in dysmenorrhoea: A placebo-controlled trial. *Ayu.* 2012;33(2):311-3.
112. Yaralizadeh M, Abedi P, Najar S, Namjoyan F, Saki A. Effect of *Foeniculum vulgare* (fennel) vaginal cream on vaginal atrophy in postmenopausal women: A double-blind randomized placebo-controlled trial. *Maturitas.* 2016;84:75-80.
113. Akha O, Rabiei K, Kashi Z, Bahar A, Zaeif-Khorasani E, Kosaryan M, et al. The effect of fennel (*Foeniculum vulgare*) gel 3% in decreasing hair thickness in idiopathic mild to moderate hirsutism, A randomized placebo controlled clinical trial. *Caspian J Intern Med.* 2014;5(1):26-9.
114. Javidnia K, Dastgheib L, Mohammadi Samani S, Nasiri A. Antihirsutism activity of Fennel (fruits of *Foeniculum vulgare*) extract. A double-blind placebo-controlled study. *Phytomedicine.* 2003;10(6-7):455-8.
115. Kim SI, Chang KS, Yang YC, Kim BS, Ahn YJ. Repellency of aerosol and cream products containing fennel oil to mosquitoes under laboratory and field conditions. *Pest Manag Sci.* 2004;60(11):1125-30.
116. Haze S, Sakai K, Gozu Y. Effects of fragrance inhalation on sympathetic activity in normal adults. *Jpn J Pharmacol.* 2002;90(3):247-53.
117. Murone AJ, Stucki P, Roback MG, Gehri M. Severe methemoglobinemia due to food intoxication in infants. *Pediatr Emerg Care.* 2005;21(8):536-8.
118. Türkyilmaz Z, Karabulut R, Sönmez K, Can Başaklar A. A striking and frequent cause of premature thelarche in children: *Foeniculum vulgare*. *J Pediatr Surg.* 2008;43(11):2109-11.
119. Skalli S, Soulaymani Bencheikh R. Epileptic seizure induced by fennel essential oil. *Epileptic Disord.* 2011;13(3):345-7.

120. Shah AH, Qureshi S, Ageel AM. Toxicity studies in mice of ethanol extracts of *Foeniculum vulgare* fruit and *Rutachalepensis* aerial parts. *J Ethnopharmacol.* 1991;34(2-3):167-172.
121. Naga Kishore R, Anjaneyulu N, Naga Ganesh M, Sravya N. Evaluation of anxiolytic activity of Ethanolic extract of *Foeniculum vulgare* in mice model. *Intern J Pharm Pharmaceut Sci.* 2012;4:584-586.
122. Ostad SN, Soodi M, Shariffzadeh M, Khorshidi N, Marzban H. The effect of fennel essential oil on uterine contraction as a model for dysmenorrhea, pharmacology and toxicology study. *J Ethnopharmacol.* 2001;76(3):299-304.
123. Rahimi R, Ardekani MR. Medicinal properties of *Foeniculum vulgare* Mill. In traditional Iranian medicine and modern phytotherapy. *Chin J Integr Med.* 2013;19(1):73-9.
124. Langhammer AJ, Nilsen OG. *In vitro* inhibition of human CYP1A2, CYP2D6, and CYP3A4 by six herbs commonly used in pregnancy. *Phytother Res.* 2014;28(4):603-10.
125. Cuzzolin L, Francini-Pesenti F, Zaffani S, Brocadello F, Pengo V, Bassi A, et al. Knowledge's about herbal products among subjects on warfarin therapy and patient-physician relationship: a pilot study. *Pharmacoepidemiol Drug Saf.* 2007;16(9):1014-7.
126. Zhu M, Wong PY, Li RC. Effect of oral administration of fennel (*Foeniculum vulgare*) on ciprofloxacin absorption and disposition in the rat. *J Pharm Pharmacol.* 1999;51(12):1391-6.
127. Subehan, Usia T, Iwata H, Kadota S, Tezuka Y. Mechanism-based inhibition of CYP3A4 and CYP2D6 by Indonesian medicinal plants. *J Ethnopharmacol.* 2006;105(3):449-55.
128. Subehan, Zaidi SF, Kadota S, Tezuka Y. Inhibition on human liver cytochrome P450 3A4 by constituents of fennel (*Foeniculum vulgare*): identification and characterization of a mechanism-based inactivator. *J Agric Food Chem.* 2007;55(25):10162-7.