



# Phyto-Nutritional Profiles of Broom Weed (*Sida acuta*) Leaf Extract

Shittu<sup>1</sup> and Alagbe JO<sup>2\*</sup>

<sup>1</sup>Department of Animal Production and Health, Ladoké Akintola University of Technology (LAUTECH) Ogbomosho, Nigeria

<sup>2</sup>Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, India

## Abstract

Medicinal plants are abundant in phytochemicals that has significant therapeutic effects. They are relatively cheap, effective and safe in prolong use. This experiment was carried out to investigate the phyto-nutritional profile of broom weed (*Sida acuta*) leaf extract. Proximate analysis *Sida acuta* leaf indicated the presence of dry matter (91.88%), crude protein (18.01%), ash (9.73%), crude fiber (6.24%), ether extract (1.77%) and energy (2760 Kcal/kg). Phytochemical screening of the extract revealed the presence of condensed tannins, hydrolysable tannins, flavonoids, saponins, phenols, oxalate, phytate, alkaloids, terpenoids and glycosides at 0.82%, 2.02%, 4.25%, 0.20%, 0.17%, 0.05%, 0.23%, 0.22%, 0.95% and 0.02% respectively and the vitamin constituents are thiamine (0.33 mg/100 g), ascorbic acid (30.17 mg/100 g), riboflavin (0.05 mg/100 g),  $\beta$ -carotene (0.79 mg/100 g) and niacin (0.41 mg/100 g). Mineral analysis showed that it contained calcium (127.6 mg/100 g), phosphorus (78.6 mg/100 g), potassium (31.6 mg/100 g), magnesium (102.1 mg/100 g), iron (2.14 mg/100 g), manganese (0.60 mg/100 g), copper (0.04 mg/100 g) and zinc (1.75 mg/100 g). Result on amino acid composition showed that they contained lysine (2.11 mg/100 g), arginine (1.77 mg/100 g), aspartic acid (2.13 mg/100 g), threonine (1.85 mg/100 g), histidine (4.71 mg/100 g), serine (2.03 mg/100 g), glycine (1.05 mg/100 g), alanine (3.31 mg/100 g), cystine (5.06 mg/100 g), valine (0.88 mg/100 g), leucine (2.04 mg/100 g), phenylalanine (4.72 mg/100 g), tyrosine (3.51 mg/100 g), isoleucine (2.84 mg/100 g), methionine (0.85 mg/100 g) and proline (1.05 mg/100 g). It was concluded that *Sida acuta* leaf extract is rich in various nutrients and phytochemicals conferring it the ability to perform multiple biological activities and as a natural alternative to antibiotics.

**Keywords:** Medicinal plants; Nutrients; *Sida acuta*; phytochemicals

## OPEN ACCESS

### \*Correspondence:

Alagbe JO, Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India, E-mail: demsonfarms@yahoo.com

Received Date: 16 Nov 2020

Accepted Date: 16 Dec 2020

Published Date: 22 Dec 2020

### Citation:

Shittu, Alagbe JO. Phyto-Nutritional Profiles of Broom Weed (*Sida acuta*) Leaf Extract. *Ann Clin Toxicol*. 2020; 3(2): 1032.

Copyright © 2020 Alagbe JO. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Introduction

The important of plants of various types cannot be over emphasized. Since the time immemorial, plant has been taking so important in various approaches. Several plant leaves are delicacy in the preparation of stew in some tribes, which when its number and types is not complete the stew look awkward and unacceptable to them. The belief is that different plant contributes different nutrient to the stew which make it delicious and nutritional. Therefore plenty plants of proven nutritional/ medicinal quality are of important to many pharmaceutical companies manufacturing a wide range of allopathic medicines, due to their phytochemical properties. This has caused increasing consideration of natural drug to an individual and most companies producing most synthetic drug.

One of the common examples is the use of plant extractive in the production of antimalarial from the plant called *Artemisia annua*. The drug is proven in the cure of malaria. Mostafa et al. [1] identified extract from *Artemisia annua* to be the most effective anti-malarial because it rich in artemisinin, which was extracted from the leaves of *Artemisia annua*. This and several others have made plant to be major source of breakthrough in the pharmaceutical industries.

It is an inevitable fact that increasing research on medicinal plants could pave the way for the discovery of novel therapeutic agents against many diseases that are outstanding diseases [2,3], which threatened human existence. Interestingly, WHO has recognized the significance of traditional medicine in the health-care sector and has assessed that approximately eighty percent of the population living in the developing countries depend on herbal medicines for their primary health care requirement [4]. There are many plants in the tropical countries that have not been harnessed, knowing their Phyto-profile will help to enable many people to be attracted to their usefulness as a result of their Phyto-properties.

Phytochemicals also called active principles or phytochemical substances Ekpo and Etim [5] are the predominant substances present in many medicinal plants which responsible for many observed physiological actions. These phytochemicals are present in many forms such as alkaloids, steroids, tannins, glycosides, volatile oils, phenols, and flavonoids Perumalsamy et al. [2] while Ekpo and Etim [5] and Asaolu [6] stated the active chemicals to consist terpenes, flavonoid, bioflavonoid, benzophenones, xanthenes as well as some metabolites such as tannins, saponins, cyanates, oxalate and anthraquinones which are non-specifically confined to various parts of a plant such as leaves, flowers, bark, seeds, fruits or roots Perumalsamy et al. [2]. All these active ingredients are used in the phototherapy and help to control or prevent many diseases. Ekpo and Etim [5] established that medicinal plants are known to owe their curative potentials to certain biological active substances, which exist in parts of the plants.

*Sida acuta* (broom weed) is one of the plants with medicinal potential qualities and present in abundance in the tropics. It is drought resistance tropical weeds that are common in almost everywhere. *Sida acuta* is an erect, branched and perennial shrub with a woody tap root, hairy branched up to 1 m high and is reproduced from their seeds. The stem is woody, rounded and slender, and is fibrous and hairy especially when young. The leaves are simple and alternate while the inflorescence is solitary and axillary with stalks up to 1.3 cm long jointed about half of the length. The flowers are yellow with five petals and the fruit is capsuled with 5-6 carpels [5]. Therefore, this study was designed to examine the phytochemical properties of *Sida acuta* to make it uses known and popular to the populace.

## Materials and Methods

### Study area

The experiment was carried out at Division of Animal Nutrition, Sumitra Research Farm, Gujarat, India during the month of December to February, 2018.

### Collection and processing of *Sida acuta* extract

Fresh and mature leaves of *Sida acuta* were collected within the farm premises and authenticated by a botanist (Liu Yung). The leaves were washed with running tap water, shade dried in a clean tray for 12 days and pulverized to coarse powder with a Panasonic electric blender Model WS-812 KJ and stored in an air tight container for further analysis. 100 g of the sample was soaked in 1000 ml water and agitated with an electric blender for 10 min, thereafter the mixture was turned into a another container, kept in the refrigerator at 4°C for 72 h and then filtered using filter paper (Whatmann No. 1) to obtain *Sida acuta* leaf extract (SAL).

### Laboratory procedures

Proximate analysis of experimental leaf was determined by standard methods as described by AOAC [7]. Mineral content of SAL was determined using an Atomic Absorption Spectrophotometer (AAS) model 236-12. Phytochemical screening of flavonoids, saponin, phytate, terpenoids, steroids and phenol of SAL and *Sida acuta* leaf was carried out using methods described by Boham and Kocipai-Abyazan (1974), Harbone (1973) and Odebiyi and Sofowora (1978) respectively. Tannins were analysed using Folin-Denis spectrophotometric method [8]. Alkaloids, phytate and oxalate were determined according to the method outlined by Day and Underwood (1986). Vitamin analysis of ascorbic acid, thiamine, riboflavin, niacin

and  $\beta$ -carotene according to the procedures outlined by [9,10,11]. Amino acids were analyzed using commercial diagnostic kits (Humburg, Braunschweig, Germany, Model RS-567A).

### Statistical analysis

All data were subjected to one -way Analysis of Variance (ANOVA) using SPSS (20.0) and significant means were separated using Duncan multiple range test of the same software. Significant was declared if  $P \leq 0.05$ .

## Results and Discussion

The proximate composition of *Sida acuta* is shown in Table 1. Proximate composition can be described as the term used in the field of animal science or feed analysis to means the six components of feed for the animal that are of major or commonly important which are moisture, crude protein, ether extract, crude fiber, crude ash and nitrogen free extracts and they are usually expressed in percentages. Proximate compositions are very important in the formulation of animal diet in other to estimate the value of what the farmer is given to his or her animals. Also for any non-conventional ingredients to be used in animal feeding or as supplement, the proximate needs to be established. According to Raimi et al. [12] plants are basic source of knowledge in our today medicine. Most of the feed given to animal are from plant kingdom, this make it so important to know the profile of all the plant around to harness their importance. *Sida acuta* had 91.88% dry matter, 18.01% crude protein while ash, crude fiber, ether extract and energy level were 9.73%, 6.24%, 1.77% and 2760.00 (Kcal/Kg), respectively. The protein composition and dry matter percentage were similar compare to the 19.13% (CP) and 91.07% (dry matter) report by Raimi et al. [12], but had significantly lower crude fiber (6.24%) compared to 9.50% reported by the authors. Difference observed in crude fiber may be as result of variation in the age at harvest or varieties of the *Sida acuta* leaf. The ADL generally increased with the growing maturity stage of growths for forages [13].

*Sida acuta* leaves had varied phytochemical properties compared to *Sida acuta* leaf extract (Table 2) [14-17]. After assessment, it was discovered that *Sida acuta* contains phytochemical properties which include the saponins, tannins, alkaloids, flavonoids, phytates, phenols, oxalate, terpenoids, glycosides and steroids. Also, the analysis revealed that *Sida acuta* leaf and extract had similar condensed tannin but varied hydrolysable tannin. *Sida acuta* leaf consists of lower (1.52%) hydrolysable tannin compared to *Sida acuta* leaf extract (2.02%). More phytochemical was discovered when compare to the findings of Senthil kumar et al. [18] that reported absence of tannins, saponins. The composition of the leaf extract from this study was lower compare to the report of Chinelo and Miracle [19] except flavonoids (3.86 mg/100 g). The authors reported highest compositions of alkaloids, flavonoids and steroids being 0.55, 2.31 and 1.85 mg/100 g respectively. The differences might be as a result of age at harvest or the variety of

**Table 1:** Proximate composition of *Sida acuta* leaf.

Parameters	% Composition
Dry matter	91.88
Crude protein	18.01
Ash	9.73
Crude fiber	6.24
Ether extract	1.77
Energy (Kcal/kg)	2760

**Table 2:** Phytochemical analysis of test material [2,3,7,26].

Parameters	<i>Sida acuta</i> leaf	<i>Sida acuta</i> leaf extract	*Permissible range (%)
Condensed tannin	0.73	0.82	1.88
Hydrolysable tannin	1.52	2.02	2.56
Flavonoids	3.86	4.25	12.1
Saponin	0.19	0.2	7.02
Phenols	0.15	0.17	-
Oxalate	1.01	0.05	0.54
Phytate	0.57	0.23	2.3
Alkaloids	0.41	0.22	2.13
Terpenoids	0.58	0.95	-
Glycosides	0.1	0.02	-
Steroids	1.42	1.05	-

**Table 3:** Vitamin composition of *Sida acuta* leaf.

Parameters	Composition (mg/100 g)	*Requirement (mg)	**% RDA
Thiamine (B <sub>1</sub> )	0.33	1.8	31
Ascorbic acid (C)	30.17	36.3	39
Riboflavin (B <sub>2</sub> )	0.05	10	34
β-carotene	0.79	1	6
Niacin (B <sub>3</sub> )	0.41	2.6	5

\*\*USDA (2010)

**Table 4:** Mineral composition of *Sida acuta* leaf.

Parameters	Composition (Mg/100 g)
<b>Macro elements</b>	
Calcium	127.6
Phosphorus	78.5
Potassium	31.6
Magnesium	20.9
Sodium	102.1
<b>Micro elements</b>	
Iron	2.14
Manganese	0.6
Copper	0.04
Zinc	1.75

the *Sida acuta*. *Abutilon indicum*, *Gossypium herbaceum*, *Hibiscus mutabilis*, *Hibiscus sabdariffa*, *Hibiscus rosa-sinensis*, *Sida acuta*, *Sida cordifolia*, *Sida rhombifolia* and several others are the genera with great ethno medicinal values [20]. The study of Richa and Sharma [21] indicated the high presence amounts of alkaloids, flavonoids, terpenoids and glycosides in methanolic leaf extract of *S. acuta* and presence of these phyto constituents in *S. acuta* showed excellent anti-bacterial and anti-malarial properties [22,23]. Alkaloid presence has been implicated to process the pharmacological property [19].

Phytochemicals are present in many plants, this help the plant to elicit the chemotherapy common to them. The was established by Edeoga et al.[24] when *Cleome nutidosperma*, *Emilia coccinea*, *Euphorbia heterophylla*, *Physalis angulata*, *Richardia bransitensis*, *Scopania dulcis*, *Sida acuta*, *Spigelia anthelmia*, *Stachytarpheta cayennensis* and *Tridax procumbens* were accessed. All the plants were found to contain alkaloids, tannins and flavonoids except for

**Table 5:** Amino acid composition of *Sida acuta* leaf [36].

Amino acids	Composition (mg/100 g)	*Reference level
Lysine	2.11	5.5
Arginine	1.77	1
Aspartic acid	2.13	-
Threonine	1.85	0.65
Histidine	4.71	0.3
Serine	2.03	-
Glycine	1.05	1.2
Alanine	3.31	-
Cystine	5.06	0.35
Valine	0.88	0.82
Leucine	2.04	1.2
Phenylalanine	4.72	0.5
Tyrosine	3.51	0.45
Isoleucine	2.84	0.6
Proline	1.05	0.2
Methionine	0.85	0.35

the absence of tannins in *S. acuta*. The absence of tannins in the *Sida acuta* was contrary to the result of this study. The differences in the report may be as a result of species assessed or methodology employed in the assessment.

Vitamins are organic nutrients that are necessary for growth and development of animals. Body needs vitamins to function properly and most vitamins cannot be produced by the animal in sufficient quantity on their own. Therefore, there is need to obtained them from the feeds fed. Plant can manufacture vitamin necessary for their metabolism, human being and animal cannot do the same except vitamin D. This may be because of lack of some necessary enzyme or pigments [25]. Animal and human need to inject the vitamin through the feed/food and *Sida acuta* has Thiamine (B<sub>1</sub>), Ascorbic acid (C), Riboflavin (B<sub>2</sub>), β-carotene and Niacin (B<sub>3</sub>) (Table 3) that can improve the animals' need of them.

The composition of thiamine (0.33 mg/100 g) in this plant was similar to the report of Olivier [26] (0.36 mg/100 g) and Nwankpa et al. [27] (0.36 mg/100 g) while the quantity of niacin (0.41 mg/100 g), ascorbic acid (30.17 mg/100 g) recorded in this study was higher when compared with the report of Olivier [26] and Nwankpa et al. [27]. Although the authors reported higher (0.12 and 0.12 mg/100 g, respectively) riboflavin compared to the report of this research (0.05 mg/100 g). According to John and Richard [28], riboflavin can be synthesized de novo by plant, yeast, and prokaryotic cells, mammals need to obtain riboflavin by consuming plant-based foods or, if necessary, supplemental sources. Riboflavin is very important vitamin in animal system; it is the precursor to the coenzymes, Flavin Mononucleotide (FMN) and Flavin Adenine Dinucleotide (FAD) [28]. The author implicated riboflavin deficiency to have possible genetic defects in human riboflavin transporter.

Higher vitamin C compare to other research results might be an added advantage as a result of numerous roles it plays in animals. According to Khalid et al. [29] vitamin C is essential for the development and maintenance of connective tissues. It plays an importance role in bone formation, wound healing and the maintenance of healthy gums. It plays other importance role such

as activation of vitamin B, folic acid, conversion of cholesterol to bile acids and tryptophan to the neurotransmitter, serotonin. It also performs the role of antioxidant that protects the body from free radicals' damages. Vitamin C is also used as therapeutic agent in many diseases and disorders; it helps to protect the immune system, reduces the severity of allergic reaction and helps to fight infections. Given plant with such quality to the animal will help to maintain healthy livestock products.

Thiamine is another vitamin present in *Sida acuta*. Thiamine is very important vitamin in animal nutrition because it cannot be produced sufficiently by ruminants; given *Sida acuta* to such ruminants will therefore improve the health status of the animal. Thiamine deficiency was identified to be common in ruminants primarily because of the inability to activate the vitamin through the help of rumen microbes, it causes cerebrocortical necrosis and neurological signs [30,31].

Table 4 shows the mineral composition of *Sida acuta* leaf (mg/100 g). The macro elements determined include calcium, phosphorus, potassium, magnesium and sodium while the micro elements include iron, manganese, copper and zinc. Except sodium and magnesium, other mineral elements observed in this experiment were higher than the reported values according to Raimi et al. [12]. Although experience has shown that phyto-nutrient contents of plants cannot be the same, so the variations are much expected. This is because many factors contributed to the soil nutrients availability and plant uptake, but the present of these minerals in this amount is an added advantage to the use of *Sida acuta* leaf meal. Calcium for example, it is the most abundant mineral in the body of both animal and man (98%). It is found in the bone and teeth. Helps in blood clotting, membrane permeability, muscle contraction, nerve functioning etc [32]. The variation in the phyto-nutrients could be as a result of different soil type or varieties. The different may also be as a result of water/or moisture content in the soil. It has been stated by Ogundola et al. [33] that type of soil, soil texture and effect of environmental exposure can influence the nutritional composition of plant. According to Paetsch et al. [34] established that plant mineral contents depends upon the interaction of number of factors which include soil, plant species, stage of maturity, yield, pasture management and climate. It was also discovered by the authors that heavy rainfall and high temperature accentuate the soil mineral deficiency.

The amino acid profile of *Sida acuta* is presented in Table 5 [35]. Amino acids are building blocks of proteins and polypeptides that are the major component of animals' muscles and tissues. Amino acid also has been identified to be part of some fluid such as milk and also it plays significant role in the biochemical and metabolic processes in cell of all the animals [36]. Considering, the contribution such as body building block, important fluid constituent, metabolic activities, growth, production, and reproduction etc, this make the determination of the constituent in some plant like *Sida acuta* very important. All the sixteen amino acids determine are very important to the animal. According to Vernon and Peter [38], plant protein food contributes approximately 65% of the per capita supply of protein on worldwide basis; therefore *Sida acuta* is a valuable plant, considering all the available amino acid. Although literatures are scarce on the amino acid profile of *Sida acuta*, but it has been documented that nitrogen is the fourth most abundant compound in plants. This nitrogen is a component of proteins nucleic acids, chlorophyll, diverse secondary compounds and many cellular structures [37].

Also, the proven performance of rabbit fed *Sida acuta* meal according to Udeh and Ekwe O [39], Naandam et al. [40], Naandam and Jerebi [41] has shown the protein quality of the leaf in the diet of rabbits.

## References

- Mostafa AE, Melissa JT, Nicholas GR, Douglas G, Pamela JW, Stephen MR. Dried whole plant *Artemisia annua* as an antimalarial therapy. PLoS One. 2012;7(12):e52746.
- Perumalsam M, Sreenath KB, Kannappan P. Identification of active pharmaceuticals of *Sida acuta* Burm. F. leaves using GC-MS and HPTLC fingerprinting. IJPSR. 2019;10(3):1194-207.
- Purchase HG, Arp LH, Domermuth CH, Pearson JEA. Laboratory manual of isolation and identification of avian pathogens. Iowa: Kendall Hunt Publishing Company, USA. 1989.
- Negash B, Addisu AG, Tarekegne T. Knowledge, attitude, and practice of complementary and alternative medicine among residents of Wayu town, Western Ethiopia. J Evid Based Complementary Altern Med. 2017;22(4):929-35.
- Ekpo MA, Etim PC. Antimicrobial activity of ethanolic and aqueous extracts of *Sida acuta* on microorganisms from skin infections. J Med Plants Res. 2009;3(9):621-24.
- Asaolu MF. Chemical composition and phytochemical screening of the seeds of *Garcinia kola*. Pakistan J Sci Ind Res. 2007;46(3):145-7.
- A.O.A.C. (1990). Association of Official Analytical Chemists. 15<sup>th</sup> Ed. Washington, D.C. USA. 1990.
- Shabbir M, Khan MR, Saeed N. Assessment of phytochemicals, antioxidant, anti-lipid peroxidation and anti-haemolytic activity of extract and various fractions of maytenusroleanus leaves. BMC Complement Altern Med. 2013;13(143):1472-6882.
- Okwu DE, Josiah C. Evaluation of the chemical composition of two Nigerian medicinal plants. Afr J Biotechnol. 2006;5(4):357-51.
- Uzomah A, Nwosu JN, Ndupuh CE. Laboratory manual in food quality control. Department of Food Science and Technology Owerri, Imo State. 2002;8-10.
- Augustine J. Simultaneous analysis of B-complexes in: Chromatographic assay of vitamin. 4<sup>th</sup> Ed. Methods of vitamin assay. 1985;1(1):87-8.
- Raimi MM, Oyekanmi AM, Adegoke BM. Proximate, phytochemical and micronutrient composition of *sida acuta*. IOSR J Applied Chem. 2014;7(2):93-8.
- Homolka P, Koukolová V, Podsedníček M, Hlaváčková A. Nutritive value of red clover and Lucerne forages for ruminants estimated by *in vitro* and *in vivo* digestibility methods. Czech J Anim Sci. 2012;10(4):454-568.
- Alagbe JO, Oluwafemi RA. Performance and haematological parameters of broiler chicks given different levels of dried lemon grass and garlic extract. Res: Agricultural Vet Sci. 2019;3(2):102-11.
- Alagbe JO, Sharma DO, Xing Liu. Effect of *Piliostigma thonningii* leaf extracts on the haematological and serum biochemical indices of broiler chickens. Noble Int J Agri Food Technol. 2019;1(2):62-9.
- Alagbe JO. Proximate mineral and phytochemical analysis of *Piliostigma thonningii* stem bark and roots. Int J Biol Physical Chem Studies. 2019;1(1):1-7.
- Olafadehan OA, Oluwafemi RA, Alagbe JO. Performance haematobiochemical parameters of broiler chicks administered Rolfe (*Daniellia oliveri*) leaf extract as an antibiotic alternative. Drug Discovery. 2020;14(33):135-45.
- Senthilkumar RP, Bhuvaneshwari V, Malayaman V, Ranjithkumar R, Sathiyavimal S. Phytochemical screening of aqueous leaf extract of *Sida acuta* Burm. F. and its antibacterial activity. JETIR. 2018;5(8):473-78.

19. Chinelo AE, Miracle OE. Phytochemical and antimicrobial investigations on various parts of *Sida acuta* Burm F. *J Ayu Herb Med*. 2018;4(2):71-5.
20. Rahman A, Gondha R. Taxonomy and traditional medicine practices on malvaceae (mallow family) of Rajshahi, Bangladesh. *Open J Bot*. 2014;1(2):19-24.
21. Richa SS, Sharma ML. Phytochemical investigations and anatomical study of three species of *Sida acuta*. *Bio life*. 2014;2(2):622-9.
22. Alka J, Kumar P. Antibacterial activity of *S. acuta* Burm. F. against human pathogens. *Asian J Pharm Clin Res*. 2012;5(3):33-35.
23. Karou D, Dicko MH, Sanon S, Simpore J, Traore AS. Antimalarial activity of *S. acuta* Burm. F. (Malvaceae) and *Pterocarpus erinaceus* Poir. (Fabaceae). *J Ethnopharmacol*. 2003;89(2):291-94.
24. Edeoga HO, Okwu DE, Mbaebie BO. Phytochemical constituents of some Nigerian medicinal Plants. *Afr J Biotechnol*. 2005;4(7):685-8.
25. Jimoh MO, Afolayan AJ, Lewu FB. Suitability of *Amaranthus* species for alleviating human dietary deficiencies. *S. Afr J Bot*. 2018;115:65-73.
26. Olivier TT, Armel JS, Francis NT. Ethnomedicinal uses, phytochemical and pharmacological profiles, and toxicity of *Sida acuta* Burm. F.: A review article. *The Pharma Innovation J*. 2017;6(6):01-06.
27. Nwankpa P, Chukwuemeka OG, Uloneme GC, Etteh CC, Ugwuezumba P, Nwosu D. Phyto-nutrient composition and ant oxidative potential of ethanolic leaf extract of *Sida acuta* in wistar albino rats. *Afr J Biotechnol*. 2015;14(49):3264-69.
28. John TP, Richard SR. Riboflavin (Vitamin B2). *Handbook of Vitamins*. 3<sup>rd</sup> EEd. Newyork: Marcel, Dekker Inc. 2001.p. 225.
29. Khalid I, Alam K, Muzaffar M. Biological significance of Ascorbic acid (vitamin C) in human health – A review. *Pak J Nutr*. 2004;3(1):5-13.
30. Jessica EM, Cailin RH, Lisa MF. Thiamine deficiency in dogs and cats. *J Am Vet Med Association*. 2013;243(5):649-56.
31. Larbier ML. Metabolism of water and minerals in “Nutrition and feeding of poultry”. Nottingham University Press, Nottingham. 1992. p. 101-118.
32. RickM. Minerals. Small Ruminant Series. 2020.
33. Ogundola AF, Bvenura C, Afolayan AJ. Nutrient and anti nutrient compositions and heavy metal uptake and accumulation in *S. nigrum* cultivated on different soil types. *Sci World J*. 2018;2018:1-20.
34. Paetsch L, Mueller CW, Rumpel C, Angst Š, Wiesheu AC, Girardin C, et al A multi-technique approach to assess the fate of biochar in soil and to quantify its effect on soil organic matter composition. *Org Geochem*. 2017;112:177-86.
35. Dean WF, Scott HM. The development of an amino acid reference diet for the early growth of chicks. *J Poult Sci*. 1965;44:803-08.
36. Kemin. Amino acid for animal health. 2019.
37. Sandra J. The occurrence of amino acid in agricultural soil and their uptake by plants. Swedish University of Agricultural Sciences. 2010.
38. Vernon R, Peter LP. Plant protein in relation to human protein and amino acid nutritions. *Am J Clin Nutr*. 1994;59(5 Suppl):1203S-12S.
39. Udeh I, Ekwe O, Aaron E. Performance of weaner rabbits fed panicum maximum, *entrosemapubescens* and *Sida acuta* supplemented with poultry growers mash. *Anim Res Int*. 2007;4(3):750-52.
40. Naandam J, Padi BAY, Bigol P, Mensah-Kumi R. Use of *Stylosanthes hamata* and *Sida acuta* as sole feeds for rabbits (*Oryctolagus cuniculus*). *Online J Anim Feed Res*. 2012;2(2):182-8.
41. Jerebi DA, Naandam J. Haematological characteristics of local rabbits fed *Sida acuta* and *Synedrella nodiflora*. *Ghana J Sci Technol Dev*. 2015;2(1):40-44.