



Post-Intubation Tracheoesophageal Fistula in Severe SARS-CoV-2 - COVID-19 Patient: Unexpected Finding during Swallowing Test

Fabrizia Elli*

¹Department of Department of Biotechnology and Life Sciences, University of Insubria, Italy

Abstract

Introduction: COVID-19 pandemic, caused by Severe Acute Respiratory Syndrome Coronavirus 2, led to an increase in intensive care admission for massive interstitial pneumonia. Many patients underwent long periods of endotracheal intubation for mechanical ventilation and subsequent tracheostomy. The most common complications related to these maneuvers are known: Laryngotracheal granulomas, webs, stenosis and malacia. Less common complications are tracheal necrosis with trachea-esophageal or tracheoarterial fistulas, caused by compression between the orotracheal tube cuff and the nasogastric tube.

Case Report: We present the case of a patient with severe COVID-19 associated pneumonia who underwent orotracheal intubation and prolonged mechanical ventilation for two months with unexplained episodes of food inhalation. During the swallowing test, it was observed inhalation, without natural passage of food from the glottic plane. Therefore, ENT specialist in swallowing, suspecting the feared tracheoesophageal fistula, requested a double contrast esophageal X-ray that confirmed the diagnostic hypothesis. Thoracic surgery team performed surgical correction by resection and tracheal anastomosis with esophago rafia.

Discussion: TEF is a very rare, but potentially fatal, complication of invasive VM. Predisposing factors are prolonged periods of invasive VM, elevated cuff pressures, oversized endotracheal tubes, patient comorbidities, local and general infections, and use of a nasogastric tube.

The endoscopic swallow test is a relatively simple test, performed at the patient's bed that can identify oropharyngeal swallowing disorders and esophageal changes, such as tracheoesophageal fistula.

OPEN ACCESS

*Correspondence:

Fabrizia Elli, Department of Biotechnology and Life Sciences, University of Insubria, Ospedale di Circolo e Fondazione Macchi, Via Guicciardini 9, 21100, Varese, Italy, E-mail: fabrizia.elli@hotmail.it

Received Date: 25 Mar 2022

Accepted Date: 14 Apr 2022

Published Date: 20 Apr 2022

Citation:

Elli F. Post-Intubation Tracheoesophageal Fistula in Severe SARS-CoV-2 - COVID-19 Patient: Unexpected Finding during Swallowing Test. *Am J Otolaryngol Head Neck Surg.* 2022; 5(5): 1192.

Copyright © 2022 Fabrizio Elli. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

The SARS-CoV-2 pandemic has spread dramatically and rapidly around the world. This disease is a highly contagious pandemic zoonosis caused by an RNA beta-coronavirus. In most patients, the disease takes a mild form with symptoms such as fever, cough, nausea, vomiting, and diarrhea. However, in some individuals, it can cause massive involvement of the lower respiratory tract resulting in interstitial pneumonia [1]. Despite the low mortality rate (1.4% to 2.3%) and the relatively low number of patients requiring intensive care (between 5 and 12%), due to its high transmissibility the infection has challenged the global healthcare system [2,3]. There has been unprecedented overcrowding of hospital facilities, particularly Intensive Care Units (ICUs). This followed the development of severe hypoxemia related to acute respiratory distress syndrome (60% to 70% of cases), shock (30%), myocardial dysfunction (20% to 30%), and acute renal failure (10% to 30%) [4].

The significant increase in the number of patients requiring admission to Intensive Care Units (ICUs) and prolonged invasive Mechanical Ventilation (MV) has brought to light an important number of complications. The most common are: Laryngotracheal granulomas, spider webs, stenosis, malacia up to tracheal necrosis with trachea-esophageal or tracheo-arterial fistulas [5].

Case Presentation

We report the case of a COVID-19 patient who underwent prolonged invasive VM for two months and developed FET. Tracheoesophageal Fistula (TEF) is a connection between the trachea and esophagus, which can be congenital or acquired. TEF following prolonged invasive VM is a rare condition, with an incidence of 0.3% to 3% [6].



Figure 1: Barium esophageal rx showed a tracheoesophageal fistula located below tracheostomy cannula (Arrowhead-TEF).



Figure 2: Bronchoscopy and gastroscopy confirmed tracheoesophageal fistula (a and b).



Figure 3: Negative postoperative barium esophageal rx.

A 58-year-old female patient with a prior history of hypertension and ovarian carcinoid presented to the Emergency Department (ED) in May 2021, complaining of dyspnea and cough for several days. On admission, she tested positive for Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). Clinical, laboratory and imaging examinations confirmed the diagnosis of pneumonia associated with SARS-CoV-2, with severe hypoxemic respiratory failure.

The patient was therefore admitted to the intensive care unit and was intubated and mechanically ventilated. However, the clinical evolution was unfavorable with persistent PaO₂/FiO₂ ratio <150 and the patient underwent percutaneous tracheostomy and PEG placement. In September 2021, during rehabilitation, performed by speech therapist specializing in swallowing disorders, uncommon swallowing alterations appeared. Therefore, the patient underwent specialized ENT evaluation by endoscopic endonasal swallowing test with food of various densities. During the examination, normal oropharyngeal transit was initially observed, with subsequent onset

of coughing and tracheal bolus ascension. The fibroendoscope was therefore removed to allow the patient to adequately clear the ingestions. At the subsequent re-evaluation, no traces of food residues were found in the laryngeal vestibule. This finding led to the suspicion of the presence of a passage between the esophagus and trachea. So, the specialist recommended a double-contrast esophageal X-ray with video-recorded swallowing study confirming a tracheo-esophageal fistula of the cervical esophageal portion. The thoracic surgery team, after identifying and evaluating the FET by bronchoscopy and gastroscopy, performed surgical correction by resection and tracheal anastomosis with esophagography, re-establishing the correct anatomical boundaries.

Discussion

Acquired TEF is a rare but potentially serious complication of prolonged intubation for mechanical ventilation. Several factors facilitate its development: High cuff pressures, oversized endotracheal tubes, patient comorbidities, corticosteroid use, local infections, nasogastric tube, need to change position (prone supine) frequently with possible excessive tube movement, repeated bronchoscopy, and high PEEP ventilation over a period of 4 weeks [7,8].

The classic presentation of TEF is represented by a sudden drop in oxygen saturation, despite adequate ventilation, associated with positive cuff leakage and gastric distension.

This was not the case in our case, but the presence of fistula was suspected only during Fiberoptic Endoscopic Evaluation of Swallowing (FEES). During the examination, there was an episode of massive inhalation with food at tracheal, level without observing bolus at subglottic glottic level in absence of frank regurgitation or alteration of the epiglottic tilt.

Despite the high morbidity and mortality of TEF, which is even higher in patients with COVID-19, surgical repair is generally required because spontaneous closure of a FET is rare [9].

In summary, a COVID-19 patient with respiratory failure developed a FET that was only suspected during swallow testing, as he had not previously given symptoms. This suggests that, alongside radiological (CT) or operative (bronchoscopy) examinations [10], a relatively simple and inexpensive examination such as endoscopic swallowing testing may have diagnostic value. The advantages of this test are the possibility to perform it at the patient's bedside, the low complications, and the reproducibility over time. So, in this way, it is possible to identify oropharyngeal swallowing disorders and esophageal alterations, such as tracheoesophageal fistula, and monitor their evolution or regression over time.

References

1. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of Coronavirus disease 2019 in China. *N Engl J Med*. 2020;382:1708-20.
2. Wang D, Hu B, Hu C. Clinical characteristics of 138 hospitalized patients with 2019 novel Coronavirus-infected pneumonia in Wuhan. *JAMA*. 2020;323(11):1061-9.
3. Kowalski LP, Sanabria A, Ridge JA, de Bree, Rinaldo A. COVID-19 pandemic: Effects and evidence-based recommendations for otolaryngology and head and neck surgery practice. *Head Neck*. 2020.
4. Phua J, Weng L, Ling L, Egi M, Man Lim C, Divatia JV, et al. Intensive care management of Coronavirus disease 2019 (COVID-19): Challenges and recommendations. *Lancet Respir Med*. 2020;8(5):506-17.

5. Nikolovski N, Kopacheva-Barsova G, Pejkovska A. Laryngotracheal stenosis: A retrospective analysis of their aetiology, diagnose and treatment. *Open Access Maced J Med Sci.* 2019;7(10):1649-56.
6. Couraud L, Ballester MJ, Delaisement C. Acquired tracheoesophageal fistula and its management. *Semin Thorac Cardiovasc Surg.* 1996;8(4):392-9.
7. Wolf M, Segal E, Yellin A, Faibel M, Talmi YP, Kronenberg J. Acquired tracheoesophageal fistula in critically ill patients. *Ann Otol Rhinol Laryngol.* 2000;109(8):731-5.
8. Mehel DM, Ozdemir D, Çelebi M, Aydemir S, Akgul G, Ozgur A. Classification of laryngeal injury in patients with prolonged intubation and to determine the factors that cause the injury. *Am J Otolaryngol.* 2020;41(3):102432.
9. Marulli G, Loizzi M, Cardillo G, Battistella L, De Palma A, Ialongo P, et al. Early and late outcome after surgical treatment of acquired non-malignant tracheo-oesophageal fistulae. *Eur J Cardiothorac Surg.* 2013;43(6):e155-e61.
10. Mooty RC, Rath P, Self M, Dunn E, Mangram A. Review of tracheoesophageal fistula associated with endotracheal intubation. *J Surg Educ.* 2007;64(4):237-40.