



Red Blood Cells Distribution Width as Biomarker of Worsening in Atherosclerosis

Calogera Pisano^{1*}, Carmela Rita Balistreri², Paolo Nardi¹, Fabio Bertoldo¹, Antonio Pellegrino¹, Renato Massoud² and Giovanni Ruvo¹

¹Department of Cardiac Surgery, Tor Vergata University Hospital, Italy

²Department of Clinical Biochemistry, Tor Vergata University Hospital, Italy

³Department of Pathobiology and Medical and Forensic Biotechnologies, University of Palermo, Italy

Keywords

Red blood cells distribution width; Inflammation; Aging; Cardio

Editorial

Red blood cell Distribution Width (RDW) is a parameter of the circulating erythrocytes' size that has reported to be associated with Cardiovascular Diseases (CVDs) [1-7]. The related mechanism of this association is not fully understood. However, RDW is an indicator of inflammation related to early inflammatory biomarkers, such as C-reactive protein (CRP), interleukin-6 (IL-6) and tumor necrosis factor- α (TNF- α) [8,9]. Accordingly, systemic chronic inflammation leads to dysfunctional bone marrow with unsuccessful production of red blood cells [10]. As a result, it determines the migration of reticulocytes into the peripheral circulation, followed by an increase in circulating levels of immature Red Blood Cells (RBCs), as well as in higher RDW levels [9]. Moreover, inflammation up-regulates the expression of complement proteins and Toll-Like Receptors (TLRs) in platelets. This contributes to platelet activation, by accelerating the onset of atherosclerosis disease [10]. Thus, inflammation is the principal pathophysiological inducer in the development of atherosclerosis, but also of other CVDs [11-14]. In addition, an increase in RDW levels also is related to augmented oxidative damage in blood circulation [15], which associates with exacerbation of pathophysiological conditions linked to the worsening of atherosclerosis [16]. This leads to hypothesize, that interplay between RDW and atherosclerosis might to be related to the inflammatory state, which can inhibit erythropoiesis both directly through the action of cytokines able to increase resistance to the effect of erythropoietin, and indirectly by the reduction of iron levels [17]. Inflammatory state determines a reduction of iron levels, which reduces the capacity of the bone marrow to generate new erythrocytes. This is due to lower absorption of iron and its largest storage [18]. In addition to inflammation, the cholesterol content in the erythrocyte membrane has been suggested to be the link between RDW and atherosclerosis. Pathological elevations of cholesterolemia lead to an excessive increase in membrane cholesterol content [19,20] and decreased erythrocyte stability and deformability. A decrease in deformability should be detrimental to the return to the trivial forms of the erythrocyte in the face of morphological changes resulting from the application of the mechanical forces of the blood circulation itself, which would generate a great multiplicity of forms. But certainly the decrease in deformability makes erythrocytes more susceptible to lysis and thus the population of younger erythrocytes would tend to suffer not only relative but also absolute increase [21], with a consequent alteration in the distribution of volumes, since the younger erythrocytes have greater volume when compared to the older ones [22]. Indeed, in a recent study with healthy volunteers, RDW showed a negative correlation with red blood cell count and blood hemoglobin levels, although no association was found between RDW and lipid levels [23-27]. The important role of RDW in the atherosclerotic process might explain the strong association between this new biomarker and the increased risk of acute coronary syndrome [28-33], ischemic cerebrovascular disease [34-38] and peripheral artery disease as recent and interesting studies have reported [39-41].

OPEN ACCESS

*Correspondence:

Calogera Pisano, Department of Cardiac Surgery, Tor Vergata University Hospital, Italy, Tel: 0039-3283297692; E-mail: bacalipi@libero.it

Received Date: 19 Jan 2018

Accepted Date: 02 Feb 2018

Published Date: 15 Feb 2018

Citation:

Pisano C, Balistreri CR, Nardi P, Bertoldo F, Pellegrino A, Massoud R, et al. Red Blood Cells Distribution Width as Biomarker of Worsening in Atherosclerosis. *Ann Atheroscler Res*. 2018; 1(2): 1007.

Copyright © 2018 Calogera

Pisano. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

1. Al-Najjar Y, Goode KM, Zhang J, Cleland JG, Clark AL. Red cell distribution width: an inexpensive and powerful prognostic marker in heart failure. *Eur J Heart Fail*. 2009;11(12):1155-62.
2. Felker GM, Allen LA, Pocock SJ, Shaw LK, McMurray JJ, Pfeffer MA, et al. Red cell distribution width as a novel prognostic marker in heart failure: data from the CHARM Program and the Duke Databank. *J Am Coll Cardiol*. 2007;50(1):40-7.

3. Van Kimmenade RR, Mohammed AA, Uthamalingam S, Van der Meer P, Felker GM, Januzzi JL. Red blood cell distribution width and 1-year mortality in acute heart failure. *Eur J Heart Fail*. 2010;12(2):129-136.
4. Uyarel H, Ergelen M, Cicek G, Kaya MG, Ayhan E, Turkkan C, et al. Red cell distribution width as a novel prognostic marker in patients undergoing primary angioplasty for acute myocardial infarction. *Coron Artery Dis*. 2011;22(3):138-44.
5. Tonelli M, Sacks F, Arnold M, Moye L, Davis B, Pfeffer M, et al. Relation Between Red Blood Cell Distribution Width and Cardiovascular Event Rate in People With Coronary Disease. *Circulation*. 2008;117(2):163-8.
6. Dabbah S, Hammerman H, Markiewicz W, Aronson D. Relation between red cell distribution width and clinical outcomes after acute myocardial infarction. *Am J Cardiol*. 2010;105(3):312-7.
7. Poludasu S, Marmur JD, Weedon J, Khan W, Cavusoglu E. Red cell distribution width (RDW) as a predictor of long-term mortality in patients undergoing percutaneous coronary intervention. *Thromb Haemost*. 2009;102(3):581-7.
8. Azab B, Torbey E, Hatoum H, Singh J, Khoueiry G, Bachir R, et al. Usefulness of red cell distribution width in predicting all-cause long-term mortality and non-ST-elevation myocardial infarction. *Cardiology*. 2011;119(2):72-80.
9. Weiss G, Goodnough LT. Anemia of chronic disease. *N Engl J Med*. 2005;352(10):1011-23.
10. Arbel Y, Shacham Y, Finkelstein A, Halkin A, Milwidsky A, Berliner S, et al. Red blood cell distribution width (RDW) and long-term survival in patients with ST elevation myocardial infarction. *Thromb Res*. 2014;134(5):976-9.
11. Bozkurt B, Mann DL, Deswal A. Biomarkers of inflammation in heart failure. *Heart Fail Rev*. 2010;15(4):331-41.
12. Drakopoulou M, Toutouzias K, Stefanadi E, Tsiamis E, Tousoulis D, Stefanadis C. Association of inflammatory markers with angiographic severity and extent of coronary artery disease. *Atherosclerosis*. 2009;206(2):335-9.
13. Pu LJ, Lu L, Shen WF, Zhang Q, Zhang RY, Zhang JS, et al. Increased glycated albumin and decrease desrage levels are related to angiographic severity and extent of coronary artery disease in patients with type 2 diabetes. *Circ J*. 2007;71(7):1067-73.
14. Tanindi A, Sahinarslan A, Elbeg S, Cemri M. Relationship Between MMP-1, MMP-9, TIMP-1, IL-6 and Risk Factors, Clinical Presentation, Extent and Severity of Atherosclerotic Coronary Artery Disease. *Open Cardiovasc Med J*. 2011;5:110-6.
15. Semba RD, Patel KV, Ferrucci L, Sun K, Roy CN, Guralnik JM, et al. Serum antioxidants and inflammation predict red cell distribution width in older women: the Women's Health and Aging Study I. *Clin Nutr*. 2010;29(5):600-4.
16. Montagnana M, Cervellin G, Meschi T, Lippi G. The role of red blood cell distribution width in cardiovascular and thrombotic disorders. *Clin Chem Lab Med*. 2011;50(4):635-41.
17. Lippi G, Targher G, Montagnana M, Salvagno GL, Zoppini G, Guidi GC. Relation between red blood cell distribution width and inflammatory biomarkers in a large cohort of unselected outpatients. *Arch Pathol Lab Med*. 2009;133(4):628-32.
18. de Back DZ, Kostova EB, van Kraaij M, van den Berg TK, van Bruggen R. Of macrophages and red blood cells; a complex love story. *Front Physiol*. 2014;5:9.
19. Balistreri WF, Leslie MH, Cooper RA. Increased cholesterol and decreased fluidity of red cell membranes (spur cell anemia) in progressive intrahepatic cholestasis. *Pediatrics*. 1981;67(4):461-6.
20. Cooper RA, Leslie MH, Knight D, Detweiler DK. Red cell cholesterol enrichment and spur cell anemia in dogs fed a cholesterol-enriched atherogenic diet. *J Lipid Res*. 1980;21(8):1082-9.
21. Franco RS. Measurement of red cell lifespan and aging. *Transfus Med Hemother*. 2012;39(5):302-7.
22. Huang YX, Wu ZJ, Mehrishi J, Huang BT, Chen XY, Zheng XJ, et al. Human red blood cell aging: correlative changes in surface charge and cell properties. *J Cell Mol Med*. 2011;15(12):2634-42.
23. Vaya A, Sarnago A, Fuster O, Alis R, Romagnoli M. Influence of inflammatory and lipidic parameters on red blood cell distribution width in a healthy population. *Clin Hemorheol Microcirc*. 2015;59(4):379-85.
24. Aulakh R, Sohi I, Singh T, Kakkar N. Red cell distribution width (RDW) in the diagnosis of iron deficiency with microcytic hypochromic anemia. *Indian J Pediatr*. 2009;76(3):265-8.
25. Tziakas D, Chalikias G, Grapsa A, Gioka T, Tentes I, Konstantinides S. Red blood cell distribution width: a strong prognostic marker in cardiovascular disease: is associated with cholesterol content of erythrocyte membrane. *Clin Hemorheol Microcirc*. 2012;51(4):2430-54.
26. Zhong Y, Tang H, Zeng Q, Wang X, Yi G, Meng K, et al. Total cholesterol content of erythrocyte membranes is associated with the severity of coronary artery disease and the therapeutic effect of rosuvastatin. *Ups J Med Sci*. 2012;117(4):390-8.
27. Kucera M, Balaz D, Kruzliak P, Ciccocioppo R, Oravec S, Rodrigo L, et al. The effects of atorvastatin treatment on the mean platelet volume and red cell distribution width in patients with dyslipoproteinemia and comparison with plasma atherogenicity indicators--a pilot study. *Clin Biochem*. 2015;48(9):557-61.
28. Forsyth AM, Braunmuller S, Wan J, Franke T, Stone HA. The effects of membrane cholesterol and simvastatin on red blood cell deformability and ATP release. *Microvasc Res*. 2012;83(3):347-51.
29. Skjelbakken T, Lappegård J, Ellingsen TS, Barrett-Connor E, Brox J, Lochen ML, et al. Red cell distribution width is associated with incident myocardial infarction in a general population: The Tromsø Study. *J Am Heart Assoc*. 2014;3(4).
30. Hu GX, Zhang J, Tian YG, Li YH, Mou L, Qiao LJ. "Diagnostic value of joint detection of homocysteine and RDW CV on acute myocardial infarction," *Eur Rev Med Pharmacol Sci*. 2016;20(19):4124-8.
31. Bujak K, Wasilewski J, Osadnik T, Jonczyk S, Kołodziejaska A, Gierlotka M, et al. The prognostic role of red blood cell distribution width in coronary artery disease: a review of the pathophysiology. *Dis Markers*. 2015;2015:824624.
32. Uyarel H, Ergelen M, Cicek G, Kaya MG, Ayhan E, Turkkan C, et al. Red cell distribution width as a novel prognostic marker in patients undergoing primary angioplasty for acute myocardial infarction. *Coron Artery Dis*. 2011;22(3):138-44.
33. Gul M, Uyarel H, Ergelen M, Karacimen D, Ugur M, Turer A. The relationship between red blood cell distribution width and the clinical outcomes in non-ST elevation myocardial infarction and unstable angina pectoris: a 3-year follow-up. *Coron Artery Dis*. 2012;23(5):330-6.
34. Ani C, Ovbiagele B. Elevated red blood cell distribution width predicts mortality in persons with known stroke. *J Neurol Sci*. 2009;277(1-2):103-8.
35. Balta S, Demir M, Demirkol S, Arslan Z, Unlu M, Celik T. Red cell distribution width is related to stroke in patients with heart failure. *Clin Appl Thromb Hemost*. 2015;21(2):190.
36. Kaya A, Isik T, Kaya Y, Enginyurt O, Gunaydin ZY, Iscanli MD, et al. Relationship between red cell distribution width and stroke in patients with stable chronic heart failure: a propensity score matching analysis. *Clin Appl Thromb Hemost*. 2015;21(2):160-5.
37. Saliba W, Barnett-Griness O, Elias M, Rennert G. The association between red cell distribution width and stroke in patients with atrial fibrillation. *Am J Med*. 2015;128(2):192.

38. Söderholm M, Borné Y, Hedblad B, Persson M, Engström G. Red cell distribution width in relation to incidence of stroke and carotid atherosclerosis: a population-based cohort study. *PLoS One*. 2015;10(5):article e0124957.
39. Bareford D, Lucas GS, Caldwell NM, Stone PC, Baar S, Stuart J. Erythrocyte deformability in peripheral occlusive arterial disease. *J Clin Pathol*. 1985;38(2):135-9.
40. Haltmayer M, Mueller T, Luft C, Poelz W, Haidinger D. Erythrocyte mean corpuscular volume associated with severity of peripheral arterial disease: an angiographic evaluation. *Ann Vasc Surg*. 2002;16(4):474-9.
41. Mueller T, Haidinger D, Luft C, Horvath W, Poelz W, Haltmayer M. Association between erythrocyte mean corpuscular volume and peripheral arterial disease in male subjects: a case control study. *Angiology*. 2001;52(9):605-13.