



Toll-like Receptor 4: Target of Lipotoxicity and Exercise-Induced Anti-inflammatory Effect?

Sihui Ma¹ and Katsuhiko Suzuki^{2*}

¹Graduate School of Sport Sciences, Waseda University, Japan

²Faculty of Sport Sciences, Waseda University, Japan

Abstract

Toll-Like Receptors (TLRs) are a main target during inflammation and metabolic syndromes as receptors for inflammatory signaling molecules, and are named numerically from 1 to 13 in humans. It is believed that TLR4 plays a key role in exercise training-induced anti-inflammatory effects, whereas new evidence shows that TLR4 is the main receptor in fatty acid-induced inflammation. In this review, we summarized classical theories, and the new discovery in the field of fatty acid-induced inflammation through TLR4. Further, we highlight the anti-inflammation ability of exercise through regulating TLR4 as we try to acquire new clues for the treatment of metabolic diseases, and understand the regulatory effect of exercise.

Toll-Like Receptor, Adipose and Muscle Inflammation and Insulin Resistance

Toll-like receptors (TLRs) belong to a pattern-recognition receptor family, linking innate and adaptive immune responses to environmental stresses; in the context of exercise, Lipopolysaccharides (LPS) are liberated to blood during exhaustive exercise and binds with the soluble LPS binding protein (LBP) that facilitates its interaction with TLR4 to cause inflammation [1]. Whilst TLRs are mainly expressed in immune cells, they are also discovered in various kinds of cells including adipocytes, muscle cells, hepatocytes and islet β cells; all of which are important regulatory cells during metabolism, and within metabolic diseases [2-4].

Among the 10+ kinds of TLRs identified, TLR4 may be the most linked with metabolic diseases. TLR4 is an essential receptor for the recognition of exogenous pathogens such as LPS, and endogenous agonists such as free fatty acids. In macrophages, downstream responses of TLR4 activation include the myeloid differentiation primary response (MyD) 88-dependent signaling pathway, and the TIR-domain-containing adapter-inducing interferon- β (TRIF)-dependent pathway. Activating MyD88 is the main regulation pathway during which nuclear factor-kappa B (NF- κ B) is activated; thus elevated inflammatory cytokines, chemokines and cell adhesion molecules contribute to the cycle of inflammation [4,5]. Elevated levels of adipocyte TLR4 and cellular infiltration by macrophages are observed in various kinds of subjects [6,7]. It is also reported that TLR4 expression is elevated in muscle during insulin resistance [8]. On the other hand, it is reported that a loss-of-function mutation in TLR4 contributes to prevent diet-induced obesity and insulin resistance [9]. It could be concluded that enhanced TLR4 expression in adipose tissue recruits immune cell infiltration and aggravates the inflammation. Subsequently, this may lead to elevated whole-body inflammatory responses, thus inducing muscle TLR4 up-regulation, therefore contributing towards insulin resistance, and diabetes.

Lipotoxicity of Adipokines, Insulin Resistance and TLR4

Insulin resistance is closely involved with a chronically inflamed state. The concept of insulin resistance is a complex process, involving free fatty acids, various adipose-origin molecules, and hormones. Free fatty acids are a main component of lipotoxicity, although their exploration is beyond the scope of this review. Lipotoxicity is a metabolic syndrome that results from the accumulation of lipid intermediates in non-adipose tissue, leading to cellular dysfunction and possibly death. For example, lipotoxicity in skeletal muscle tissue is closely linked to decrease glucose uptake, thus contributing to insulin resistance [10]. Furthermore, palmitate, a classical kind of free fatty acid, is shown to interfere with the key transcription factor FoxO1 in hepatocytes, negating the function of insulin, thus failing to suppress hepatic gluconeogenesis and lipid synthesis in the liver [11].

OPEN ACCESS

*Correspondence:

Katsuhiko Suzuki, Waseda University,
Faculty of Sport Sciences, 359-1192
Mikajima, Tokorozawa, Japan,
E-mail: katsu.suzu@waseda.jp

Received Date: 10 May 2018

Accepted Date: 11 Jun 2018

Published Date: 26 Jul 2018

Citation:

Ma S, Suzuki K. Toll-like Receptor 4:
Target of Lipotoxicity and Exercise-
Induced Anti-inflammatory Effect?. *Ann
Nutr Food Sci.* 2018; 2(4): 1027.

Copyright © 2018 Katsuhiko
Suzuki. This is an open access
article distributed under the Creative
Commons Attribution License, which
permits unrestricted use, distribution,
and reproduction in any medium,
provided the original work is properly
cited.




Figure 1: Exercise interacting with TLR4 through regulation of bioactive properties.

The excess of free fatty acids causes lipotoxicity, whereas adipose-origin inflammatory cytokines, named adipokines, also contribute to this situation. Tumor Necrosis Factor (TNF)- α and Interleukin (IL)-6 are classical cytokines secreted by various cells, including macrophages and adipocytes during a chronic inflammatory state. They are classical ligands of TLR4, thus decreasing the translocation and activity of Glucose Transporter (GLUT)1 and GLUT4 to the cell membrane in skeletal muscle, leading to insulin resistance [12]. Interestingly, it is reported that blocking IL-6 signaling does not improve insulin resistance, despite the prevention of macrophage recruitment, which may be attributed to adipose-origin IL-6 secretion [13]. Leptin, resistin and adiponectin are classical adipokines linked to adipocyte metabolism and cytokine secretion network, generated from adipose tissue. Among these, leptin is reported to ameliorate insulin resistance in lipodystrophic mice independent of hepatocyte leptin receptors, and increases insulin sensitivity through activating the AMPK signaling pathway [14,15]. However, in obese subjects, leptin resistance usually exists. Several studies conducted recently claim that increased circulating resistin is associated with insulin resistance in type 2 diabetes mellitus [16,17]. Leptin resistance usually appears alongside insulin resistance, whilst the promotion of TLR4 mediated lipotoxicity, and inhibition of resistin is shown to reduce TLR4 signaling, thus ameliorating insulin resistance [18,19]. Adiponectin is an anti-inflammatory adipokine. It is reported that adiponectin suppresses TLR4/MyD88 signaling in a rat model, contributing to prevent insulin resistance [20]; increased resistin and TLR4 are linked with down-regulation of adiponectin, thus contributing towards insulin resistance [21]. Adipokines are closely linked to insulin resistance, whilst the function of TLR4 plays a key, but complimentary role in the above sessions.

Anti-Inflammatory and Anti-Diabetes Effect of Exercise through Regulation of TLR4

Endurance exercise has complicated interactions with the immune system and inflammation, including the dynamics of stress hormones, systematic cytokine release and neutrophil activation, which was narrated by Suzuki and colleagues, and then extended by followers [22-24]. The anti-inflammatory effects of acute and chronic exercise have been also reported in recent years [23-26]. Obesity induces a phenotypic switch in adipose tissue macrophage polarization, whilst aerobic exercise reverses this switch, partly by regulating TLR4 expression in adipocytes and macrophages, or by down-regulating the overexpression of TLR4, TLR2, MyD88 and NF- κ B [26]. On the other hand, resistance-trained elderly women displayed significantly

lowered TLR4 expression [27]. Muscular cell-origin IL-6 is reported to possess inflammatory regulation abilities; exercise with concurrent IL-6 infusion inhibits endotoxin-induced TNF- α production [28,29]. It is reported that the increase of TLR4 by ligands may decrease with the increase of the stimulation, therefore multiple exercise bouts may decrease the amount of activated TLR4 [30]. The reduction of free fatty acids in the blood by exercise may also contribute to the anti-inflammatory function (Figure 1). Reduced circulating monocytes by regular exercise is another probability for the exercise-induced anti-inflammatory phenomenon, since monocytes are one of the main cells to express TLR4 [31]. In a recent study, both acute and chronic exercise indirectly activated the leptin-AMPK-ACC signaling pathway and increased insulin sensitivity in the liver of type 2 diabetic rats through TLR4 mediation [32]. Exercise is reported to decrease resistin; therefore TLR4 may also be down-regulated by exercise. Figure 1 concisely describes the pathway how exercise is interacted with TLR4 through regulation of bioactive properties.

The effect of exercise on adiponectin is contradictory, with some reports of no influence, and other reports of increased adiponectin concentrations, despite the belief of exercise to improve insulin sensitivity [34-37]. Therefore, the interaction between TLR4, exercise, and various adipokines is an intriguing area of research, requiring further investigation [38-40].

Conclusion

The relationships between TLR4 and inflammation, and exercise and anti-inflammation are undeniably strong; TLR4 may be the key link between these relationships. The immediate function of adipokines requires in depth investigations. However, as TLR4 is extensively expressed in various cell types, tissue-specific knock-out models may be useful to study which tissue(s) and their magnitudes contribute to exercise-induced TLR-4-dependent anti-inflammation responses.

Acknowledgements

This work was partly supported by a Strategic Research Foundation at Private Universities 2015-2019 (S1511017) and the Scientific Research (A) (15H01833) from the Ministry of Education, Culture, Sports, Science and Technology of Japan. Sihui Ma received a scholarship from China Scholarship Council. Also, we would like to thank Dr. Llion Roberts, School of Allied Health Sciences & Menzies Health Institute Queensland, Griffith University, for English editing.

References

- Hung YL, Suzuki K. The pattern recognition receptors and lipopolysaccharides (LPS)-induced systemic inflammation. *International Journal of Research Studies in Medical and Health Sciences*. 2017; 2(7):1-7.
- Velloso LA, Folli F, Saad MJ. TLR4 at the crossroads of nutrients, gut microbiota, and metabolic inflammation. *Endocr Rev*. 2015;36(3):245-71.
- Ghosh S, Lertwattanarak R, Garduño Jde J, Galeana JJ, Li J, Zamarripa F, et al. Elevated muscle TLR4 expression and metabolic endotoxemia in human aging. *J Gerontol A Biol Sci Med Sci*. 2015;70(2):232-46.
- Yang L, Ma S, Wan Y, Duan S, Ye S, Du S, et al. In vitro effect of 4-pentylphenol and 3-methyl-4-nitrophenol on murine splenic lymphocyte populations and cytokine/granzyme production. *J Immunotoxicol*. 2016;13(4):548-56.
- Jiang Q, Yi M, Guo Q, Wang C, Wang H, Meng S, et al. Protective effects of polydatin on lipopolysaccharide-induced acute lung injury through TLR4-MyD88-NF- κ B pathway. *Int Immunopharmacol*. 2015;29(2):370-6.

6. Das UN. Is obesity an inflammatory condition? *Nutrition*. 2001;17(11):953-66.
7. Kim KA, Gu W, Lee IA, Joh EH, Kim DH. High fat diet-induced gut microbiota exacerbates inflammation and obesity in mice via the TLR4 signaling pathway. *PLoS One*. 2012;7(10):e47713.
8. Reyna SM, Ghosh S, Tantiwong P, Meka CS, Eagan P, Jenkinson CP, et al. Elevated toll-like receptor 4 expression and signaling in muscle from insulin-resistant subjects. *Diabetes*. 2008;57(10):2595-602.
9. Tsukumo DM, Carvalho-Filho MA, Carvalheira JB, Prada PO, Hirabara SM, Schenka AA, et al. Loss-of-function mutation in Toll-like receptor 4 prevents diet-induced obesity and insulin resistance. *Diabetes*. 2007;56(8):1986-98.
10. Brøns C, Grunnet LG. MECHANISMS IN ENDOCRINOLOGY: Skeletal muscle lipotoxicity in insulin resistance and type 2 diabetes: a causal mechanism or an innocent bystander? *Eur J Endocrinol*. 2017;176(2):R67-78.
11. Li Z, Lai ZW, Christiano R, Gazos-Lopes F, Walther TC, Farese RV. Global analyses of selective insulin resistance in hepatocytes due to palmitate lipotoxicity. *Mol Cell Proteomics*. 2018;17(5):836-49.
12. Daniele G, Mendoza RG, Winnier D, Fiorentino TV, Pengou Z, Cornell J, et al. The inflammatory status score including IL-6, TNF- α , osteopontin, fractalkine, MCP-1 and adiponectin underlies whole-body insulin resistance and hyperglycemia in type 2 diabetes mellitus. *Acta Diabetologica*. 2014;51(1): 123-31.
13. Kraakman MJ, Kammoun HL, Allen TL, Deswaerte V, Henstridge DC, Estevez E, et al. Blocking IL-6 trans-signaling prevents high-fat diet-induced adipose tissue macrophage recruitment but does not improve insulin resistance. *Cell Metab*. 2015;21(3):403-16.
14. Cortés VA, Cautivo KM, Rong S, Garg A, Horton JD, Agarwal AK. Leptin ameliorates insulin resistance and hepatic steatosis in *Agpat2*^{-/-} lipodystrophic mice independent of hepatocyte leptin receptors. *J Lipid Res*. 2014;55(2):276-88.
15. Minokoshi Y, Kim YB, Peroni OD, Fryer LG, Müller C, Carling D, et al. Leptin stimulates fatty-acid oxidation by activating AMP-activated protein kinase. *Nature*. 2002;415(6869):339-43.
16. Santilli F, Liani R, Di Fulvio P, Formoso G, Simeone P, Tripaldi R, et al. Increased circulating resistin is associated with insulin resistance, oxidative stress and platelet activation in type 2 diabetes mellitus. *Thromb Haemost*. 2016;116(12):1089-99.
17. Supák D, Melczer Z, Cseh K. Elevated serum acylated (biologically active) ghrelin and resistin levels associate with pregnancy-induced weight gain, insulin resistance and antropometric data in the fetus. *European Journal of Obstetrics and Gynecology and Reproductive Biology*. 2016;206:e111.
18. Holland WL, Bikman BT, Wang LP, Yuguang G, Sargent KM, Bulchand S, et al. Lipid-induced insulin resistance mediated by the proinflammatory receptor TLR4 requires saturated fatty acid-induced ceramide biosynthesis in mice. *J Clin Invest*. 2011;121(5):1858-70.
19. Santos SH, Andrade JM, Fernandes LR, Sinisterra RD, Sousa FB, Feltenberger JD, et al. Oral angiotensin-(1-7) prevented obesity and hepatic inflammation by inhibition of resistin/TLR4/MAPK/NF- κ B in rats fed with high-fat diet. *Peptides*. 2013;46:47-52.
20. Mandal P, Roychowdhury S, Park PH, Pratt BT, Roger T, Nagy LE. Adiponectin and heme oxygenase-1 suppress TLR4/MyD88-independent signaling in rat Kupffer cells and in mice after chronic ethanol exposure. *J Immunol*. 2010;185(8):4928-37.
21. Benomar Y, Amine H, Crépin D, Al Rifai S, Riffault L, Gertler A, et al. Central resistin/TLR4 impairs adiponectin signaling, contributing to insulin and FGF21 resistance. *Diabetes*. 2016;65(4):913-26.
22. Suzuki K, Totsuka M, Nakaji S, Yamada M, Kudoh S, Liu Q, et al. Endurance exercise causes interaction among stress hormones, cytokines, neutrophil dynamics, and muscle damage. *J Appl Physiol*. 1999;87(4):1360-7.
23. Suzuki K, Nakaji S, Yamada M, Liu Q, Kurakake S, Okamura N, et al. Impact of a competitive marathon race on systemic cytokine and neutrophil responses. *Med Sci Sports Exerc*. 2003;35(2):348-55.
24. Suzuki K. Cytokine response to exercise and its modulation. *Antioxidants*. 2018;7(1):17.
25. Petersen AMW, Pedersen BK. The anti-inflammatory effect of exercise. *J Appl Physiol*. 2005;98(4): 1154-62.
26. Kawanishi N, Yano H, Mizokami T, Suzuki K. Mechanisms of chronic inflammation improvement by exercise: focus on immune response of local tissue. *The Journal of Physical Fitness and Sports Medicine*. 2013;2(4):487-92.
27. Flynn MG, McFarlin BK, Phillips MD, Stewart LK, Timmerman KL. Toll-like receptor 4 and CD14 mRNA expression are lower in resistive exercise-trained elderly women. *J Appl Physiol*. 1985;95(5): 1833-42.
28. Starkie R, Ostrowski SR, Jauffred S, Febbraio M, Pedersen BK. Exercise and IL-6 infusion inhibit endotoxin-induced TNF- α production in humans. *FASEB J*. 2003;17(8):884-6.
29. Petersen AM, Pedersen BK. The role of IL-6 in mediating the anti-inflammatory effects of exercise. *J Physiol Pharmacol*. 2006;57:43-51.
30. Nomura F, Akashi S, Sakao Y, Sato S, Kawai T, Matsumoto M, et al. Endotoxin tolerance in mouse peritoneal macrophages correlates with down-regulation of surface toll-Like receptor 4 expression. *J Immunol*. 2000;164(7):3476-9.
31. Markofski MM, Flynn MG, Carrillo AE, Armstrong CL, Campbell WW, Sedlock DA. Resistance exercise training-induced decrease in circulating inflammatory CD14+ CD16+ monocyte percentage without weight loss in older adults. *European Journal of Applied Physiology*. 2014;114(8):1737-48.
32. Yi X, Cao S, Chang B, Zhao D, Gao H, Wan Y, et al. Effects of acute exercise and chronic exercise on the liver leptin-AMPK-ACC signaling pathway in rats with type 2 diabetes. *J Diabetes Res*. 2013;2013:946432.
33. Kadoglou NP, Perrea D, Iliadis F, Angelopoulou N, Liapis C, Alevizos M. Exercise reduces resistin and inflammatory cytokines in patients with type 2 diabetes. *Diabetes Care*. 2007;30(3):719-21.
34. Nassiss GP, Papantakou K, Skenderi K, Triandafillopoulou M, Kavouras SA, Yannakoulia M, et al. Aerobic exercise training improves insulin sensitivity without changes in body weight, body fat, adiponectin, and inflammatory markers in overweight and obese girls. *Metabolism*. 2005;54(11):1472-9.
35. Kriketos AD, Gan SK, Poynten AM, Furler SM, Chisholm DJ, Campbell LV. Exercise increases adiponectin levels and insulin sensitivity in humans. *Diabetes Care*. 2004;27(2):629-30.
36. Jürimäe J, Purge P, Jürimäe T. Adiponectin is altered after maximal exercise in highly trained male rowers. *Eur J Appl Physiol*. 2005;93(4):502-5.
37. Hulver MW, Zheng D, Tanner CJ, Houmar JA, Kraus WE, Slentz CA, et al. Adiponectin is not altered with exercise training despite enhanced insulin action. *Am J Physiol Endocrinol Metab*. 2002; 283(4):E861-E5.
38. Ma S, Huang Q, Yada K, Liu C, Suzuki K. An 8-week ketogenic low carbohydrate, high fat diet enhanced exhaustive exercise capacity in mice. *Nutrients*. 2018;10(6):673.
39. Ma S, Suzuki K. Potential application of ketogenic diet to metabolic status and exercise performance: a review. *EC Nutrition*. 2018;13(7):496-9.
40. Tanisawa K, Suzuki K, Ma S, Kondo S, Okugawa S, Higuchi M. Effects of ingestion of different amounts of carbohydrate after endurance exercise on circulating cytokines and markers of neutrophil activation. *Antioxidants*. 2018;7(4):51.