



Abrikossoff Tumor of the Stomach (Granular Cell Tumor)

Daniel Castresana¹, Mohammed Muqheet Adnan^{1*}, David Martin² and Gulshan Parasher¹

¹Department of Gastroenterology, University of New Mexico, Mexico

²Department of Pathology, University of New Mexico, Mexico

Clinical Image

A 49 year old M with h/o abdominal pain underwent an Upper Endoscopy that revealed a 1.5 cm submucosal gastric lesion on the greater curvature in the antrum. An EUS revealed a 14 x 16 mm mass arising from the muscularis propria. Core biopsies with a 22 gauge needle were performed which histologically and immunohistochemically confirmed granular cell tumor (GCT). Immunostaining for S-100 protein and CD 68 were positive, supporting the tumor's proposed Schwann cell origin. GCT are benign tumors most commonly seen in the esophagus (75%), Colon (21%) and are very rarely seen in the stomach (4%) [1,2]. Colorectal and gastric GCT's are thought to have more infiltrative growth [1]. Immunolabelling for S-100 protein, CD56, CD68, SOX-10 and Inhibin- α is usually positive [1]. GCT can become malignant if rapidly grow to a size of 4 cm or more, have increased cellular atypia or high mitotic activity defined as a Ki-67 index of 10% or more [3]. Surgical en bloc or wedge resection in the past and more commonly endoscopic submucosal dissection (ESD) has been described for both benign and malignant tumors [2-5]. Most of the gastric GCT were either surgically resected [3] or endoscopically dissected [2], we opted for continued endoscopic surveillance as the biopsies were consistent with a small benign tumor.

OPEN ACCESS

*Correspondence:

Mohammed Muqheet Adnan, Department of Gastroenterology, University of New Mexico, Mexico, E-mail: muqheetadnan@gmail.com

Received Date: 20 Jan 2017

Accepted Date: 30 Jan 2017

Published Date: 01 Feb 2017

Citation:

Castresana D, Adnan MM, Martin D, Parasher G. Abrikossoff Tumor of the Stomach (Granular Cell Tumor). *J Gastroenterol Hepatol Endosc.* 2017; 2(1): 1005.

Copyright © 2017 Adnan MM. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1: Core biopsies show a granular cell tumor involving muscular tissue (muscularis).




Figure 2: The neoplastic cells contain small hyperchromatic nuclei and abundant eosinophilic cytoplasm (left). A fragment of uninvolved gastric mucosa is also observed (right black arrow).




Figure 3: Immunohistochemical stains for S100 was positive supporting the diagnosis of granular cell tumor.

References

1. An S, Jang J, Min K, Kim MS, Park H, Park YS, et al. Granular cell tumor of the gastrointestinal tract: histologic and immunohistochemical analysis of 98 cases. *Hum Pathol.* 2015; 46: 813-819.
2. Kahng DH, Kim GH, Park DY, Jeon MS, Yi JW, Choi YY, et al. Endoscopic resection of granular cell tumors in the gastrointestinal tract: a single center experience. *Surg Endosc.* 2013; 27: 3228-3236.
3. Patti R, Almasio PL, Di Vita G. Granular cell tumor of stomach: a case report and review of literature. *World J Gastroenterol.* 2006; 12: 3442-3445.
4. Min KW, Lee KG, Han H, Jang SM, Paik SS. Gastric granular cell tumour clinically mimicking carcinoid tumour treated by endoscopic submucosal dissection. *ANZ J Surg.* 2014; 84: 985-986.
5. Pertile D, Scabini S, Romairone E, Scordamaglia R, Rimini E, Ferrando V. Gastric Abrikossoff tumor (granular cell tumor): case report. *G Chir.* 2010; 31: 433-434.