



Ear and Pubic Keloids in an Adolescent Girl

Harouna Moussa¹, Ibrahim Mamadou Abdoul Kadir^{2*}

¹Department of Dermatology, Dosso Regional Hospital, Niger

²Department of Medicine, Dosso Regional Hospital, Niger

Introduction

Keloids are frequent connective tissue tumors characterized by a significant proliferation of fibroblasts and collagen with a recurrent and extensive tendency. They can be secondary to a solution of cutaneous continuity: Surgical wound; vaccination; burn; hard inflammatory lesion. Sometimes spontaneous, especially in black people. We report here a case in a young patient.

Observation

This is a 20 year old married patient, melanoderma, with no known pathological history who presented to our department with multifocal slightly pruritic tumor lesions evolving for 4 years on the auricle and the pubis. The patient and her companions thought of a mystical disease. The interrogation revealed that these lesions occurred after an ear piercing and folliculitis lesions on the pubic area. The examination revealed a good general condition, firm non sensitive normochromic tumor lesions of variable size with smooth surface, protruding and adherent to the deep plane located on the pubis and bilaterally on the auricle. The CBC and blood glucose levels were normal. We proceeded to a complete removal of these lesions and infiltration sessions of corticosteroids (Triamcinolone) after healing every 2 weeks with periodic control of blood glucose and blood pressure. After 6 infiltration sessions, the result was spectacular, to the great satisfaction of the patient and the Dermatology Department team. No recurrence after one year of follow-up.

OPEN ACCESS

*Correspondence:

Ibrahim Mamadou Abdoul Kadir,
Department Medicine, Dosso Regional
Hospital, Niger,

E-mail: kader.ibrahim@yahoo.fr

Received Date: 17 Jun 2021

Accepted Date: 07 Jul 2021

Published Date: 09 Jul 2021

Citation:

Moussa H, Kadir IMA. Ear and
Pubic Keloids in an Adolescent Girl. *Am
J Otolaryngol Head Neck Surg.* 2021;
4(6): 1141.

Copyright © 2021 Ibrahim Mamadou
Abdoul Kadir. This is an open access
article distributed under the Creative
Commons Attribution License, which
permits unrestricted use, distribution,
and reproduction in any medium,
provided the original work is properly
cited.




Figure 1: Right earkeloids.



Figure 2: Left earkeloids.



Figure 3: Pubic keloids.



Figure 5: Keloids of the left ear after excision and infiltration.



Figure 4: Keloids of the right ear after excision and infiltration.



Figure 6: Pubic keloids after excision and infiltration.