



Effects of High Iodine Containing Low Osmolar Contrast Agent (Visipaque) on Thyroid Function Tests

Niknamian S*

Department of Defense, Violet Cancer Institute, Federal Health Professionals, USA

Abstract

Objective: Investigating high iodine containing low osmolar contrast agent (visipaque) effects on thyroid function tests and thyroid sonography characteristics.

Methods: 65 euthyroid cases and 92 controls composed the samples in baseline. Thyroid function tests, Urine Iodine Concentration (UIC) and thyroid sonography were conducted for both groups before and 1 and 3 months after angiography. Serum levels of T4, T3, T3RU, TSH, TPO-Ab and UIC were measured, and hypo/hyperthyroid is prevalence was compared between groups.

Results: Mean T3, T4 and TSH changes 1 month after angiography were insignificant in both groups ($P=0.61$, $P=0.4$ and $P=0.14$, $P=0.23$ in cases and controls for T3 and T4). Medians among cases and controls were 12.8 and 16.75 $\mu\text{g}/\text{dl}$, respectively, at baseline. These values varied to 28.45 and 15.2 $\mu\text{g}/\text{dl}$, and 12.95 and 14.2 $\mu\text{g}/\text{dl}$ one month and 3 months after angiography in case and control groups, respectively. UIC increase one month after angiography was significant among cases ($P=0.002$). TPO-Ab+ were same 3 months after angiography. Thyroid volume changes were significant among cases ($P<0.001$) and insignificant among controls ($P=0.680$). No significant difference was seen between cases and controls in overt hypothyroidism; however, a considerable change was seen in thyroid volume and UIC one month after angiography among cases. The hypothyroidism rate among cases was insignificant which may be either related to few cases or short half-life of visipaque (2.1 h) so that 97% of injected dose was excreted in urine within 24 h.

Conclusion: Thyroid function test is not recommended before angiography in patients without previous thyroid records.

Keywords: Thyroid disease; Hyperthyroidism; Thyroid function test; Visipaque

OPEN ACCESS

Introduction

***Correspondence:**

Niknamian S, Department of Defense,
Violet Cancer Institute, Federal Health
Professionals. VA, USA,
E-mail: so.niknamian@gmail.com

Received Date: 02 Jul 2020

Accepted Date: 05 Sep 2020

Published Date: 09 Sep 2020

Citation:

Niknamian S. Effects of High Iodine Containing Low Osmolar Contrast Agent (Visipaque) on Thyroid Function Tests. *Ann Integr Oncol.* 2020; 1(1): 1005.

Copyright © 2020 Niknamian S. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Thyroid diseases are the second most common cause of endocrine disorders following diabetes [1,2]. Coronary angiography and angioplasty are important diagnostic and therapeutic procedures for the coronary artery diseases. Iodine containing contrasts which is used in angiography may lead to thyroid dysfunction due to high contents of iodine. Thyroid dysfunction can increase symptomatic coronary artery diseases and cardiovascular mortality [3]. Although, another study showed that administration of iodine containing contrast agents which is usually used for coronary angiography on following thyroid hormone concentrations is not accompanied with hyperthyroidism in patients with normal results of thyroid analysis [4]. Medications containing iodine mostly induce thyroid dysfunction. Usual sources of iodine such as antiarrhythmic drugs (amiodarone), eye drops and eye ointments, topical ointments, tooth pastes and multivitamins are disinfectants [5,6]. Although, a case study demonstrated that consumption of high-dose drugs containing iodine can be considered as a predisposed to thyroid dysfunction [7]. Iodine containing radio-contrast is the most conventional compounds used in medicine [8]. For example, increased ischemic heart diseases need to cardiac catheterization and CT scans lead to increase in consumption of iodine containing contrasts [9]. A single-dose injection for CT-scan releases 7000 μg iodine which composes almost 45 folds of needed daily iodine [2]. Iodine overload condition can lead to decrease the hormone synthesis in thyroid gland by down regulation of Na/I co-transporters [10]. However, high amount of iodine intake can impair thyroid regulation and induce thyroid dysfunction [11]. Iodine containing contrasts includes low-osmolar agents, isosmolar agents and high-osmolar agents. The isosmolar and hypo-osmolar agents are suitable for patients with chronic kidney diseases, although they have higher iodine levels than high-osmolar agents [12].




Figure 1: Data of people enrollment.

One study has reported that after radio contrast intake, iodine thyroid pool in normal thyroid remains high up to 4 to 8 weeks and urinary iodine increases up to 300% and returns to normal level after 43 days [12]. On average, each patient receives about 50cc to 100cc and 30cc iodine containing contrast in CT-scan and angiography, respectively [13]. In angiographic centers, 50cc and 30cc of visipaque (Iodixanol) is typically used for CT-angiography and angiography, respectively, as isosmolar contrast agent with iodine content of 320 mg/ml [1].

We investigated the thyroid function tests, thyroid autoimmunity marker (TPOAb), thyroid ultrasonography and urinary iodine patients undertaking coronary angiography, CT-angiography in Iran, an iodine sufficient area according to WHO reports [12].

Materials and Methods

Patients

This study was performed at angiographic center of Isfahan Endocrine and Metabolism Research Center in 2016-2017. Patients were selected from candidates for coronary angiographic or coronary CT angiography based on continuous enrollment. Inclusion criteria were age over 18 years old, candidate for elective angiography or CT angiography, euthyroid subjects and patient's consent for study. Our study was approved by ethics committee of Isfahan University of Medical Sciences. The medical records include history of previous thyroid diseases or malignancy or taking any medications. Accordingly, history of other disorders such as diabetes, hypertension and dyslipidemia was evaluated. Then, thyroid examination and baseline laboratory tests- including blood and urine samples-were conducted. Next, the thyroid sonography was carried out to evaluate the thyroid and neck soft tissue. The next day, angiography or CT-angiography was done for the case group. The patients were explained by the symptoms of hyperthyroid is more hypothyroid is and they were asked to report any symptoms of hyper/hypothyroidism. The same physical examinations, laboratory tests and thyroid sonography were repeated at 1 and 3 months after angiography or CT-angiography in case and control groups. Data of people enrollment is shown in

consort diagram (Figure 1).

Laboratory studies

The Fasting Plasma Glucose (FPG), lipid profile includes (TG, Cholesterol, LDL-c, HDL-c), Blood Urea Nitrogen (BUN), Creatinine (Cr) TSH, T4, T3, T3RU, Thyroid Peroxidase Antibody (TPO-Ab), were determined before coronary angiography.

Analytic approach

Statistical analysis was performed using SPSS version 24. Descriptive data was presented based on mean, Standard Deviation (SD), frequency, percentage and qualitative data using Chi-square, independent sample T-test and one way-ANOVA for comparing quantitative data between the two groups. ANOVA test was repeated for comparing changes in variables over time within each group or between groups. Median (range) was used for the variables without normal distribution such as UIC. Also, for evaluation of some parameters such as TSH and TPO-Ab that have no normal distribution, we used log-transformation.

Results

One hundred patients under the angiography or CT coronary angiography and 100 non-cardiovascular patients were enrolled. During the three month period of follow up, 35 patients in the case group were excluded (Figure 1) and data analysis was done on 65 patients and 92 controls who completed the study. The mean age of case and control groups members was 60.1 and 59.6 years, respectively ($p=0.81$); 38 (58.5%) of cases and 48 (52.2%) of controls were male ($p=0.44$). Demographic characteristics are presented in Table 1.

T3 mean at baseline was 1.65 (0.63) nmol/l and 1.7 (0.31) nmol/l in case and control groups, respectively; it was 1.82 (0.48) nmol/l and 1.78 (0.33) nmol/l in case and control groups, respectively, one month after angiography; and it was 1.65 (0.5) nmol/l and 1.81 (0.13) nmol/l for case and controls groups three months after angiography; as a result, T3 changes were statistically insignificant in both groups ($p=0.61$ and $p=0.4$ in case and control groups). Furthermore, T4 at baseline was 9.11 (2.26) μ g/dl and 8.78 (1.58) μ g/dl in case and control groups; it varied to 9.15 (2.16) μ g/dl and 8.62 (1.48) μ g/dl 1 month after angiography and also 8.76 (1.89) μ g/dl and 8.83 (6.15) μ g/dl three months after angiography in case and control groups respectively, indicating no significant change ($P=0.07$ and $P=0.8$ for case and control groups). Baseline TSH among cases and controls was 2.84 (2.1) μ u/ml and 3.15 (2.95) μ u/ml, respectively; it varied to 2.98 (1.06) μ u/ml and 3.22 (2.39) μ u/ml 1 month later and 4.14 (1.5) μ u/ml and 3.49 (2.58) μ u/ml 3 months later in case and control groups, respectively, indicating no significant TSH changes ($P=0.14$ and $P=0.23$ among cases controls) (Table 2).

The mean of serum TSH, T4, T3 and TPO Ab concentrations in different thyroid functions are presented in (Tables 3a and 3b). Mean of serum TSH level was statistically increased in one month after study in both groups but other functional thyroid hormones (T3 and T4) were not different among exposed and non-exposed groups. Moreover, 14 cases and 28 controls were become hypothyroid (mild and overt hypothyroidism) and no statistically significant difference was seen between the two groups ($p=0.07$). Also the mean level of TSH was not different between the case and control groups in 1 and 3 months after intervention. Distribution of TSH level (including range, median and 25 to 75 percentiles) in 1 and 3 months after intervention are shown in Figure 2.

Table 1: Demographic characteristics and baseline values of patients under angiography and control groups.

Variables	Groups	Case	Control	P
Sex	Male	38 (58.5)	48 (52.2)	0.44
N (%)	female	27 (41.5)	44 (47.8)	
age(year)		60.1 (\pm 10.3)	59.6 (\pm 14)	0.81
BMI (kg/m ²)		26.88 (\pm 3.06)	26.74 (\pm 3.94)	0.81
Family history of thyroid disorders		12 (18.5)	27 (29.3)	0.12
History of radiotherapy of head and neck		3 (4.6)	1 (1.1)	0.31
Thyroid function tests at base line	T4 (μg/dl)	9.11 (\pm 2.26)	8.78 (\pm 1.58)	0.54
	T3(μg/dl)	1.65 (\pm 0.63)	1.7 (\pm 0.31)	0.28
	T3RU	0.9 (\pm 0.2)	0.85 (\pm 0.12)	0.056
	TSH (mIU)	2.84 (\pm 2.1)	3.15 (\pm 2.95)	0.68
	TPO Ab +	14 (21.5%)	29 (31.5%)	0.65
	UIC (μg/dl)	12.43 (\pm 7.55)	13.54 (\pm 8.64)	0.44

T3: Three Iodothyronin, T4: Thyroxin, TSH: Thyroid Stimulating Hormone, T3RU: T3 Resin Uptake, UIC: Urinary Iodine Concentration

'P value <0.05


TPO Ab thyroid peroxidase antibody, as considered positive when level >37 IU/mI

Table 2: The mean changes of thyroid function test in the before and after angiography in the two groups.

Variables	Groups	Baseline	1 st month	3 rd months	P
T4 (μg/dl)	Case	9.11 (2.26)	9.15 (2.16)	8.76 (1.89)	0.07
	control	8.78 (1.58)	8.62 (1.48)	8.83 (6.15)	0.8
T3 (nmol/l)	Case	1.65 (0.63)	1.82 (0.48)	1.65 (0.5)	0.61
	control	1.7 (0.31)	1.78 (0.33)	1.81 (.13)	0.4
T3RU (μu/ml)	Case	2.84 (2.1)	2.98 (1.06)	4.14 (1.5)	0.14
	control	3.15 (2.95)	3.22 (2.39)	3.49 (2.58)	0.23
TSH median (range)	Case	1.7 (1.2-3)	1.8 (1.2-3.35)	2.3 (1.5-3.2)	0.027*
	control	2.3 (1.43-3.1)	2.5 (1.93-3.5)	2.55 (1.8-4.2)	0.009*


*mean \pm SD; *median (percentile 25 to 75); T3: Three Iodothyronin; T4: Thyroxin; TSH: Thyroid Stimulating Hormone

'P value <0.05

**Figure 2:** Distribution of TSH level (including range, median and 25 to 75 percentile) in 1 and 3 months after intervention.

In addition, frequency of TPO-Ab+ in case and control group was 14 and 29 (21.5% vs. 31.5), respectively, and no statistically difference was found between the two groups (P=0.17).

Median of urinary iodine was not different between exposed and non-exposed groups at baseline (P=0.93) as well as 1 and 3 months after intervention (Figure 2). Median of urinary iodine before exposure was 12.8 μg/dl and 16.75 μg/dl among cases and controls, respectively, indicating no statistically significant difference between

**Figure 3:** Median, range and 25 to 75 percentile of TSH level in hypothyroid patients in the case and control groups.

two groups (p=0.059). Median of urinary iodine one month after exposure was 28.45 μg/dl and 15.2 μg/dl among cases and controls, respectively, indicating a statistically significant difference between the two groups (p=0.002); while it was 12.95 μg/dl and 14.2 μg/dl among cases and controls, respectively, 3 months after exposure, indicating no statistically significant difference between the two group (p=0.13) (Figure 3). Mean value of thyroid volume in case group at baseline was 5.908 (1.36 to 8.8) cc which increased to 6.244.7 (1.039.6) cc one month later and decreased to 5.630 (1.323) cc three

Table 3a: Thyroid function tests 1 month after exposure.

Scale	Case group				Control group			
	Euthyroid (n=51)	Mild Hypo (n=10)	Overt-hypo (n=4)	P	Euthyroid (n=64)	Mild Hypo (n=24)	Overt-hypo (n=4)	P
T3 (nmol/l)	1.8 (0.4)	1.7 (.55)	1.9 (51)	0.74	1.8 (0.31)	1.4 (0.35)	1.35 (.35)	0.14
T4 (μg/dl)	9.8 (6.7)	9.5 (5.5)	13.8 (5.9)	0.12	9.1 (3.9)	8.8 (6.5)	9.04 (6.3)	0.06
TSH (μu/l)	1.8 (0.8)	8.2 (1.6)	13.7 (2.5)	<0.001	2.3 (0.6)	5.6 (1.7)	14.6 (0.8)	<0.001
T3RU	0.96 (0.1)	1.03 (0.3)	0.7 (0.1)	0.08	0.9 (0.1)	0.92 (0.2)	0.7 (0.001)	0.12
N (%) TPO+	10 (19.6%)	3 (30%)	1 (25%)	0.66	20 (31%)	8 (33.3%)	1 (2%)	0.43
UIC (μg/dl)	29.68 (14.2)	26.93 (17.6)	27.9 (3.3)	0.93	29.4 (14.2)	18.6 (10.9)	14.7 (9.7)	0.44

T3: Three Iodothyronine; T4: Thyroxin; TSH: Thyroid Stimulating Hormone; TPOAb: Thyroid Peroxidase Antibody as considered positive when level >37 IU/Ml, T3RU:

T3 Resin Uptake; UIC: Urinary Iodine Concentration

*P value <0.05

Table 3b: Thyroid function tests 3 month after exposure.

Scale	Case group				Control group			
	Euthyroid (n=51)	Mild Hypo (n=10)	Overt-hypo (n=4)	P	Euthyroid (n=64)	Mild Hypo (n=24)	Overt-hypo (n=4)	P
T3	1.67 (0.48)	1.58 (0.48)	1.6 (0.26)	0.82	1.7 (0.5)	2.1 (2)	1.5 (0.27)	0.28
T4	9.2 (1.8)	7.3 (1)	6.6 (0.5)	<0.1	8.3 (1.7)	10.6 (2.9)	7.5 (1.5)	0.27
TSH	1.95 (0.8)	6.2 (1.7)	27 (8.6)	<0.001	2.2 (0.7)	5.2 (1.5)	14.2 (2.1)	<0.001
TTRU	0.93 (0.1)	0.88 (0.15)	0.85 (0.1)	0.19	0.9 (0.1)	0.9 (0.2)	0.87 (0.1)	0.95
N (%) TPO +	11 (21.6%)	3 (30%)	1 (25%)	0.84	18 (28.1%)	8 (33.3%)	2 (50%)	0.61
UIC	17.8 (11.7)	7.1 (2.1)	17.4 (11.7)	0.38	13 (5)	14.1 (5.7)	17.8 (10.1)	0.57

T3: Three Iodothyronine; T4: Thyroxin; TSH: Thyroid Stimulating Hormone; TPO-Ab: Thyroid Peroxidase Antibody, as considered positive when level >37 IU/Ml, T3RU:

T3 Resin Uptake; UIC: Urinary Iodine Concentration

*P value <0.05

Table 4: Mean changes of thyroid volume (mm³) in the two groups.

Time	Groups	
	Case	control
Baseline	5908.6 ± 1368.8	5611.9 ± 1477.6
1 month later	6244.7 ± 1039.6	5662 ± 1394
3 months later	5630 ± 1323	5639.9 ± 1403
P	<0.001	0.68

months later. Thyroid volume changes in case group were statistically significant ($P<0.001$) but they were insignificant among the controls ($P=0.680$) cc. Similarly, mean thyroid volume at baseline was 5.611.9 (1.477.6) cc and it was 5.662 (1.394) cc and 5.639 (1.403) cc one and three months later, respectively, respectively (Tables 4 and 5).

Discussion

The patients with cardiovascular diseases usually considerably need angiography and angioplasty for management of coronary disease, which lead to exposure to iodine containing contrast media and cause some problems for such patients such as congestive heart failure and cardiac arrhythmia [2]. It is also reported that contrast agents increase thyroid dysfunction such as hyperthyroidism and hypothyroidism, and cardiopulmonary arrest after radio-contrasts injection [14,15]. The result from a large epidemiological study in Iran showed this country as an endemic iodine deficient area [16].

Our results showed the mean urine iodine as 20/3 mg and 20/1 mg/d for Euthyroid and hypothyroid subjects, respectively, and in a study conducted in Isfahan, the UIC median for iodine sufficiency is reported 18 μg/dl [16,17]. It is also shown that contrast Media are independent risk factors for mortality and cardiovascular events

Table 5: Mean thyroid volume (CC) based on thyroid status in case and control groups in 1 and 3 months after exposure.

Time	Thyroid volume	Groups	
		case	control
One month after	Normal	6.403 (0950)	5675.7 (1353.3)
	Mild hypo	5.725 (1018)	5515.8 (1947.6)
	Overt	4.326.9 (383.4)	5603 (1929)
three months after	P	0.001	0.96
	Normal	5.762.3 (1334.9)	6529.3 (1364.7)
	Mild hypo	5.375.2 (795.1)	5873.6 (1492.1)
Overt	4.092.4 (493.1)	5046 (1813.9)	
	P	0.044	0.62

[18]. Also, in a prospective cohort conducted on 101 euthyroid patients, hyperthyroidism was arisen among such these patients [19]. Additionally, evaluation of thyroid function tests 1,3,6,12 months after angiography indicate several new cases of hyperthyroidism where severity of baseline thyroid dysfunction is increased [20]. Some studies found no finding about any effect or significant change in thyroid function tests [17]. Due to different results of studies and lack of similar study in Iran, we decided to evaluate effect of contrast agents on thyroid function tests as our main goal. Previous studies have not investigated thyroid sonographic features and UIC and their association with thyroid dysfunction. In our study, no significant difference was found in terms of mean changes of T4 ($P=0.47$), total T3 ($P=0.7$) and TSH ($P=0.95$) between the two groups. Median of UIC was 12.8 μg/dl and 16.75 μg/dl among cases and controls, respectively, at baseline; and it increased to 28.45 μg/dl and 15.2 μg/dl and also 12.95 μg/dl and 14.2 μg/dl one and three months after intervention,

respectively, among cases in controls, indicating a statistically significant change after one month ($P=0.002$). Mean values of TPO-Ab+ three months after angiography were same as the baseline. Thyroid volume changes in case group were statistically significant ($P<0.001$) while they were insignificant among controls ($P=0.680$). There was no significant difference between case and control group in overt hypothyroidism while a significant difference was seen in terms of thyroid volume and size and UIC one month after angiography in both groups. Consequently, the use of iodine containing contrast media in angiography in cardiovascular diseases may be associated with thyroid dysfunction and cardiac dysfunctions such as cardiac arrhythmia among such patients; this is why we were motivated for this study. Among the patients, we found some insignificant changes in thyroid volume and mean TSH level among members of exposed group; therefore, follow up of thyroid function tests among euthyroid patients is not recommended.

Conclusion

In conclusion, thyroid volume changes were significant among cases ($P<0.001$) and insignificant among controls ($P=0.680$). No significant difference was seen between cases and controls in overt hypothyroidism, however, a considerable change were seen in thyroid volume and UIC one month after angiography among cases. The hypothyroidism rate among cases was insignificant which may be either related to few cases or short half-life of visipaque (2.1 h).

References

- Aminorroaya A, Meamar R, Amini M, Feizi A, Tabatabaei A, Imani EF. Incidence of thyroid dysfunction in an Iranian adult population: The predictor role of thyroid auto antibodies: Results from a prospective population-based cohort study. *Eur J Med Res.* 2017;22(1):21.
- Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. The Colorado thyroid disease prevalence study. *Arch Intern Med.* 2000;160(4):526-34.
- Cappola AR, Fried LP, Arnold AM, Danese MD, Kuller LH, Burke GL, et al. Thyroid status, cardiovascular risk, and mortality in older adults. *JAMA.* 2006;295(9):1033-41.
- Koroscil MT, Pelletier PR, Slauson JW, Hennessey J. Short-term effects of coronary angiographic contrast agents on thyroid function. *Endocr Pract.* 1997;3(4):219-21.
- Thomsen HS, Faber J. Iodine-based contrast media cause hyperthyroidism—another important adverse reaction to contrast media? SAGE Publications. 2012;53(10).
- Seminara M, Stephanie B, Daniels M, FACE, Gilbert H. Amiodarone and the thyroid. *Endocr Pract.* 1998;4(1):48-57.
- Hoang T, Mai V, Clyde P, Shakir M. Over-the-counter-drug-induced thyroid disorders. *Endocr Pract.* 2013;19(2):268-74.
- Toprak O, Cirit M, Bayata S, Yeşil M. Review of the radiocontrast nephropathy risk profiles and risk stratification. *Anatol J Cardiol.* 2004;4(4):331-5.
- Katzberg R, Haller C. Contrast-induced nephrotoxicity: Clinical landscape. *Kidney Int.* 2006;69:S3-S7.
- Eng PH, Cardona GR, Fang S-L, Previt M, Alex S, Carrasco N, et al. Escape from the acute Wolff-Chaikoff effect is associated with a decrease in thyroid sodium/iodide symporter messenger ribonucleic acid and protein. *Endocrinology.* 1999;140(8):3404-10.
- Zimmermann MB, Aeberli I, Torresani T, Bürgi H. Increasing the iodine concentration in the Swiss iodized salt program markedly improved iodine status in pregnant women and children: A 5-y prospective national study. *Am J Clin Nutr.* 2005;82(2):388-92.
- Lee SY, Rhee CM, Leung AM, Braverman LE, Brent GA, Pearce EN. A review: Radiographic iodinated contrast media-induced thyroid dysfunction. *J Clin Endocrinol Metab.* 2015;100(2):376-83.
- Mann DL, Felker GM. Heart Failure: A companion to Braunwald's heart disease: Elsevier Health Sciences; 2014.
- Kornelius E, Chiou J-Y, Yang Y-S, Peng C-H, Lai Y-R, Huang C-N. Iodinated contrast media increased the risk of thyroid dysfunction: A 6-year retrospective cohort study. *J Clin Endocrinol Metab.* 2015;100(9):3372-9.
- Alkuju S, Pyram R, Odeyemi O. In the eye of the storm: Iodinated contrast medium induced thyroid storm presenting as cardiopulmonary arrest. *Heart & Lung: J Acute Crit Care.* 2013;42(4):267-9.
- Azizi F, Hatami H, Janghorbani M. Epidemiology and control of common diseases in Iran. Tehran: Eshtiagh Publications. 2000;602-16.
- Molen AJVD, Thomsen HS, Morcos SK. Contrast media safety committee, European Society of Urogenital Radiology. Effect of iodinated contrast media on thyroid function in adults. *Eur Radiol.* 2004;14(5):902-7.
- Marraccini P, Bianchi M, Bottoni A, Mazzarisi A, Coceani M, Molinaro S, et al. Prevalence of thyroid dysfunction and effect of contrast medium on thyroid metabolism in cardiac patients undergoing coronary angiography. *Acta Radiologica.* 2013;54(1):42-7.
- Ozkan S, Oysu AS, Kayatas K, Demirtunç R, Eren M, Uslu H, et al. Thyroid functions after contrast agent administration for coronary angiography: A prospective observational study in euthyroid patients. *Anadolu Kardiyol Derg.* 2013;13(4):363-9.
- Egorov V, Sviridenko N, Platonova N, Arbuzova M, Zlotnikova O, Buziashvili I, et al. Disturbances of thyroid function after coronary angiography. *Kardiologiiia.* 2005;46(4):46-9.