



## Gustatory Changes Following Mandibular Third Molar Surgery: A Systematic Review of the Literature

Abdurrahman A Al-Samman\* and Omar S Al-Nuaim

Department of Oral Surgery, Ninawah Health Directorate, Iraq

### Abstract

The purpose of this study was to systematically review studies to measure the incidence of gustatory changes because of LN damage following Mandibular Third Molar (M3M) removal. A computerized search of the several databases and references cited in the various studies was performed to identify eligible articles. The primary predictor variable was taste changes after M3M surgery. Five published studies acceptable for detailed analysis according to predetermined inclusion and exclusion criteria. In the 5 selected articles, incidences of gustatory changes were 0% to 38.3% and 2.9% of the pooled studies. Taste disturbance as a result LN deficit in M3M surgery in not uncommon complication especially in patients with high difficulty index score.

**Keywords:** Wisdom tooth surgery; Taste disturbance; Gustatory deficit; Neuro-sensory deficit

### Introduction

The removal of Mandibular Third Molars (M3M) is the most common surgical procedure associated with Lingual Nerve (LN) damage [1,2].

LN is anatomically related to third molar area; within the periosteum in the lingual or distal side of third molar; that makes it at risk when lingual flap is reflected, lingual flap retractor is used, and/or when tooth extracted using lingual split technique [3-5]. Moreover, the anatomic variation on LN and the fact that panoramic radiography cannot preoperatively predicts of LN injury, the surgeon is often not able to avoid this complication [6].

LN injury may result in a variety of neurosensory deficits like anesthesia, paresthesia, dysesthesia, hypoesthesia and/or a change in taste perception of food and drink (hypogeusia or ageusia) [7,8]. It is very disabling complication causing many problems for patients like tongue biting, drooling, burns from hot food and drinks, and a burning sensation of the tongue [7].

This systematic review aimed to identify all available relevant studies in the literature related to gustatory changes because of LN damage in M3M surgery, and to determine the incidence and time course of these changes.

### Materials and Methods

Electronic search, including PubMed, Scopus, and the Cochrane Library were performed. In addition; the online databases of the Journal of Oral and Maxillofacial Surgery; International Journal of Oral and Maxillofacial Surgery; British Journal of Oral and Maxillofacial Surgery; Oral Surgery; Oral Medicine; Oral Pathology; Oral Radiology as well. Search performed for all years available up until 10<sup>th</sup> July 2018. Only articles published in English were included. The following key words/terms were used in search: third molar or wisdom tooth combined with paresthesia, dysesthesia, taste, gustatory function, neurosensory deficit, neurosensory disturbance and sensory nerve impairment.

The authors carried out the search individually. The abstract of each article was viewed and the full text was downloaded if it was considered relevant to the inclusion criteria of the study:

- Articles relevant to lingual nerve deficit caused by M3M surgery;
- Articles list the gustatory deficit that the subjects experienced based on clinical objective taste testing.
- Articles must be randomized clinical trials, controlled clinical trials, or prospective clinical studies.

### OPEN ACCESS

#### \*Correspondence:

Abdurrahman A Al-Samman,  
Department of Oral Surgery,  
Ninawah Health Directorate, Iraq, Tel:  
00964772 1933236;  
E-mail: abd\_alsamman@yahoo.com

Received Date: 12 Nov 2018

Accepted Date: 22 Nov 2018

Published Date: 27 Nov 2018

#### Citation:

Al-Samman AA, Al-Nuaim OS.  
Gustatory Changes Following  
Mandibular Third Molar Surgery: A  
Systematic Review of the Literature.  
World J Oral Maxillofac Surg. 2018;  
1(3): 1015.

**Copyright** © 2018 Abdurrahman A Al-Samman. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Table 1:** Articles for the final review and the incidence of taste disturbance and other LN deficits presented in these articles.

Study	Study design	Number of M3M surgeries	Age of patients, Mean (SD)	Type of surgery	Number (%) of patients with taste disturbance	Number (%) of patients with other LN deficits	Neuro-sensory evaluation
Shafer et al., [12]	Pros.	17 patient (bilateral) N=34	15 years to 28 years, 21.1 year (3.7)	Standard buccal approach	1 (2.9%) patient with temporary deficit	1 (2.9%) patient with temporary deficit	Preop, 1 and 6 months after surgery
Gulicher and Gerlach [13]	Pros.	687 patient, N=1106	About 25.8 year	Standard buccal approach With lingual periosteal elevator to protect LN	13 (1.2%) patients with temporary Deficit 1 (0.09%) of them permanent deficit.	23 (2.1%) patients with temporary deficit 3 (0.3%) of them permanent deficit.	After surgery 6 months
Akal et al., [14]	Pros.	27 patient (bilateral) N=54	17 years to 28 years, 21.32 year (3.6)	Standard buccal approach	0	0	Preop, 1 and 6 months after surgery
Ridaura-Ruiz et al., [15]	Pros.	16	≥ 55 years, 29.9 year (11.2)	Standard buccal approach	0 (only salt taste tested)	0	Preop., 1 week, and 1 month after surgery
Anand et al., [16]	RCT	60	NA	Standard buccal approach	23 (38.3%)	Not assessed	Preop. and 1 week after surgery
<b>Total</b>		<b>1270</b>			<b>37 (2.9%)</b>	<b>*24 (2%)</b>	

M3M: Mandibular third Molar

LN: Lingual Nerve

NA: Not Available

\*Results of only 4 studies

• Studies were also selected if contain data of interest among other findings.

A second-round search was carried out using the references of all included articles that met the inclusion criteria. Any disagreement between authors solved by consensus. Due to a limited number of studies, and the considerable differences in study design, it was not possible to determine the risk of bias of included studies.

Then, the variable of interest in the results of all selected studies were analyzed as “mixed group” of studies to assess taste deficits among patients undergoing M3M extraction.

## Results

A computer searches resulted in 191 potentially relevant articles. A review of the reference lists in these articles yielded another 23 articles, giving a total of 214 articles.

In the first round of evaluation, 206 articles were excluded from the study: eighty-two articles only assessed paresthesia and/or dysesthesia of LN. Twenty-five articles were review, case reports, case-series, and technical notes. Twenty-five articles related to mental/inferior alveolar nerve deficit. Twenty-three duplicated articles. Twenty-one articles were not related to wisdom tooth surgery. Eleven articles were published in non-English language. Seven articles assessed taste function of LN after repair surgery or maxillofacial trauma. Abstract not available in 6 articles. Five articles were letter to editor, and one animal study.

Only eight articles considered eligible and passed through the second round of evaluation. In this round, three articles were excluded; two retrospective studies evaluated taste function in subjective way, and correlate patients with taste deficits with neurosensory dysfunction [9,10]. Another article assessed taste function subjectively 3 months to 12 months after surgery [11].

Five studied were suitable for final review [12-16]. One of the selected studies was randomized clinical trials and 4 were prospective clinical studies (Figure 1 and Table 1).

All studies carried out using the standard buccal approach (no chisel, no lingual split technique, and no lingual flap reflection) and in

one of them, the periosteal elevator used as LN protector [13].

The incidence of temporary taste dysfunction in the included studies ranged from 0% to 38.3%, and the combined totals were 2.9%. In contrast, the reported rates of other forms of LN deficits; in four of included studies; were 0% to 2.1%; giving a total of 2%.

## Discussion

The LN carries different types of nerve fibers, and joined by the chorda tympani branch of the facial nerve, which contains gustatory, thermosensitive, and mechanosensitive afferent fibers from papillae on the dorsal surface of the tongue [17]. The Chorda tympani-lingual nerve runs medially to the mandible at third molar region where they being most susceptible to damage during surgical procedures [14].


Thorough knowledge of LN anatomy is crucial to avoid or reduce the risk of iatrogenic nerve damage, it is important to know that LN spatial position differ widely from individual to individual, and in the same individual [18]. Also, the level of alveolar process and the inclination of its lingual surface in the M3M region can influence the LN position [19].

Trauma to the LN cannot be predicted by preoperative panoramic radiography, and different methods have been used to evaluate LN spatial position, such as magnetic resonance imaging, ultrasound, but they did not accurately identify and assess anatomic relations of LN to the neighboring structures [20-23].

Knowledge about the incidence of LN disturbances is particularly important for the surgeon as well as for the patient when evaluating the risk-benefit ratio in M3M surgery [24].

The rate of lingual nerve temporary deficit following M3M surgery ranges 0.1% to 22% [25]. This wide range due to different risk factors like increasing age, M3M eruption status, pattern of impaction, type of surgical technique, raising of lingual flap, and surgeons' experience. LN may also be damaged as a result of nerve compression by edema, by syringe needle, local anesthetics, and suture at the 3M extraction site [14,20,26,27].

The postoperative evaluation of sensory disturbance is necessary to evaluate the type of the problem and to test the over time-potential



improvement [7,28]. The sense of taste could be surveyed objectively by using a kit containing different concentrations of sugar (sucrose), salt (NaCl), citric acid, and quinine hydrochloride.

The sensory tests should be as objectively as possible to avoid patients' bias in subjective reports that may influence results [29,30]. This explains why reviewers excluded three studies from this review [9-11].

Although Anand et al., [16] in their study not assessed other forms of LN deficits; they reported a highest incidence (38.3%) of taste changes compared with other studies, especially in cases with high extraction difficulty score that need a high level of surgeon's experience. Similarly, researchers found a highly significant correlation between the level of M3M extraction difficulty and incidence of LN deficit [31]. Moreover, Hillerup and Stoltze recorded a comparable result of taste deficit (32.6%) when they assessed taste function subjectively [11].

Akal et al., [14] tested the whole-mouth gustatory function and found non-significant increase in thresholds for all taste sensations compared with preoperative values. The non-documentation of surgical difficulty score and the type of impaction might explain these findings. Similar findings were reported by Ridaura-Ruiz et al., [15]. They observed only slight increase in the salt recognition threshold one week after surgery, which was not significant statistically. Noting that this study comprised limited number of cases and it only consider the salt taste recognition as a sign of gustatory deficit. However salty tastes have more predilections for increased taste threshold compared with other tastes as reported by other studies [11,14,15].

Gülicher and Gerlach surveyed postoperative sensory impairment over a period of 6 months in more than 680 patients with more than 1100 case of M3M surgery [13]. They reported 1.2% and 0.09% temporary and permanent hypoguesia respectively. Surgeons routinely inserted periosteal elevator on the lingual aspect of the mandible to protect LN and this may accounts for a higher percent

(2.1%) of other types of LN sensory disturbance were recorded illustrating that not all cases of LN paresthesia or dysesthesia should associated with taste deficit.

Differences among included studies may be related to the fact that that nerve fiber size and myelination may play a role in neurosensory disturbances after LN injury [32].

Assessment of gustatory deficit and recovery was not attempted largely in M3M studies. For future studies, the authors recommend a standardized methodology and follow- up for assessing gustatory deficit following third molar surgery through randomized clinical trials with large sample size to evaluate incidence, type of this neurosensory disturbance and the associated risk factors.

## Conclusion

This literature review of prospective studies found that taste disturbance as a result LN deficit in M3M surgery in not uncommon complication especially patients with high difficulty index score.

## References

1. Bagheri SC, Meyer RA, Khan HA, Kuhmichel A, Steed MB. Retrospective review of microsurgical repair of 222 lingual nerve injuries. *J Oral Maxillofac Surg.* 2010;68(4):715-23.
2. Tay AB, Zuniga JR. Clinical characteristics of trigeminal nerve injury referrals to a university centre. *Int J Oral Maxillofac Surg.* 2007;36(10):922-7.
3. Pogrel MA, Renaut A, Schmidt B, Ammar A. The relationship of the lingual nerve to the mandibular third molar region: an anatomic study. *J Oral Maxillofac Surg.* 1995;53(10):1178-81.
4. Brann CR, Brickley MR, Shepherd JP. Factors influencing nerve damage during lower third molar surgery. *Br Dent J.* 1999;186(10):514-6.
5. Rood JP. Permanent damage to inferior alveolar and lingual nerves during the removal of impacted mandibular third molars. Comparison of two methods of bone removal. *Br Dent J.* 1992;172(3):108-10.

6. Chossegros C, Guyot L, Cheynet F, Belloni D, Blanc JL. Is lingual nerve protection necessary for lower third molar germeotomy? A prospective study of 300 procedures. *Int J Oral Maxillofac Surg.* 2002;31(6):620-4.
7. Fielding AF, Rachiele DP, Frazier G. Lingual nerve paresthesia following third molar surgery: a retrospective clinical study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1997;84(4):345-8.
8. Jones RH. Microsurgical repair of nerves injured during third molar surgery. *Aust Dent J.* 1992;37(4):253-61.
9. Hillerup S. Iatrogenic injury to oral branches of the trigeminal nerve: records of 449 cases. *Clin Oral Investig.* 2007;11(2):133-42.
10. Kjølle GK, Bjørnland T. Low risk of neurosensory dysfunction after mandibular third molar surgery in patients less than 30 years of age. A prospective study following removal of 1220 mandibular third molars. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2013;116(4):411-7.
11. Hillerup S, Stoltze K. Lingual nerve injury in third molar surgery I. Observations on recovery of sensation with spontaneous healing. *Int J Oral Maxillofac Surg.* 2007;36(10):884-9.
12. Shafer DM, Frank ME, Gent JF, Fischer ME. Gustatory function after third molar extraction. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1999;87(4):419-28.
13. Güllicher D, Gerlach KL. Sensory impairment of the lingual and inferior alveolar nerves following removal of impacted mandibular third molars. *Int J Oral Maxillofac Surg.* 2001;30(4):306-12.
14. Akal UK, Kuçukyavuz Z, Nağcı R, Yılmaz T. Evaluation of gustatory function after third molar removal. *Int J Oral Maxillofac Surg.* 2004;33(6):564-8.
15. Ridaura-Ruiz L, Figueiredo R, Valmaseda-Castellon E, Berini-Aytes L, Gay-Escoda C. Sensibility and taste alterations after impacted lower third molar extractions. A prospective cohort study. *Med Oral Patol Oral Cir Bucal.* 2012;17(5):759-64.
16. Anand R, Shankar DP, Manodh P, Devadoss P, Aparna M, Neelakandan RS. Short-Term Evaluation of Gustatory Changes After Surgical Removal of Mandibular Third Molar-A Prospective Randomized Control Trial. *J Oral Maxillofac Surg.* 2018;76(2):258-66.
17. Zur KB, Mu L, Sanders I. Distribution pattern of the human lingual nerve. *Clin Anat.* 2004;17(2):88-92.
18. Pippi R, Spota A, Santoro M. Prevention of Lingual Nerve Injury in Third Molar Surgery: Literature Review. *J Oral Maxillofac Surg.* 2017;75(5):890-900.
19. Dias GJ, de Silva RK, Shah T, Sim E, Song N, Colombage S, et al. Multivariate assessment of site of lingual nerve. *Br J Oral Maxillofac Surg.* 2015;53(4):347-51.
20. Chossegros C, Guyot L, Cheynet F, Belloni D, Blanc JL. Is lingual nerve protection necessary for lower third molar germeotomy? A prospective study of 300 procedures. *Int J Oral Maxillofac Surg.* 2002;31(6):620-4.
21. Miloro M, Halkias LE, Slone HW, Chakeres DW. Assessment of the lingual nerve in the third molar region using magnetic resonance imaging. *J Oral Maxillofac Surg.* 1997;55(2):134-7.
22. Olsen J, Papadaki M, Troulis M, Kaban LB, O'Neill MJ, Donoff B. Using ultrasound to visualize the lingual nerve. *J Oral Maxillofac Surg.* 2007;65(11):2295-300.
23. Chan HL, Leong DJ, Fu JH, Yeh CY, Tatarakis N, Wang HL. The significance of the lingual nerve during periodontal/implant surgery. *J Periodontol.* 2010;81(3):372-7.
24. Gargallo-Albiol J, Buenechea-Imaz R, Gay-Escoda C. Lingual nerve protection during surgical removal of lower third molars. a prospective randomised study. *Int J Oral Maxillofac Surg.* 2000;29(4):268-71.
25. Leung YY, Cheung LK. Risk factors of neurosensory deficits in lower third molar surgery: an literature review of prospective studies. *Int J Oral Maxillofac Surg.* 2011;40(1):1-10.
26. Harn SD, Durham TM. Incidence of lingual nerve trauma and postinjection complications in conventional mandibular block anesthesia. *J Am Dent Assoc.* 1990;121(4):519-23.
27. Pogrel MA, Thamby S. Permanent nerve involvement resulting from inferior alveolar nerve blocks. *J Am Dent Assoc.* 2000;131(7):901-7.
28. Jones RH. Repair of the trigeminal nerve: a review. *Aust Dent J.* 2010;55(2):112-9.
29. Renton T, Thexton A, McGurk M. New method for the objective evaluation of injury to the lingual nerve after operation on third molars. *Br J Oral Maxillofac Surg.* 2005;43(3):238-45.
30. Renton T, Thexton A, McGurk M. Objective evaluation of iatrogenic lingual nerve injuries using the jaw-opening reflex. *Br J Oral Maxillofac Surg.* 2005;43(3):232-7.
31. Cheung LK, Leung YY, Chow LK, Wong MC, Chan EK, Fok YH. Incidence of neurosensory deficits and recovery after lower third molar surgery: a prospective clinical study of 4338 cases. *Int J Oral Maxillofac Surg.* 2010;39(4):320-6.
32. Holland GR, Robinson PP. Axon populations in cat lingual and chorda tympani nerves. *J Dent Res.* 1992;71(8):1468-72.