



## How to Help Managing Emotions Felt by Bipolar Patients?

**Bourin M\***

*Neurobiology of Anxiety and Mood Disorders, University of Nantes, France*

### Abstract

The regulation of the control of the emotional experience is one of the determining parameters of the adaptation of an individual to his environment. Bipolar disorder is a psychiatric pathology in which disturbances in the processing of emotions are particularly marked during acute episodes and persist during the interictal period. Treatment with mood stabilizers generally improves the perception of emotions, while antidepressants are brakes on this perception. Finally, simple psychological measures can be useful in controlling emotions.

**Keywords:** Bipolar disorder; Emotions; Euthymic state; Mood stabilizers

### Introduction

The management of emotions is the lot of many aspects of mental health, in particular bipolar disorder [1]. Some of the emotions we face include anger, depression, despair, loneliness, indifference, fear, sadness, love, happiness, confidence, peace, strength, and relaxation [2]. We often feel several emotions at the same time. Then there are times when we feel empty and nothing seems to bother us.

Trying to understand how we feel is a lot harder than it looks because we often feel more than one emotion at a time and also have trouble explaining our emotions to others. We may know that we are depressed, or hypomanic, or in a mixed state, but determining the emotional details of these states is extremely tricky [3]. When friends, family members or even doctors ask the patient how he is feeling, he does not always know what to say or to answer. Writing often helps us understand our emotional state. We have also learned to ask ourselves a series of questions, such as: How and why do we feel what we feel. Often there are not always answers to these questions when treating bipolar disorder because the patient cannot figure out what they are feeling.

Current classifications of mental disorders do not take into account the euthymic phase of bipolar disorder. However, it has now been established that many patients have comorbid conditions and/or residual symptoms during these phases [4].

Studies on impulsivity in bipolar patients have demonstrated the interest of this type of approach for improving our understanding of the modifications occurring during episodes and for a better characterization of the euthymic period [5].

Surprisingly few studies have addressed the emotional reactivity of bipolar patients. The mood episodes characterizing bipolar disorder are most often defined on the basis of emotional tone (sadness/euphoria). Most bipolar patients spontaneously report having excessively strong emotional reactions, even between depressive or manic episodes [6]. They often say that they are more sensitive than others to events. Several elements suggest that emotional reactivity is disturbed in bipolar patients even during euthymic periods. Indeed, it is linked to stress management during these phases [7]. Stressful events have been shown to have deleterious effects, increasing the frequency of relapses in patients with bipolar disorder [8]. On the other hand, bipolar patients experience considerable daily variation in affect, even when they are euthymic. Similarly, having “highs and lows” during euthymic periods is predictive of recurrent depressive episodes.

### Emotional Hyper-Reactivity during Euthymic Periods

During euthymic periods, bipolar patients experience more intense emotions than patients with personality disorders or control subjects, resulting in a higher level of emotional variability [9].

These results were obtained in studies using self-administered questionnaires according to the classic technique of showing the subject positive, neutral or negative images, ensuring that the patients have a good understanding of the images shown to them and assess subjective and objective factors related to the emotions triggered by images [10]. Thus, by this method of

### OPEN ACCESS

#### \*Correspondence:

Michel Bourin, Neurobiology of Anxiety and Mood Disorders, University of Nantes, 98, rue Joseph Blanchart, 44100 Nantes, France;

E-mail: michel.bourin@univ-nantes.fr

Received Date: 07 Jun 2023

Accepted Date: 20 Jun 2023

Published Date: 24 Jun 2023

#### Citation:

Bourin M. How to Help Managing Emotions Felt by Bipolar Patients?. *Am J Med Public Health.* 2023; 4(3): 1046.

Copyright © 2023 Bourin M. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

emotional induction, the emotional reactivity of the subjects can be studied objectively, by comparing it with that of control subjects who consider the same stimuli as neutral. Subjective ratings of arousal and startle reflex when viewing neutral images are greater in euthymic bipolar patients than in control subjects, demonstrating their higher level of emotional reactivity.

Bipolar patients present with episodes of hypomania or depression interspersed with sub syndromic features between episodes. These minimal thymic elements may result from underlying emotional hyper-reactivity, leading to emotional reactivity that becomes chronic and corresponds to sub syndromic symptoms [11]. The tone of these symptoms in a sub depressive or euphoric and irritable mode may depend on the environmental situation, but also on the character traits of the subjects or even on comorbidities, including, in particular, anxiety.

This emotional hyper-reactivity seems to increase considerably during manic states, associated with all types of stimuli (positive, neutral and negative). In euthymic periods, this emotional hyper-reactivity seems to be limited to neutral situations [12]. This can be explained by the fact that the positive and negative images induce a stronger reactivity which does not make it possible to discriminate the differences by reaching a point of saturation.

### Is Emotional Reactivity Hereditary?

It would be useful to conduct similar studies to measure emotional reactivity in parents of bipolar subjects. The presence of the same characteristics in patients and their relatives would identify this emotional hyper-reactivity as an endophenotype associated with susceptibility to bipolar disorder [14]. Emotional hyper-reactivity may also be a risk factor for developing acute episodes in bipolar patients.

Emotional hyper-reactivity may explain the particularly acute sensitivity of bipolar patients to life events. Bipolar subjects not only have a high risk of recurrence when faced with major life events, but they also appear to be particularly susceptible to minor life events [14]. It was found that events that seemed minor and could be considered almost neutral were nevertheless likely to disrupt social routine and trigger bouts of mania.

Is emotional reactivity an endophenotype of bipolar disorder? If so, it could be used to identify subjects at risk of developing bipolar disorder.

### What Effect do Treatments have on Emotional Reactivity?

A good response to mood stabilizer treatment is most often associated with a decrease in emotional reactivity [15]. In contrast, bipolar patients treated for long periods (several years) with antidepressants have been shown to tend to progress to a chronic state of emotional reactivity, mainly to dysphoric episodes (unstable mood) associated with insomnia [16]. This reflects their very often unexplained discomfort, and paradoxically makes it difficult to stop antidepressants, which must be done by gradually reducing the doses, adding or increasing a mood stabilizer [17].

Studies of emotional reactivity can thus contribute to improving our understanding of the mechanisms underlying mood changes. Neurobiological disturbances involved in bipolar disorders; suggest that these are characterized by an increased sensitivity to emotionally

salient environmental information and an inability to regulate their mood [18]. Additionally, bipolar patients appear to have an inability to differentiate between relevant and irrelevant emotional stimuli.

Neuroimaging studies in children and adults show the involvement of the prefrontal cerebral regions, and more particularly the dorsolateral prefrontal cortex and the anterior cingulate cortex in the genesis of emotions [19]. Thus, the definition of bipolar disorder could be refined in light of enduring dysfunction in emotional reactivity, rather than simply as recurrence of episodes. A mild emotional hyper-reactivity characterizes the inter-crisis periods and partly explains the high prevalence of sub-syndromic symptoms. It has also been shown that bipolar mood episodes can also be defined by emotional reactivity, not just affect tone (euphoria/depressed mood). Manic and mixed states are characterized by high emotional hyper-reactivity and this concept, extended to depressive episodes, makes it possible to distinguish two types of bipolar depression. Indeed, mixed bipolar patients can experience, at the same time or in a very short interval, emotions as different as euphoria, sadness, irritability or anxiety. Thus, bipolar disorder must be considered as a global emotional dysregulation, fluctuating from inhibition to excitation [20,21].

### Can we Improve this “Disease of Emotions”?

When it comes to emotions, certain actions can be taken to help manage emotions:

1. Exercise increases the synthesis of brain neurotransmitters that serve to regulate emotions.
2. Spend time with people and talk to them about how you feel.
3. Various forms of talk therapy are beneficial for emotional situations (talk groups).
4. Don't get carried away with negative thoughts, even if it's really easy. The patient will improve over time in positive thinking.
5. Meditation is known to help keep people calm and feel at peace.
6. Depending on the music, it can help relax people or bring them joy.
7. Being grateful helps you achieve all the good things you have in life.
8. Whether indoors or outdoors, playing and having fun with others can help you feel better.
9. Sometimes a simple distraction like watching TV or reading a book can help lift your mood.
10. Writing or keeping a diary of your emotions helps keep track of feelings over a period of time.

It is very important for bipolar people to keep track of their emotions and their evolution over time [22]. Other close people can also help determine the emotional state: In fact, to understand how we feel, they can observe what we say and how we act. Body language provides information about emotional state; it says more about a person's emotions than verbal communication. Paying attention to your own body language is important, especially when you don't know how you feel, it can be the key to determining your emotional state.

Emotion monitoring and tracking is useful, especially for doctors.

During consultations, psychiatrists and therapists often begin by saying:

" So how are you?"

The patient usually does not know how to answer this question, but if one keeps track of his moods, this attention contributes to the treatment and general progress. If one can manage/monitor one's emotions then one can properly inform the doctor of the progress being felt and the treatment can be more accurate and helpful.

## References

1. Bourin M. Psychoeducation of bipolar disorder patients and their relatives. *Arch Depress Anxiety*. 2018;4(1):001-2.
2. Fredrickson BL. The role of positive emotions in positive psychology. The broaden-and-build theory of positive emotions. *Am Psychol*. 2001;56:218-26.
3. Emilien G, Septien L, Brisard C, Corruble E, Bourin M. Bipolar disorder: how far are we from a rigorous definition and effective management? *Prog Neuropsychopharmacol Biol Psychiatry*. 2007;31:975-96.
4. McIntyre RS, Alda M, Baldessarini RJ, Bauer M, Berk M, Correll CU, et al. The clinical characterization of the adult patient with bipolar disorder aimed at personalization of management. *World Psychiatry*. 2022;21:364-87.
5. Santana RP, Kerr-Gaffney J, Ancane A, Young AH. Impulsivity in bipolar disorder: State or trait? *Brain Sci*. 2022;12(10):1351.
6. M'Bailara K, Demotes-Mainard J, Swendsen J, Mathieu F, Leboyer M, Henry C. Emotional hyper-reactivity in normothymic bipolar patients. *Bipolar Disord*. 2009;11:63-9.
7. Henry C, Phillips M, Leibenluft E, M'Bailara K, Houenou J, Leboyer M. Emotional dysfunction as a marker of bipolar disorders. *Front Biosci (Elite Ed)*. 2012;4:2622-30.
8. Kim EY, Miklowitz DJ, Biuckians A, Mullen K. Life stress and the course of early-onset bipolar disorder. *J Affect Disord*. 2007;99:37-44.
9. Wenzel M, Althen H, Veeh J, Reif A. Euthymic patients with predominantly manic polarity avoid happy faces in a dot probe task. *Int J Bipolar Disord*. 2022;10(1):16.
10. Askim K, Knardahl S. The influence of affective state on subjective-report measurements: Evidence from experimental manipulations of mood. *Front Psychol*. 2021;12:601083.
11. Grunze H, Born C. The impact of subsyndromal bipolar symptoms on patient's functionality and quality of life. *Front Psychiatry*. 2020;11:510.
12. M'Bailara K, Atzeni T, Colom F, Swendsen J, Gard S, Desage A, et al. Emotional hyperreactivity as a core dimension of manic and mixed states. *Psychiatry Res*. 2012;197(3):227-30.
13. Guglielmo R, Miskowiak KW, Hasler G. Evaluating endophenotypes for bipolar disorder. *Int J Bipolar Disord*. 2021;9(1):17.
14. Aldinger F, Schulze TG. Environmental factors, life events, and trauma in the course of bipolar disorder. *Psychiatry Clin Neurosci*. 2017;71(1):6-17.
15. Ahn SW, Baek JH, Yang SY, Kim Y, Cho Y, Choi Y, et al. Long-term response to mood stabilizer treatment and its clinical correlates in patients with bipolar disorders: A retrospective observational study. *Int J Bipolar Disord*. 2017;5(1):24.
16. McInerney SJ, Kennedy SH. Review of evidence for use of antidepressants in bipolar depression. *Prim Care Companion CNS Disord*. 2014;16(5):10.4088/PCC.14r01653.
17. Bourin M. Are antidepressants useful in bipolar disease? *Arch Depress Anxiety*. 2017;3(2):058-9.
18. Koenders MA, Dodd AL, Karl A, Greene MJ, Elzinga BM, Wright K. Understanding bipolar disorder within a biopsychosocial emotion dysregulation framework. *J Affect Disord Rep*. 2020;2:100031.
19. Marsh R, Gerber AJ, Peterson BS. Neuroimaging studies of normal brain development and their relevance for understanding childhood neuropsychiatric disorders. *J Am Acad Child Adolesc Psychiatry*. 2008;47(11):1233-51.
20. Stein DJ, Horn N, Ramesar R, Savitz J. Bipolar disorder: Emotional dysregulation and neuronal vulnerability. *CNS Spectr*. 2009;14(3):122-6.
21. Bourin M. Treating the cognitive impairment in bipolar patients. *J Neurosci Neurosurg*. 2018;1(1):107.
22. Fulford D, Peckham AD, Johnson K, Johnson SL. Emotion perception and quality of life in bipolar I disorder. *J Affect Disord*. 2014;152-154:491-7.