# 9

# Adenoviral Epidemic Keratoconjunctivitis and the Challenges for a Timely Treatment

#### Gabriel Gonzalez<sup>1</sup> and Koki Aoki<sup>2\*</sup>

<sup>1</sup>Department of Bioinformatics, Hokkaido University, Japan <sup>2</sup>Department of Ophthalmology, Hokkaido University Graduate School of Medicine, Japan

## **Letter to Editor**

Adenoviruses are cause of ocular morbidity worldwide in the form of Epidemic Keratoconjunctivitis (EKC). The EKC consequences in patients, such as reduced vision and corneal erosion, can last between weeks and months. Besides, many adenoviruses are characterized by high infectivity and structural stability that enable their easy spread by direct contact with infected patients or with contaminated surfaces, creating high risks of nosocomial outbreaks [1]. Additionally, the EKC infections by emergent novel recombinant adenoviral types point to a growing number of EKC threats [2-5]; therefore, efficient and proper diagnosis of adenoviral infections and the subsequent accurate type identification are challenges for clinicians. The spotting of the early symptoms of EKC, its timely treatment, and its swift containment are challenges to be overcome. Although there is no available commercial remedy against EKC, promising research is being performed to develop clinical treatments [6]. Likewise, bioinformatics analyses comparing genomic variation across adenoviral types aim to better understanding the cellular pathways affected by the adenoviral infections [7]. The characterization of host-pathogen interactions will lead us to the improvement of treatment and prevention practices. In short, despite our growing knowledge about adenoviral EKC, much remains to be clarified to convert this understanding into practical applications to alleviate the health and economic burden of EKC in our society.

#### References

1. Rutala WA, Peacock JE, Gergen MF, Sobsey MD, Weber DJ. Efficacy of hospital germicides against adenovirus 8, a common cause of epidemic keratoconjunctivitis in health care facilities. Antimicrob Agents Chemother. 2006;50(4):1419-24.

### **OPEN ACCESS**

#### \*Correspondence:

Koki Aoki, Department of Ophthalmology, Faculty of Medicine and Graduate School of Medicine, Hokkaido University, Sapporo Japan, E-mail: mdaoki@itg.bcj.ne.jp Received Date: 20 Sep 2017 Accepted Date: 12 Oct 2017 Published Date: 19 Oct 2017

#### Citation:

Gonzalez G, Aoki K. Adenoviral Epidemic Keratoconjunctivitis and the Challenges for a Timely Treatment. J Clin Ophthalmol Eye Disord. 2017; 1(3): 1015.

**Copyright** © 2017 Koki Aoki. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

- Aoki K, Ishiko H, Konno T, Shimada Y, Hayashi A, Kaneko H, et al. Epidemic keratoconjunctivitis due to the novel hexon-chimeric-intermediate 22,37/H8 human adenovirus. J Clin Microbiol. 2008;46(10):3259-69.
- 3. Kaneko H, Aoki K, Ishida S, Ohno S, Kitaichi N, Ishiko H, et al. Recombination analysis of intermediate human adenovirus type 53 in Japan by complete genome sequence. J Gen Virol. 2011;92(6):1251-9.
- Kaneko H, Aoki K, Ohno S, Ishiko H, Fujimoto T, Kikuchi M, et al. Complete genome analysis of a novel intertypic recombinant human adenovirus causing epidemic keratoconjunctivitis in Japan. J Clin Microbiol. 2011;49(2):484-90.
- Kaneko H, Iida T, Ishiko H, Ohguchi T, Ariga T, Tagawa Y, et al. Analysis of the complete genome sequence of epidemic keratoconjunctivitis-related human adenovirus type 8, 19, 37 and a novel serotype. J Gen Virol. 2009;90(6):1471-6.
- Caraballo R, Saleeb M, Bauer J, Liaci AM, Chandra N, Storm RJ, et al. Triazole linker-based trivalent sialic acid inhibitors of adenovirus type 37 infection of human corneal epithelial cells. Org Biomol Chem. 2015;13(35):9194-205.
- Gonzalez G, Koyanagi KO, Aoki K, Watanabe H. Interregional coevolution analysis revealing functional and structural interrelatedness between different genomic regions in Human mastadenovirus D. J Virol. 2015;89(12):6209-17.