



Exercise Training and Hypertension: Ready for Prime-Time?

Paolo Severi^{1,2}, Stefania D Emidio^{1†}, Massimo Armeni³, Veronica Bravi⁴ and Massimo Leggio^{2*}

¹Department of Physical Medicine, Salus Infirmorum Clinic, Italy

²Department of Medicine and Rehabilitation, San Filippo Neri Hospital – Salus Infirmorum Clinic, Rome, Italy

³Department of Physical Medicine, Institute for Craniosacral Therapies, Italy

⁴Department of Physical Medicine, Still Osteopathic Institute, Italy

Editorial

A sedentary lifestyle has been characterized as an independent risk factor for cardiovascular disease [1]. Many studies have shown the inverse association between physical activity level and the incidence of cardiovascular diseases [2-4]. Low aerobic fitness is a strong predictor for future cardiovascular disease and all-cause mortality in both healthy and cardiovascular disease patients, including those with hypertension [5-7]. Various mechanisms could be involved in the cardiovascular protective effects of physical activity, including improvement in endothelial function, a decrease in sympathetic neural activity and a reduction in arterial stiffness [8,9].

Hypertension is one of the most important risk factors for cardiovascular disease, and has been ranked as the leading cause for death and disability worldwide: therefore, adequate control of blood pressure is important for public health [10]. Lowering of blood pressure and prevention of hypertension is in first instance preferable by lifestyle changes. These include weight loss, moderation of alcohol intake, a diet with increased fresh fruit and vegetables, reduced saturated fat, reduced salt intake, reduced stress, and, finally, increased physical activity [11,12]. With regard to the latter, former guidelines predominantly recommended aerobic exercises such as walking, jogging, and cycling for lowering blood pressure.

Even if drug therapy remains the mainstay of hypertension management, on the other hand blood pressure control goes beyond adherence with drug therapy as there are other associated factors [13], and achieving adequate blood pressure control with antihypertensive medication remains an elusive goal for many patients anyway [14]. Furthermore, medical management of hypertension is often complicated by concomitant comorbid conditions such as dyslipidaemia, hyperinsulinaemia, glucose intolerance, reduced arterial compliance, sympathetic over-activity and obesity; for instance, some antihypertensive agents adversely affect other cardiovascular risk factors, and adherence to medication is often a problem. However, lifestyle changes improve multiple risk factors without any side effects [15,16], and physical exercise has been demonstrated as a positive and effective adjunct to other lifestyle measures in the prevention and management of hypertension [17]. As a consequence, physical activity is recommended as prevention, treatment, and control of all stages of hypertension [18-21], and therefore both the American Heart Association and the American College of Sports Medicine have endorsed the inclusion of resistance training as an integral part of an exercise program for promoting health and preventing cardiovascular disease [22,23], in fact, continuous exercise training is the type of physical activity most frequently recommended to hypertensive subjects [19], and in some patients regular aerobic exercise also reduces or eliminates the need for antihypertensive medication [24].

In a previous systematic review with meta-analysis provided by Cornelissen et al. [25] the authors reviewed the effect of resistance training on blood pressure and other cardiovascular risk factors in adults, and the findings suggested that both moderate-intensity dynamic resistance training and low-intensity isometric resistance training may cause a reduction in systolic and diastolic blood pressure; furthermore, dynamic resistance training favorably affected some other cardiovascular risk factors such as an increase in peak VO₂ and a reduction in body fat and plasma triglycerides. The clinical importance of these blood pressure reductions can be estimated from large, prospective intervention studies investigating morbidity and mortality outcomes that suggest that small reductions in resting systolic and diastolic blood pressure of 3 mmHg can reduce coronary heart disease risk by 5%, stroke

OPEN ACCESS

*Correspondence:

Massimo Leggio, Department of Medicine and Rehabilitation, Cardiac Rehabilitation Operative Unit, San Filippo Neri Hospital – Salus Infirmorum Clinic, Via della Lucchina 41, 00135 Rome, Italy, Tel: +3906302511; E-mail: mleggio@libero.it

Received Date: 22 Oct 2016

Accepted Date: 16 Dec 2016

Published Date: 02 Jan 2017

Citation:

Severi P, Emidio SD, Armeni M, Bravi V, Leggio M. Exercise Training and Hypertension: Ready for Prime-Time?. *Sports Med Rehabil J.* 2017; 2(1): 1011.

Copyright © 2017 Leggio M. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

by 8%, and all-cause mortality by 4% [26-28].

Moreover, given that the association between blood pressure and cardiovascular risk has no lower threshold, reductions of this magnitude in individuals with even optimal blood pressure at baseline still seem to have clinical significance [11], and this strongly underlines the potential of resistance training as adjuvant therapy for the prevention and treatment of high blood pressure. In addition, the positive effect of resistance training on other cardiovascular risk factors suggest that a better physical fitness is associated with a lower risk of all-cause mortality and cardiovascular events and is independently associated with longevity [18,29]: aerobic exercise not only reduces blood pressure, it also lowers levels of low-density lipoprotein cholesterol, reduces insulin resistance and glucose intolerance, and often is associated with reduced body weight [30]. Finally, it must be reminded that although genetic predisposition is a risk factor for hypertension, studies have shown that behavioural factors, such as sedentary lifestyle, overshadow genetic predisposition as a cause of hypertension [31].

To prescribe resistance training as a potential tool in the control of blood pressure, one should know how different training characteristics influence the blood pressure response. Aerobic or endurance exercises are dynamic physical exercises, involving large muscle groups, which increase cardio-respiratory fitness and help in weight control. They are those in which the exercising muscles make use of oxygen. Aerobic exercises are effective in lowering blood pressure or preventing hypertension [19]. These endurance activities, such as walking, swimming, cycling and low-impact aerobics (dancing or rope skipping), are the core of the exercise programme for managing hypertension. The blood pressure responses to aerobic exercises depend on the activities engaged in. Several studies have shown that high-intensity aerobic interval training improves aerobic fitness and reduces several cardiovascular risk factors more than moderate intensity continuous training [32-36]. Although a large body of evidences suggests that vigorous training compared to moderate- and low-intensity training elicits more cardiovascular benefits, also in hypertensive patients [37,38], several meta-analyses indicate little or no intensity-dependent effect of exercise training for the reduction of the blood pressure [18,19,39]. In studies where low and high intensity training programmes have been directly compared, the lower intensity programmes were either more effective [40] or as effective [41] as the higher intensity programmes, but more recent reviews [28,42] reported no influence of exercise intensity on blood pressure reduction following exercise treatment. Further, aerobic exercise training of 60–85% of age-predicted (220-age in years) maximal heart rate may be as, or more, effective as high-intensity exercise in lowering BP in hypertensive patients [43]. As a consequence, no specific guidelines delineate exercise intensity and frequency, but a recent study performed by Molmen-Hansen et al. [44] study indicates that the exercise-induced lowering effect of blood pressure is intensity dependent: in addition, high-intensity aerobic interval training was superior to moderate intensity continuous training in terms of improved cardiac and endothelial function, aerobic capacity, and heart rate recovery, and so aerobic interval training should be viewed as an effective method to lower blood pressure and improve other cardiovascular risk factors. Moreover, a growing body of evidences suggests and our results also confirm the positive impact of exercise training on both left and right ventricular systo-diastolic function, in terms of subclinical improvement, in pharmacologically treated hypertensive patients [9], and it must be reminded that given

the poor blood pressure control in the general population and the enormous prevalence of hypertension the possible means and process of aerobic exercise complementing antihypertensive drug therapy in order to achieve higher blood pressure control rates would place a large number of people at decreased risk for cardiovascular morbidity and mortality and so would have enormous implications and would be of considerable and growing global public health importance.

In conclusion, there is an ample evidence in the literature that aerobic exercise lowers blood pressure in individuals with hypertension. This is expected to encourage physicians to recommend or refer people with hypertension, especially those who require more than two antihypertensive drugs to achieve blood pressure control, for aerobic exercise. Increased attention needs to be placed on strategies to maintain or improve fitness, efforts to encourage physical activity should urgently be intensified and supported, and sedentary lifestyle should be viewed as one of several major modifiable risk factors in the prevention and management of cardiovascular disease. Considering the dearth of studies on the possible additive or complementary effect of aerobic exercise on antihypertensive drug therapy in achieving blood pressure control, more empirical studies are needed to make an assertion about the role of aerobic exercise, in conjunction with antihypertensive drug therapy, in achieving blood pressure control.

References

1. National Heart Institute. Physical activity and cardiovascular health. *JAMA*. 1996; 273: 241-246.
2. Tanaka H, Dinanno FA, Monahan KD, Clevenger CM, DeSouza CA, Seals DR. Aging, habitual exercise, and dynamic arterial compliance. *Circulation*. 2000; 102: 1270-1275.
3. Kakiyama T, Sugawa J, Murakami H, Maeda S, Kuno S, Matsuda M. Effects of short-term endurance training on aortic distensibility in young males. *Med Sci Sports Exerc*. 2005; 37: 267-271.
4. Cameron JD, Dart MD. Exercise training increase total systemic arterial compliance in humans. *Am J Physiol*. 1994; 266: H693-H701.
5. Kokkinos P, Manolis A, Pittaras A, Doumas M, Giannelou A, Panagiotakos DB, et al. Exercise capacity and mortality in hypertensive men with and without additional risk factors. *Hypertension*. 2009; 53: 494-499.
6. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood E. Exercise capacity and mortality among men referred for exercise testing. *N Eng J Med*. 2002; 346: 793-801.
7. Boman K, Gerds E, Wachtell K, Dahlof B, Nieminen MS, Olofsson M, et al. Exercise and cardiovascular outcomes in hypertensive patients in relation to structure and function of left ventricular hypertrophy: the LIFE study. *Eur J Cardiovasc Prev Rehabil*. 2009; 16: 242-248.
8. Guimaraes GV, Ciolac EG, Carvalho VO, D'Avila VM, Bortolotto LA, Bocchi EA. Effects of continuous vs. interval exercise training on blood pressure and arterial stiffness in treated hypertension. *Hypertens Res*. 2010; 33: 627-632.
9. Leggio M, Mazza A, Cruciani G, Sgorbini L, Pugliese M, Bendini MG, et al. Effects of exercise training on systo-diastolic ventricular dysfunction in patients with hypertension: an echocardiographic study with tissue velocity and strain imaging evaluation. *Hypertens Res*. 2014; 37: 649-654.
10. Caldaroni E, Lombardi M, Severi P, Leggio M. Ambient Air Pollution and Hypertension: A Relationship that Strikes Around the Clock. *Arch Clin Hypertens*. 2016; 1: 044-045.
11. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. National High Blood Pressure Education Program Coordination Committee. Seventh report of the Joint National Committee on

- Prevention, Detection, Evaluation and Treatment of High Blood pressure. Hypertension. 2003; 42: 1206-1252.
12. Mancia G, De Backer G, Dominiczak A, Cifkova R, Fagard R, Germano G, et al. Management of Arterial Hypertension of the European Society of Hypertension; European Society of Cardiology: Guidelines for the Management of Arterial Hypertension: the Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens.* 2007; 25: 1105-1187.
 13. Mant J, McManus RJ. Does it matter whether patients take their antihypertensive medication as prescribed? The complex relationship between adherence and blood pressure control. *J Hum Hypertens.* 2006; 20: 551-553.
 14. Halm J, Amoako E. Physical activity recommendation for hypertension management: does healthcare providers' advice make a difference? *Ethn Dis.* 2008; 18: 278-282.
 15. Vatten LJ, Nilsen TI, Holmen J. Combined effect of blood pressure and physical activity on cardiovascular mortality. *J Hypertens.* 2006; 24: 1939-1946.
 16. Burke V, Beilin LJ, Cutt HE, Mansour J, Williams A, Mori TA. A lifestyle programme for treated-hypertensives-improved-health-related behaviours and cardiovascular risk factors, a randomized control trial. *J Clin Epidemiol.* 2007; 60: 133-141.
 17. Cox KK. Exercise and blood pressure: applying findings from the laboratory to the community settings. *Clin Exp Pharmacol Physiol.* 2006; 33: 868-871.
 18. Cornelissen VA, Fagard RH. Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors. *Hypertension.* 2005; 46: 667-675.
 19. Pescatello LS, Franklin BA, Fagard R, Farquhar WB, Kelley GA, Ray CA. American College of Sports Medicine position stand. Exercise and hypertension. *Med Sci Sports Exerc.* 2004; 36: 533-553.
 20. Graham I, Atar D, Borch-Johnsen K, Boysen G, Burell G, Cifkova R, et al. European guidelines on cardiovascular disease prevention in clinical practice: full text. Fourth Joint Task Force of the European Society of Cardiology and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of nine societies and by invited experts). *Eur J Cardiovasc Prev Rehabil.* 2007; 14: S1-S113.
 21. Kokkinos PF, Giannelou A, Manolis, Pittaras A. Physical activity in the prevention and management of high blood pressure. *Hellenic J Cardiol* 2009; 50: 52-59.
 22. Williams MA, Haskell WL, Ades PA, Amsterdam EA, Bittner V, Franklin BA, et al. American Heart Association Council on Clinical Cardiology; American Heart Association Council on Nutrition, Physical Activity, and Metabolism. Resistance exercise in individuals with and without cardiovascular disease: 2007 update: a scientific statement from the American Heart Association Council on Clinical Cardiology and Council on Nutrition, Physical Activity and Metabolism. *Circulation.* 2007; 116: 572-584.
 23. Braith RW, Stewart KJ. Resistance exercise training: its role in the prevention of cardiovascular disease. *Circulation.* 2006; 113: 2642-2650.
 24. Appel LJ. Lifestyle modification as a means to prevent and treat high blood pressure. *J Am Soc Nephrol.* 2003; 14: S99-S102.
 25. Cornelissen VA, Fagard RH, Coeckelberghs E, Vanhees L. Impact of resistance training on blood pressure and other cardiovascular risk factors. A meta-analysis of randomized, controlled trials. *Hypertension.* 2011; 58: 950-958.
 26. Lewington S, Clarke R, Qizilbash N, Peto R, Collins R. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies, Prospective Studies Collaboration. *Lancet.* 2002; 360: 1903-1913.
 27. Stamler R. Implications of the INTERSALT study. *Hypertension.* 1991; 17: 116-120.
 28. Whelton PK, He J, Appel LJ, Cutler JA, Havas S, Kotchen TA, et al. Primary prevention of hypertension: clinical and public health advisory from the National High Blood Pressure Education Program. *JAMA.* 2002; 288: 1882-1888.
 29. Kodama S, Saito K, Tanaka S, Maki M, Yachi Y, Asumi M, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women. *JAMA.* 2009; 301: 2024-2035.
 30. Dengel DR, Hagberg JM, Pratley RE, Rogus EM, Golberg AP. Improvements in blood pressure, glucose metabolism, and lipoprotein lipids after aerobic exercise plus weight loss in obese, hypertensive middleaged men. *Metabolism.* 1998; 47: 1075-1082.
 31. Douglas JG, Bakris GL, Epstein M, Ferdinand KC, Ferrario C, Flack JM, et al. Management of high blood pressure in Americans. Consensus statement of the Hypertension in Africans and Americans Working Group of the International Society on Hypertension in Blacks. *Arch Intern Med.* 2003; 163: 525-541.
 32. Swain DP, Franklin BA. Comparison of cardioprotective benefits of vigorous versus moderate intensity aerobic exercise. *Am J Cardiol.* 2006; 97: 141-147.
 33. Wisloff U, Stoylen A, Loennechen JP, Bruvold M, Rognmo O, Haram PM, et al. Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study. *Circulation.* 2007; 115: 3086-3094.
 34. Rognmo O, Hetland E, Helgerud J, Hoff J, Slordahl SA. High intensity aerobic interval exercise is superior to moderate intensity exercise for increasing aerobic capacity in patients with coronary artery disease. *Eur J Cardiovasc Prev Rehabil.* 2004; 11: 216-222.
 35. Wenger HA, Bell GJ. The interactions of intensity, frequency and duration of exercise training in altering cardiorespiratory fitness. *Sports Med.* 1986; 3: 346-356.
 36. Tjonna AE, Lee SJ, Rognmo O, Stolen TO, Bye A, Haram PM, et al. Aerobic interval training versus continuous moderate exercise as a treatment for the metabolic syndrome: a pilot study. *Circulation.* 2008; 118: 346-354.
 37. Cornelissen VA, Verheyden B, Aubert AE, Fagard RH. Effects of aerobic training intensity on resting, exercise and post-exercise blood pressure, heart rate and heart-rate variability. *J Hum Hypertens.* 2010; 24: 175-182.
 38. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation.* 2007; 116: 1081-1093.
 39. Cornelissen VA, Arnout J, Holvoet P, Fagard RH. Influence of exercise at lower and higher intensity on blood pressure and cardiovascular risk factors at older age. *J Hypertens.* 2009; 27: 753-762.
 40. Roger MW, Probst MM, Gruber JJ, Berger R, Boone JB. Differential effects of exercise training intensity on blood pressure and cardiovascular responses to stress in borderline hypertensive humans. *J Hypertens.* 1996; 14: 1369-1375.
 41. Tashiro E, Miura S, Koga M, Sasaguri M, Ideishi M, Ikeda M, et al. Crossover comparison between the depressor effects of low and high work-rate exercise in mild hypertension. *Clin Exp Pharmacol Physiol.* 1993; 20: 689-696.
 42. Fagard RH. Exercise characteristics and the blood pressure responses to dynamic physical training. *Med Sci Sports Exerc.* 2001; 33: S484-S492.

43. Kokkinos P, Pittaras A, Manolis A, Panagiotakos D, Narayan P, Manjoros D, et al. Exercise capacity and 24-h blood pressure in prehypertensive men and women. *Am J Hypertens.* 2006; 19: 251-258.
44. Molmen-Hansen HE, Stolen T, Tjonna AE, Aamot IL, Ekeberg IS, Tyldum GA, et al. Aerobic interval training reduces blood pressure and improves myocardial function in hypertensive patients. *Eur J Prev Cardiol.* 2012; 19: 151-160.