

PICC: Just Secure It

Agrati S1 and Ambrosoli AL2*

¹University of Insubria, Varese, Italy

²Department of Anesthesia, Perioperative Medicine and Intensive Care Unit, F. Del Ponte Woman and Children's Hospital, Italy

Clinical Image

Securing a pediatric Peripherally Inserted Central Catheter (PICC) properly is crucial to prevent displacement and ensure appropriate hygiene during dressing changes in order to reduce the risk of infection [1]. An accurate placement significantly lowers complications, enhancing overall patient well-being and safety, particularly essential in pediatric care. The introduction of innovative solutions, such as SecurAcath, allows the achievement of these goals. Its position maneuvers and medications are easy and quick, there are different sizes from 3 Fr to 8 Fr depending on the diameter of the chosen PICC. The right placement of a SecurAcath is reported in the picture above (Figure 1). Once positioned it is possible to disinfect all the skin around the insertion site and eventually place an anti-decubitus device under it. This contributes to improving infection control, simplifying care procedures, and last but not least enhancing outcomes for pediatric patients.



Figure 1: PICC fixed with SecurAcath and medicated with a transparent patch and anti-decubitus device.

OPEN ACCESS

*Correspondence:

Andrea Luigi Ambrosoli, Department of Anesthesia, Perioperative Medicine and Intensive Care Unit, F. Del Ponte Woman and Children's Hospital, ASST Sette Laghi, Varese, via del Ponte n.19,

Italy
Received Date: 07 Feb 2024
Accepted Date: 20 Feb 2024
Published Date: 24 Feb 2024

Citation:

Agrati S, Ambrosoli AL. PICC: Just Secure It. J Surg Tech Proced. 2024; 8(1): 1066.

Copyright © 2024 Ambrosoli AL. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

1. Pet GC, Eickhoff JC, McNevin KE, Do J, McAdams RM. Risk factors for peripherally inserted central catheter complications in neonates. J Perinatol. 2020;40(4):581-8.