



Vasa Vasorum and Coronary Artery Disease

Zenshio Onouchi*

Department of Pediatrics, Kyoto Prefectural University of Medicine, Graduate School of Medical Science, Japan

Commentary

Some arterial diseases reveal endothelial cell dysfunction, in which vasa vasorum play an important role. One of them is the atherosclerosis. Another is Kawasaki disease, we proposed, in which middle-sized arteritis, aortitis, and their aneurysms are due to vasa vasorum vasculitis. In these diseases, many studies have accumulated on the pathogenesis of endothelial cell dysfunction. However, these studies may be adapted to the endothelial cells of every artery, and coronary artery is not specific to them.

Coronary artery has vasa vasorum recognized unique origination and distribution which worse reduced blood flow through vasa vasorum vasculitis due to reduced host Bessel, coronary blood flow, by the decrease in the distensibility of the coronary artery wall.

We proposed a novel treatment of these diseases based on our hypothesis that predilection of coronary artery is attributed to medial ischemic necrosis due to involvement of vasa vasorum.

OPEN ACCESS

*Correspondence:

Zenshio Onouchi, Department of Pediatrics, Kyoto Prefectural University of Medicine, Graduate School of Medical Science, Japan,
E-mail: zen@mrj.biglobe.ne.jp

Received Date: 05 Feb 2018

Accepted Date: 15 Feb 2018

Published Date: 23 Feb 2018

Citation:

Onouchi Z. Vasa Vasorum and Coronary Artery Disease. *Ann Cardiol Cardiovasc Med.* 2018; 2(1): 1012.

Copyright © 2018 Zenshio Onouchi.

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.