



Rectal Carcinoma Metastasis to the Thyroid Gland - The Role of Radiotherapy: Case Report

Danijela Scepanovic*, Mojmir Masar, Pavol Bires, Marek Paluga and Margita Pobjakovska

Department of Radiation Oncology, National Cancer Institute of Slovakia, Bratislava, Slovakia

Abstract

A very rare occurrence of thyroid metastases from rectal cancer is described in the literature. Therefore, there are no therapeutic standards; however, surgery still has a major role in the treatment of these patients. Radiotherapy has a controversial role and its significance is rather palliative. We presented a male patient with metastatic rectal carcinoma to the thyroid gland who has been treated with surgery and palliative radiotherapy in our institution.

Forty-nine months after rectal resection preceded by neoadjuvant radiochemotherapy, metastasis of rectal cancer to the thyroid gland was diagnosed. The patient underwent thyroidectomy followed by palliative radiotherapy. Unfortunately, 5 months after the end of radiotherapy, the residual tumor progressed and compromised swallowing function. Therefore, percutaneous endoscopic gastrostomy was placed. The patient died 10 months after the end of radiotherapy.

The role of radiotherapy in our patient had a short-term effect and a controversial role. Nevertheless, we decided to show this case due to the rare occurrence of metastases to the thyroid gland from rectal cancer. We recommend to think on their potential occurrence enabling early diagnosis and more successful treatment.

Keywords: Thyroid metastases; Rectal carcinoma; Radiotherapy

Introduction

The thyroid gland is rarely the site of metastasis of malignant tumors, and metastatic tumors are reported to represent only 2% to 3% of all malignant tumors of the thyroid gland [1]. Because thyroid metastases have been reported in 1% to 24% of cancer patients in autopsy studies their incidence may be higher [2]. The incidence of thyroid metastasis among rectal cancer patients is extremely rare as confirmed by a study of Lièvre et al. between 1993 and 2004 that identified 6 cases among 5,862 patients with colorectal cancer (0.1%) [2,3]. The most common malignant tumors that metastasize to the thyroid gland are kidney, breast and lung cancer [3,4].

Case Presentation

In June 2017 a 71-year-old male patient underwent total thyroidectomy due to metastasis in the thyroid gland from rectal cancer, followed by radiotherapy.

Previously on January 2013, this patient was treated with neoadjuvant radiochemotherapy for low rectal carcinoma (50.4 Gy/28 fractions concomitantly with capecitabine). The distal edge of the tumor was 4.5 cm from the anus, clinical stage of the disease was cT3cN1M0. In March 2013 he underwent surgery (abdominoperineal resection). The postoperative stage was pT2pN0M0. He continued treatment with adjuvant chemotherapy (capecitabine) for the next 6 months.

In May 2015, 20 months after completion of the adjuvant chemotherapy, a left pulmonary metastasis was diagnosed and wedge resection of the lung was conducted (upper lobe of the lung), followed by chemotherapy with regimen XELOX (capecitabine plus oxaliplatin) up to 8 cycles (from June to December 2015).

Sixteen months after completion of XELOX chemotherapy, our patient experienced the symptoms such as hoarseness and difficulty swallowing.

Computed Tomography (CT) scan of the neck showed a lesion in the left lobe of the thyroid gland (50 mm × 29 mm × 35 mm) and spread of this mass to the tracheoesophageal groove (Figure 1). Otolaryngological examination (ENT) showed the left Recurrent Laryngeal Nerve (RLN) paresis as a consequence of tumor spread into the tracheoesophageal groove. Ultrasonography and Fine

OPEN ACCESS

*Correspondence:

Danijela Scepanovic, Department of Radiation Oncology, National Cancer Institute of Slovakia, Bratislava, Klenova, Bratislava, Slovakia, E-mail: danijelascepanovic@hotmail.com

Received Date: 06 Jun 2020

Accepted Date: 30 Jun 2020

Published Date: 03 Jul 2020

Citation:

Scepanovic D, Masar M, Bires P, Paluga M, Pobjakovska M. Rectal Carcinoma Metastasis to the Thyroid Gland - The Role of Radiotherapy: Case Report. *Am J Head Neck Cancer*. 2020; 1(1): 1001.

Copyright © 2020 Danijela

Scepanovic. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Figure 1: Preoperative CT: Lesion in the left lobe of the thyroid gland.




Figure 2: a) Postoperative CT: Residual tumor, b) CT tumor progression after radiotherapy.

Needle Aspiration (FNA) of the thyroid mass were performed.

In April 2017, the malignant changes suggesting metastasis were cytologically proven. Marker values were: TSH (Thyroid-Stimulating Hormone) - 0.14 mIU/l, fT4 (Free Thyroxine) - 14.7 pmol/l, TG (Thyroglobulin) - 59 ng/ml, CEA (Carcinoembryonic Antigen) - <0.5 ng/ml.

In June 2017 (45 months after completion of adjuvant chemotherapy) the patient underwent thyroid surgery (thyroidectomy without parathyroid gland removal using neuro-electro-stimulator - debulking).

Histopathology and immunohistochemical examination (CDX-2 and Thyroglobulin were positive and Calcitonin, CK19, HBME1, p53 and TTF were negative) confirmed the thyroid metastasis of rectal cancer. Also, metastases were diagnosed in Delphian lymph node, both lobes of thyroid gland, as well as tracheal and esophageal infiltration.

Postoperative CT showed (one month after thyroid surgery): in lower part of the left neck to observe inhomogeneous tumor formation with size 40 mm × 30 mm, with infiltration of the adjacent neck muscles, and spread to the tracheoesophageal groove (Figure 2a).

Thereafter, the patient received palliative radiotherapy at our department. We conducted 3D conformal external beam radiotherapy on the left neck side with a total dose of 40 Gy, hypofractionated regimen with a daily dose of 2.5 Gy (EQD2 [equivalent dose when delivered in 2 Gy/fraction] = 41.67 Gy, BED10 [Biologically Effective Dose] = 50 Gy), by three fields' technique. We used a multi leaf collimator (80 leaves, 40 × 40) and 6-MV X-ray energy on a linear accelerator (Figure 3).

Five months after the end of radiotherapy, the patient had tumor progression (Figure 2b). Subsequently, he received 2 cycles of palliative chemotherapy bFOL (5-fluorouracil bolus/leucovorin + oxaliplatin), however, without effect.




Figure 3: Radiotherapy plan - 3D conformal external beam radiotherapy.

Despite the use of all treatment modalities, the patient clinical condition deteriorated, he could not swallow and eat. Gastroscopy was performed and showed a stenosis just below Killian's dehiscence caused by extramural oppression. In March 2018 the push - PEG (Percutaneous Endoscopic Gastrostomy) was placed. The patient died 11 months after thyroidectomy and 10 months after radiotherapy.

Discussion

In general, thyroid metastases are less common [5]. The incidence of thyroid metastasis among rectal cancer patients is extremely rare as confirmed by a study of Lièvre et al. between 1993 and 2004 that identified 6 cases among 5,862 patients with colorectal cancer (0.1%) [2,3]. This statement is also confirmed by studies from the USA and UK on 15 cases of metastatic thyroid tumors showed that the kidney was the most common primary source of thyroid metastasis, however, only one case showed thyroid metastasis from colorectal cancer [6].

The explanation for this phenomenon can be found in the fact that colorectal cancer spreads to the thyroid gland through the portal vein, vena cava, and pulmonary vein, however, sometimes the special physiology and pathology of the vertebral venous system enable such tumors to bypass the portal vein, pulmonary vein, and vena cava and to be transferred directly to the thyroid or any part of the body, without entering the thoracic and abdominal cavity [7].

In patients with advanced stage of the disease and location of primary tumor in the distal rectum thyroid metastases usually occur and then their occurrence are associated with a poor prognosis and high mortality [8,9]. Thyroid metastases from rectal cancer can occur from 0 to 8 years after rectal resection [5].

The presented patient corresponded with these data. He had a stage III cancer in the distal part of the rectum and thyroid metastasis was diagnosed 49 months after rectal resection.

As already mentioned, thyroid metastases occur in patients with locally advanced rectal cancer and are often diagnosed with other distant metastases, most often with lung metastases (either synchronous or metachronous), this combination is recorded in up to 60% of patients with thyroid metastases [7,9].

Even in our patient thyroid metastasis from the rectal carcinoma occurred after lung metastasis (metachronous with 23 months interval). He underwent pulmonary resection 26 months after rectal resection. In our case report, there was no physiological exception of the spread of rectal cancer and the lung was not skipped, just lung and thyroid metastasis appeared metachronous.

Although there are no treatment standards for thyroid metastases, surgical resection is the treatment of choice, even in extensive metastatic disease. The extent of surgery in terms of total thyroidectomy or less radical resection does not affect the overall survival of patients [10]. The other modalities did not show a significant effect. Colorectal metastases to the thyroid gland do not uptake radioactive iodine, thereupon radioactive iodine therapy is not indicated [6]. Thyroid metastases from rectal cancer is an aggressive disease that is most often locally advanced at the time of diagnosis, and then radiotherapy and chemotherapy are performed, as well as, in patients with comorbidities who are not candidates for surgery, however, most often have a palliative role [6].

Because the thyroid metastasis appeared many years after initial diagnosis, its diagnosis was even more difficult during routine patient follow-up. However, patient did not report any symptoms related to the thyroid function, nor did the diagnostic methods show any changes in the thyroid gland. Actually, he was euthyroid just before thyroidectomy. Clinical features, such as a hoarseness, and dysphagia appeared during the course of disease progression abruptly. Our patient underwent thyroidectomy to avoid the appearance of expected dyspnea when the trachea was also oppressed. Although this aggressive surgery was performed, a residual tumor remained. Then we decided to apply palliative radiotherapy, with mild hypofractionation. The role of radiotherapy in local tumor control was limited and its effect was short term, even subsequent palliative chemotherapy did not control tumor growth.

Five months after the end of radiotherapy, it was essential to place push - PEG due to tumor progression that caused the patient unable to swallow. Our patient died 10 months after the end of radiotherapy.

Conclusion

In routine clinical practice, the occurrence of thyroid metastases from rectal cancer is underestimated because their occurrence is rare. Although thyroid metastases from rectal cancer are not common, we should think about them because when they appears clinically, it is already a very aggressive disease that is difficult to control.

Optimal treatment of thyroid metastases from rectal cancer is still not standard defined, but surgery certainly plays a major role in preventing respiratory and swallowing compression due to rapid growth of metastatic lesions with compression on the trachea and oesophagus. However, the role of radiotherapy in our patient had a short-term effect and a controversial role. Nevertheless, we decided to show this case due to the rare occurrence of thyroid metastases from the rectal cancer and recommended to think on their potential occurrence enabling early diagnosis and more successful treatment.

References

1. Ozawa T, Saito S, Matsuura S, Kishi H, Maeda M, Watanabe T. Thyroid gland metastasis of rectal cancer. *J Surg Case Rep.* 2015;2015(15):1-4.
2. Minami S, Inoue K, Irie J, Mine T, Tada N, Hirabaru M, et al. Metastasis of colon cancer to the thyroid and cervical lymph nodes: A case report. *Surg Case Rep.* 2016;2(1):108.
3. Coelho MI, Albano MN, Costa Almeida CE, Reis LS, Moreira N, Almeida CMC. Colon cancer metastasis to the thyroid gland: A case report. *Int J Surg Case Rep.* 2017;37:221-4.
4. Amenduni T, Carbone A, Bruno R. Precocious and isolated thyroid metastasis of colorectal adenocarcinoma and incidental thyroid papillary microcarcinoma. *Endocrine.* 2014;47(3):969-70.
5. Moria K, Koinumaa K, Nishinob H, Horiea H, Kawarai Lefora A, Sata N. Rectal cancer with a metastasis to the thyroid gland: A case report. *Int J Surg Case Rep.* 2020;66:39-42.
6. Wood K, Vini L, Harmer C. Metastases to the thyroid gland: the Royal Marsden experience. *Eur J Surg Oncol.* 2004;30(6):583-8.
7. Batson OV. The function of the vertebral veins and their role in the spread of metastases. *Ann Surg.* 1940;112(1):138-49.
8. Keranmu A, Zheng H, Wu Y, Zhao J, Xu X, Liu F, et al. Comparative study of single-center patients with thyroid metastases from colorectal cancer and previously reported cases in the literature. *World J Surg Oncol.* 2017;15(1):88.
9. Fujita T, Ogasawara Y, Doihara H, Shimizu N. Rectal adenocarcinoma metastatic to the thyroid gland. *Int J Clin Oncol.* 2004;9(6):515-9.
10. Calzolari F, Sartori PV, Talarico C, Parmeggiani D, Beretta E, Pezzulo L, et al. Surgical treatment of intrathyroid metastases: Preliminary results of a multicentric study. *Anticancer Res.* 2008;28(5B):2885-8.