



Understanding Lower Respiratory Tract Infection: A Comprehensive Review of Associated Risk Factors

Hussain SM*, Rumaisa N and Keshav J

Pharmacology, Nagaon, Assam, India

Abstract

Lower Respiratory Tract Infections (LRTIs), sometimes known as chest infections, impact the lungs and bronchial tubes, among other airways and tissues in the chest. Bronchitis and pneumonia are the two most typical types of chest infections. Most cases of pneumonia are caused by bacteria, whereas viruses cause bronchitis in most cases. Up to one-third of acute stroke patients experience chest infections as a common consequence. Of all the medical problems that accompany a stroke, these infections have the highest attributable mortality due to their considerably increased risk of death. Chest infections continue to be a major worldwide health concern because of the high rates of morbidity and mortality they cause in all age groups. Since treatment and prevention measures are more difficult to come by in low- and middle-income countries, diseases including tuberculosis, bronchitis, and pneumonia are among the most hazardous. This thorough analysis integrates data from recent epidemiological research and therapeutic trials to examine the many risk factors linked to lower respiratory tract infections and chest infections. The review also discusses how access to healthcare and socioeconomic position affect the frequency and consequences of chest infections. This article seeks to clarify these risk factors to improve knowledge of the etiology of chest infections and to direct the creation of focused preventative and therapeutic plans.

Keywords: Chest infection (Pneumonia); Risk Factors; COPD; TB

Introduction

A typical consequence of acute stroke that affects up to one-third of patients is a chest infection. With a three-fold greater risk of death, chest infections have the greatest attributable mortality of all medical conditions, after stroke. A longer hospital stays and a higher chance of being discharged to a nursing home are also linked to chest infections. Pneumonia and bronchitis are the two most prevalent types of chest infections while viruses typically cause bronchitis, bacteria are the primary cause of pneumonia.

Worldwide, respiratory disorders have a significant negative impact on health. Across the globe, five of these illnesses rank among the most frequent causes of serious illness and death. The third greatest cause of death globally, moderate to severe Chronic Obstructive Pulmonary Disease (COPD) is thought to affect 65 million individuals, of whom 3 million dies from it annually. With over 334 million people affected and 14% of children globally suffering from asthma, it is the most common chronic illness among children. 10.4 million cases of Tuberculosis (TB) were reported in 2015; 1.4 million of those cases resulted in death. Worldwide, lung cancer is the most common cancer that kills people, taking the lives of 1.6 million people per year and still rising [1].

Furthermore, several respiratory conditions have a significant but little-understood impact. Over 100 million individuals experience breathing disorders related to sleep. Pulmonary hypertension affects millions of people. Over 50 million individuals suffer from lung ailments related to their jobs. More than 2.4 million fatalities annually are attributed to lower respiratory tract infections, which are particularly common in low- and middle-income nations [2,3].

More people die from lower respiratory tract infections than from malaria, TB, and HIV combined. For both individuals over 65 and children under the age of five who do not belong to the neonatal era leads to death. With 650,000 children under five years old dying from pneumonia in 2019, this age group saw over 15% of all mortality. Furthermore, among those over the age of 70, lower respiratory tract infections resulted in almost a million fatalities. It is also one of the most common causes of hospitalization and the second biggest cause of years lost to premature death. Pneumonia can have long-term effects that include worsening lung health, the development

OPEN ACCESS

*Correspondence:

Sarkar Mubashir Hussain,
Pharmacology, Nagaon, Assam, India,

Received Date: 22 Jul 2024

Accepted Date: 12 Aug 2024

Published Date: 17 Aug 2024

Citation:

Hussain SM, Rumaisa N, Keshav J.
Understanding Lower Respiratory Tract
Infection: A Comprehensive Review of
Associated Risk Factors. *Ann Pulm Res
Med.* 2024; 2(1): 1006.

Copyright © 2024 Hussain SM. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

of chronic respiratory diseases like bronchiectasis, and chronic cardiovascular disorders like heart failure, cardiac arrhythmias, and coronary syndromes, in addition to death and hospitalization [4].

Excessive risk of chest infection during the early phase of stroke has been associated in previous studies with many factors. These include diabetes mellitus and more severe neurologic impairment. A poor swallowing technique could also play a significant role; dysphagia or a failed initial swallow assessment is linked to a higher risk of chest infection following a stroke. Numerous other risk variables, such as dental health and the presence of oral pathogens, have also been linked to chest infections in studies of elderly and non-acute stroke patients. Pneumonia, which has been a major concern for millennia, is classified as a respiratory illness that affects every section of the respiratory tract. When it comes to medical conditions, there are two main pneumonia that was acquired in a hospital, or HAP, and pneumonia that was acquired in a community, or CAP.

The phrase "infection cause" describes the variety of bacteria, fungi, viruses, and parasites that can result in sepsis and other upper respiratory tract infections. Pneumonia has a very poor survival probability when detected much later. Pneumonia is more common in children 0 to 5 years old and in older adults (above 50 or in their 60s). The immunological response of the host and the virus are the main factors that determine the likelihood of infection.

Community-Acquired Pneumonia (CAP) continues to be a significant contributor to mortality and morbidity. It is crucial to implement preventive efforts that identify and address modifiable risk factors to minimize deaths linked to CAP. There are few population-based studies on CAP risk factors. Alcoholism, heart disease, lung disease, immunosuppressive medication, and in a Finnish study with people over 60, additional illnesses were found to be separate causes of pneumonia risk.

Comparable outcomes found on UK study that included significance of cigarette smoking. Additional risk variables found in a Spanish study included low body mass index, prior respiratory infections, and prior pneumonia. Studies using population-based cohorts from the United States of America validated these results and emphasized the impact of diabetes, asthma, and excessive weight gain. Additional CAP risk variables that have been proposed in this research have either not been consistently seen or could not be statistically confirmed.

Adults are significantly more likely to experience morbidity and mortality from Community-Acquired Pneumonia (CAP), it has an annual incidence of 1.07 to 1.2 per 100 individuals. The elderly are disproportionately affected by it; in those 65 years of age or older, the incidence is 14 per 100 persons annually. Hospitalization for pneumonia is linked to increased long-term mortality, and CAP has a substantial long-term impact. It is widely acknowledged that a few risk factors, including underlying medical illnesses and other lifestyle choices, might raise the likelihood of acquiring Community-Acquired Pneumonia (CAP). Understanding these risk variables would enable us to target these particular risk groups with treatments to lower the CAP rate. Based on the OR values in people who have been bedridden for a long time for a lung infection, age, BMI, and History of smoking which is considered as minor risk factors. Diabetes, antibiotic use, and staying up late were deemed to be moderate risk factors. Hormone levels, consciousness problems, nasal feeding, ventilator use, invasive procedures, and length of hospital stay were among the high-risk

factors. In conclusion, increased focus should be placed on the medium-risk factors (diabetes, antibiotic use, length of stay in bed), the low-risk factors (age, BMI, smoking history), and in particular the high-risk factors (hormones, consciousness disorders, nasal feeding, ventilator use, invasive surgeries, length of hospitalization times) to prevent pulmonary infections in patients who are bedridden for an extended period.

Pneumocystis carinii pneumonia, sometimes referred to as *Pneumocystis jirovecii* Pneumonia (PJP) or simply *Pneumocystis pneumonia* is an opportunistic fungal infection that is commonly linked to pneumonia in patients who are nearing the end of their HIV treatment. However, since the introduction of Antiretroviral Treatments (ART) in the treatment of HIV disease, its incidence has been declining also affects non-HIV patients undergoing therapy for cancer hematopoietic stem cell transplantation (high-dose chemotherapy), Solid Organ transplantation (SO), chronic inflammatory illnesses, and other immunosuppressive medications such high dose steroids.

Epidemiology

The most prevalent infectious disease-related cause of death worldwide is Lower Respiratory Tract Infections (LRTIs), which come in fifth place overall. According to epidemiological research, the most common Long-Term Respiratory Infections (LRTIs) are thought to be pneumonia, influenza, bronchitis (including Acute Exacerbations of Chronic Obstructive Pulmonary Disease [AECOPD]), and bronchiolitis.

For every 100 hospital admissions in the US, there are 0.6 to 1.0 nosocomial pneumonia episodes. The majority of instances involve individuals who are not intubated; for intubated patients, the rates increase by six to twenty times. Nosocomial pneumonia is the most frequent nosocomial infection-related cause of death. Crude mortality rates for nosocomial pneumonia patients might range from 20 to 176; generally speaking, individuals in intensive care units using mechanical ventilation were at higher risk of passing away. Global Burden of Diseases (GBD) data from 2019 show that Lower Respiratory Tract Infections (LRTIs), which include pneumonia and bronchiolitis, afflicted 489 million people. The 2019 GBD states that kids under the age of seventy are usually impacted by pneumonia. For the year 2019, there were 489 million incident cases and 11 million frequent cases of LRTI. For every 1,000 people over the age of 70, there were 155.4 LRTI incidences worldwide, according to the 2016 GBD study.

Pneumonia that is obtained outside of a hospital in people who have not been hospitalized in the month prior to the onset of symptoms is known as Community-Acquired Pneumonia (CAP), according to the Etiology of Pneumonia in the Community (EPIC) study carried out in the United States of America. Pneumonia developed after spending at least two days in a hospital and without any prior signs of sickness incubation is known as Hospital-Acquired Pneumonia (HAP). The term for aspiration pneumonia that manifests pneumonia linked to a ventilator that occurs more than 48 h following endotracheal intubation (VAP) [5].

Respiratory illness brought on by breathing in oral or gastrointestinal secretions. It is better to think of it as a component of the range between CAP and HAP rather than as a unique entity. One does not contract pneumonia at a hospital that is related to medical care (HCAP) pneumonia. The highest incidence of CAP was seen in

adults. Adults aged 80 years and beyond (16.4 instances per 1,000 persons) and those aged 65 to 79 (6.3 cases per 1,000 persons). There were 2.4 cases of CAP for every 1,000 adult cases [6].

Categories of Pneumonia

According to anatomy

- Bronchopneumonia: This variant of pneumonia impacts the lungs' bronchus.
- Lobar pneumonia: is an infection that affects only the lobes in the lungs.
- Interstitial pneumonia: is an infection of the interstitial space, which is the space between the alveoli and the surrounding structure.

According to clinical types

- Community-Acquired Pneumonia (CAP): Pneumonia that has been diagnosed in a person who has not yet been admitted to the hospital. Both bacteria—*Haemophilus influenza*, *Mycoplasma pneumoniae*, and *Streptococcus pneumoniae*—are the causative agents of Community-Associated Influenza (CAP) [7].

- Ventilator-acquired pneumonia is a kind of Hospital-Acquired Pneumonia (HAP). HAP is defined as pneumonia that appears two days after being admitted to the hospital and does not show any signs of microbial growth. This type, which manifests within a few days of admission, is caused by Methicillin-Resistant *Staphylococcus aureus* (MRSA) and Methicillin-Susceptible *Staphylococcus aureus* (MSSA) [8].

- Aspiration Pneumonia: Aspiration is described as respiratory difficulties such as choking or airway blockage. Any foreign object that obstructs the airway and is breathed into the lungs might cause aspiration pneumonia.

- Healthcare-Associated Pneumonia (HCAP): Similar to other cases of pneumonia, this one is contracted in an in-hospital care unit. The pathogenic microorganisms in this case may be the same as those found in HAP.

Mechanism of Action of Pneumonia

Pneumonia is a lung inflammation that mostly affects the alveoli. The mechanism of action involves several important phases; Ingress of Pathogens: The most prevalent causes of pneumonia include fungus, viruses, and bacteria. Usually, inhaling airborne droplets, aspirating oropharyngeal contents, or hematogenous dissemination are the ways that infections enter the lungs.

Infection and colonization: Once within the lungs, the infections stay hidden from the host immune system by adhering to the respiratory epithelium. They proliferate and disperse, causing infection. Certain bacteria, such as *Streptococcus pneumoniae*, possess capsules that hinder immune cells from phagocytosing them.

Immune Response: Alveolar macrophages are activated in response to an infection by the immune system, releasing cytokines and chemokines. Alveolar filling with fluid, pus, and cellular debris, as well as inflammation, are caused by the recruitment of neutrophils and other immune cells.

Consolidation and Inflammation: When there is inflammation, the alveoli fill with exudate, which results in consolidation—a solid, airless state of lung tissue. This interferes with gas exchange, leading

to symptoms such as fever, cough, chest discomfort, and dyspnea.

Resolution or Complications: Ideally, the infection is eradicated by the immune system and the alveoli gradually return to normal. Nevertheless, depending on the pathogen's virulence and the host's immunological condition, problems such abscess formation, pleural effusion, or Acute Respiratory Distress Syndrome (ARDS) may occur.

Risk Factors

Pneumonia types differ in their risk factors. A few of these include low birth weight, poverty and socioeconomic factors, undernutrition, zinc deficiency, crowding, low birth weight, exposure to indoor air pollution, and the existence of underlying comorbidities like heart or lung disease and immunodeficiency states like HIV infection, neuromuscular and gastrointestinal disorders like reflux. Children can develop CAP as a result of premature birth, malnutrition, indoor air pollution, inadequate breastfeeding, and other factors; adults can develop risk factors from diseases of the respiratory tract like COPD, heart-related disorders, and liver infections like liver cirrhosis [9,10].

Men and females typically have different morphologies, which makes men more susceptible to CAP. One can categorize the factors that contribute to pneumonia risk into those that are hospital-related or patient-related. One of the risk factors for HAP is oropharyngeal colonization by bacteria. The intensity of this oropharyngeal colonization necessitates placement in intensive care units. The risk of CAP is higher in children under the age of five (46), elderly adults 13, and people with comorbidities who are over 65. The primary risk factors for CAP in children are undernourishment, air pollution in the home, ambient particulate matter, premature birth, and inadequate breastfeeding. Adults with respiratory disorders such as COPD, cardiovascular illness, chronic liver disease, and diabetes mellitus are the most typical concomitant conditions that increase CAP risk. The sociodemographic attributes of the participants: The current study included 403 children under the age of five (50.6% female and 49.4% male) who had 100% of their parents' or guardians' responses. The respondents were divided into three age groups: 1 to 3 (49.9%), less than one (36.2%), and 4 to 5 (13.9%). Sixty-eight percent of the study participants lived in rural areas. There were 299 (51.9%) families with four members or fewer, 31% with five members or more, and 17.1% with fewer than three members. 21% of the respondents were from homes making more than 2500 birr, from families, 37.5% making less than 1000 birr (19.03 USD), and 42.4% of the respondents came from families making between 1000 and 2500 birr (19.03–47.57 USD) a month. Of the fathers, 44.7% were illiterate and of the moms, 113 (47.9%) were. Of the dads, sixty (14.9%) and sixty-two (15.4%) had a diploma or higher in education. 3.2% of the mothers were students, and more than half (63%) were stay-at-home moms.

Pneumonia Risk Factors: A study of univariate logistic regression revealed that Infants who were exclusively breastfed for less than two years were 1.75 times more likely to get pneumonia than those who were nursed for more than two years. The multivariate logistic analysis showed that children from homes using wood for cooking were 3.5 times more likely to get pneumonia than children from households using children from homes with wood stoves had a 3.5-fold higher risk of pneumonia than children from homes with electricity. Compared to exclusive breast feeders, children who received mixed nursing had a 5.2-fold higher risk of pneumonia. The risk of pneumonia was twice as high for children who began receiving supplemental food before the age of six months as for those

who began beyond that period. Another risk factor for Spontaneous Pneumothorax (SP) and Spontaneous Pneumomediastinum (SPM) is cigarette smoking [11,12].

Acute Respiratory Illnesses (ARIs) are a primary cause of illness, and death in both developed and poor countries. Every year, almost 13 million children under the age of five die. Worldwide; 95% of these deaths occur in underdeveloped nations, and ARI accounts for one-third of all deaths. Seven out of ten deaths in the under-five age group in poor nations are attributable to ARI [13].

This study's main objective was to look into the risk factors for pneumonia. The goals were to determine the standard of living among families whose children had pneumonia, to assess the nutritional status of children under five about pneumonia, and to create baseline data and statistical information about pneumonia in children under five for future use. In this study, forty moms who had taken their sick, pneumonia-stricken children under five to the Mohamed Al Amin Hamid Pediatric Hospital in Omdurman City participated.

According to the study's findings, there were 23 male children (57.50%) and 17 female children (42.50%). There was a correlation between gender and pneumonia, as evidenced by the gender distribution, which showed that pneumonia was more common in male children than in female children.

The study's findings showed a correlation between family income and pneumonia, with pneumonia being more common in households with lower incomes (26, 65.00%). According to Park's 2007 study, children from poorer socioeconomic origins usually have higher rates of respiratory illnesses.

The following children's lower respiratory tract infections are linked to risk factors below five: Child mortality considered for a long time as a trustworthy a measure of the nation's socioeconomic development. Even though India's baby and under-five mortality rates have dramatically dropped, there are still reasons to be concerned, especially in the less developed regions of the country. Acute respiratory infections continue to be among the main reasons for dying in our country for children below five, accounting for 15.9% of fatalities due to complications associated with preterm delivery. Hospital data due to a high death rate indicate that as much as 13% of inpatient fatalities in pediatric department in India are related to Acute Respiratory Infections (ARIs).

Worldwide, the most common Respiratory Syncytial Virus (RSV) is often identified infection in pediatric Acute Lower Respiratory Infections (ALRIs) and a leading cause of hospital admissions. In children under five, an estimated 33.1 million new RSV-ALRI episodes occurred globally in 2015; around 10% of these patients required hospitalization due to their severity. RSV-related deaths in children younger than five years old are estimated to have occurred in 2015 in roughly 59,600 hospital deaths worldwide and 118,200 deaths overall; poor countries accounted for 99% of these deaths [14,15].

Although more severe types of sickness and mortality are disproportionately higher in developing nations, ALRI incidence is high even in the industrialized world. It has been demonstrated that several variables, including low birth weight, the timing of breastfeeding initiation, weaning with supplemental food, vaccination status, and others, affect the risk of ALF in children under five years old [16,17]. When speculating on the possible etiology of ALRI, it is also important to consider additional sociodemographic

risk factors, including exposure to biomass fuels, kerosene lamps, parental education, overcrowding, socioeconomic level, and passive smoking. With the use of suitable health education campaigns and other focused community development initiatives, preventative efforts would benefit from an understanding of risk factors.

Recognizing risk factors for respiratory tract infection:

Factors related to demographics: A person's risk of RTIs is influenced by their age, sex, and socioeconomic status, among other demographic factors. The immune systems of infants, young children, and the elderly are impaired, making them especially susceptible. There have also been noted gender disparities in the susceptibility to respiratory infections, with men frequently exhibiting more severe symptoms.

Environmental elements: The risk of respiratory infections is influenced by a variety of environmental factors, such as work hazards, allergies, and air pollution. Particulate matter and nitrogen dioxide are two examples of airborne contaminants that can weaken respiratory defenses and aggravate pre-existing respiratory disorders. Additional dangers are posed by indoor contaminants such mold, tobacco smoke, and second-hand smoke, particularly in poorly ventilated areas. Biological agents, chemicals, and dust exposure at work raise the risk of respiratory illness in certain occupational groups.

Individual behaviors like smoking, not washing your hands properly, and not getting enough vaccinations all have a significant impact on your risk of RTI. Smoking tobacco increases the vulnerability to infections by harming respiratory epithelial cells and reducing mucociliary clearance.

Clinical Presentation and Diagnosis

Cough, fever, chills, dyspnea, pleuritic chest discomfort, and exhaustion are typical symptoms. Consider the patient's medical history, including any recent respiratory infections, contact with sick people, smoking history, and underlying medical issues.

Physical assessment such as using auscultation, check for unusual breath sounds like crackles or wheezing. Blood cultures and the A higher White Blood Cell count (WBC) on a Complete Blood Count (CBC) suggests infection, are examples of laboratory testing.

Microbiological testing, sputum gram staining, and CT (Computed Tomography) scans are examples of imaging methods.

Recommended Diagnostic Testing

Radiographic imaging

For the detection of pneumonia, a low-cost, easily accessible plain chest radiograph is recommended. The appearance of when combined with clinical characteristics, a fresh infiltrate on a chest radiograph is regarded as the "gold standard" for choosing patients for clinical studies. A chest radiograph can provide further details regarding the location, intensity, and side effects of pneumonia, including cavitation, pleural effusion, multi-lobe disease, and acute respiratory distress condition.

Microbiological examination

Regular blood tests and radiographs are not sensitive enough or specific enough to back up treatment recommendations for a particular patient. An extremely low white Blood Cell Count (WBC) could indicate neutropenia, which could indicate serious pneumonia,

or it could raise suspicions about particular infections if the patient has a history of immune suppression.

Other cultures and blood culture

Although the usage of broad-spectrum antibiotics, hospital stays, and expenses can all rise as a result of false-positive blood culture results, most studies have demonstrated that pneumococcal bacteremia is associated with an increased mortality rate. Only in cases of severe CAP are blood cultures recommended by the IDSA/ATS recommendations due to their limited yield and reduced therapeutic value.

Management of Community-Acquired Pneumonia

When is it appropriate to take a chest radiograph in a public setting?

A chest X-ray is not necessary for people who have been suspected of having CAP unless:

- The diagnosis is uncertain and the radiograph would aid in the differential diagnosis and acute illness care.
- Treatment for suspected CAP has not shown satisfactory results at review.
- The patient is thought to be at risk for underlying lung pathology, including lung cancer.

When is the right time for a hospital to take a chest radiograph?

As soon as it is practical, a chest radiograph should be taken for every hospitalized patient who is suspected of having CAP to support or contradict the diagnosis. As soon as a chest radiograph is obtained, the goal of any service should be to diagnose CAP so that the patient can receive antibiotics within four hours of being admitted to the hospital.

While recuperating, when should a second chest radiograph be taken?

Before being discharged from the hospital, individuals who have made a satisfactory clinical recovery from CAP do not need to have their chest radiographs taken again. For every patient, a chest radiograph should be scheduled approximately every six weeks, regardless of whether they have been admitted to the hospital, show persistent symptoms or physical indicators, or are more likely to have an underlying cancer (particularly in smokers and older adults). Individuals who have persistent symptoms and abnormalities on radiographs ought to be assessed for additional investigations, which may include bronchoscopy, six weeks after the end of medication.

When a patient is admitted to the hospital, what standard investigations ought to be conducted?

Upon admission, all patients ought to undergo the following examinations:

- Arterial blood gases and oxygenation saturation as required by the BTS Emergency Oxygen Use Guidelines for Adult Patients. A reliable diagnosis can be made using a chest radiograph [16].
- Using electrolytes and urea to help in severity evaluation.
- C-reactive protein as a starting point and diagnostic tool.

Empirical selection of antibiotics for community-treated adults

The recommended medication for individuals receiving community-based treatment is still amoxicillin, 500 mg three times per day. For people who are hypersensitive to penicillin, either doxycycline or clarithromycin is a suitable substitute medication. Individuals exhibiting indicators of a moderate or severe infection ought to be admitted to the hospital at least once.

- o When possible, it is possible to confirm the diagnosis using a chest X-ray.

- o Five days should be sufficient for patients undergoing community-based treatment for low-severity pneumonia; Extended courses could be required for patients who are hospitalized or those with more severe pneumonia.

- o If after three days the symptoms don't go better as predicted, think about extending the course for more than five days.

- o The therapy of community-dwelling individuals with CAP is determined by the evaluation of adult patients using clinical judgment informed by the CRB-65 score. Each of the following attributes carries one point on the CRB-65 death risk score.: Difficulty (average 30/min). A blood pressure reading (SBP<60 mmHg). More than 65 years old.

- o Look for *Staphylococcus aureus* and *Legionella* infection risk factors.

1. Exposure to artificial water systems such as spa pools, cooling towers, air conditioning units, and recent trip are among the possible causes of *Legionella*.

2. These can include people who are chronically sick or incapacitated, residents of nursing homes, aspiration, and recent influenza infections (*S. aureus*). Approximately every four years, there are epidemics of mycoplasma pneumoniae infection. Clinical judgment is used to determine the severity in children and young adults.

If symptoms and signs are severe, admit the patient to the hospital right now.

Depending on how serious the illness is, patients with Risk Elements for Resistance (RFR) are treated differently. It is not advised to utilize empirical broad-spectrum medication in non-severe CAP patients with reduced function. Samples of sputum from these patients should be taken. If a resistant pathogen is discovered, therapy may be focused on it; however, if the patient has shown a good response to conventional CAP treatment, it seems sensible to keep treating the patient without treating the isolated germs. In individuals with severe CAP and RFR, empirical use of broad-spectrum treatment makes sense.

Fever management: In our neurosurgical intensive care unit, we conducted routine testing on the patient's sputum, blood, and urine when a fever was noticed. The infection control unit assessed and managed every case of infection.

Antibiotic timing: There is evidence linking postponing antibiotic medication to a higher risk of mortality. As a result, an accurate diagnosis needs to be made quickly. According to two significant studies, shorter hospital stays and higher survival rates in cases of community-acquired pneumonia are linked to prompt

diagnosis and early antibiotic therapy [18,19].

Therapy duration: As of right now, an outpatient with mild-to-moderate CAP should receive treatment for no more than seven days if they have not experienced fever for 48 h to 72 h or indications of an extrapulmonary infection.

The Aftereffects of a Lower Respiratory Infection

There is no known overall death rate from lower respiratory tract infections acquired in the community, although, among patients who require hospital admission, the rate of death ranges from five to fifteen percent. Less frequent side effects include localized bronchiectasis, empyema, and the formation of an abscess. Even though a pleural effusion is typically sterile, it should always be tapped for Gram staining and culture. Chest radiographs typically show pleural effusions.

Having an underlying bronchial tumor or a staphylococcal infection are common cause of emphysema.

Endocarditis, osteomyelitis, septic arthritis, and cerebral abscess are a few examples of concurrent infection sites that can arise when pneumonia develops as a part of septicemia. Ventilation and special care may be necessary if severe pneumonia becomes rapidly fatal [20,21].

Prevention

Treatment or prevention of HIV infection; prophylactic antibiotics for immunocompromised children; prevention of HIV transmission from mother to child; full immunization; improving living conditions to prevent crowding; avoidance of tobacco smoke exposure from conception through childhood; and reduction of indoor air pollution are some of the strategies that can be employed to prevent or lessen the effects of childhood respiratory diseases; and improving breastfeeding and childhood nutrition, both of which improve immune status. A few of these interventions are also suitable for adult respiratory illness prevention. Even though vaccination is one of the biggest advancements in contemporary public health, many children remain unvaccinated against diseases that can be prevented, especially in low- and middle-income nations. The majority of vaccine-preventable diseases are found in countries with the lowest immunization rates, which also have the highest rates of childhood death.

For many of these pediatric illnesses to be controlled and eradicated, vaccinations are necessary. Important advancements in the prevention of pneumonia have been made with the development of novel combination vaccines against *Haemophilus influenzae* type b and *Streptococcus pneumoniae* (pneumococcus).

To avoid lower respiratory tract infections, vaccinations against germs including *S. pneumoniae*, *H. influenzae* type b, and whooping cough (pertussis) are quite effective. Influenza can be prevented with the use of the influenza vaccination. Other virus-related vaccinations, like the one against measles, are almost completely eradicating the illness. Although antibiotic-resistant bacteria can complicate therapy, antibiotics have made most cases of bacterial pneumonia treatable. Among the important areas of intervention are hygienic measures to stop the transmission of resistance strains amongst people, include hand cleanliness, screening for the existence of resistant strains, and isolating positive patients. Usage of antibiotics that are now on

the market with caution in both humans and animals, giving them only, when necessary, for the right diagnosis, and at the right dosage, intervals, and length of time [21,22].

Conclusion

Globally, chest infections continue to be a major health concern, impacting a wide range of communities to differing degrees of severity. The many facets of chest infections, including their etiology, epidemiology, treatment, and clinical symptoms, have been brought to light by this thorough review. In summary, to effectively battle chest infections and lessen their impact on global health, a multidisciplinary approach combining medical professionals, researchers, and policymakers is needed. We can create a healthier future with lower morbidity and mortality from chest infections by being aware of the comprehensive risk factors and taking preventative action.

References

1. Coronado K, Herrada C, Rojas DA. Role of the fungus pneumocystis in IL1 β pathway activation and airways collagen deposition in elastase-induced COPD animals. *Int J Mol Sci.* 2024;25(6):3150.
2. Langford B. Antibiotic prescribing recommendations in COVID-19: A systematic survey (Doctoral dissertation). 2023.
3. Chhetri KR, Chouhan APS, Singh V, Shaida B. Radio imaging of pneumonia: A comprehensive review. *J Commun Dis.* 2022;54(3):75-87.
4. Soriano JB, Kendrick PJ, Paulson KR, Gupta V, Abrams EM, Adedoyin RA, et al. Prevalence and attributable health burden of chronic respiratory diseases, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet Respir Med.* 2020;8(6):585-96.
5. Mandell LA, Niederman MS. Aspiration pneumonia. *N Eng J Med.* 2019;380(7):651-63.
6. NICE. Pneumonia in adults: diagnosis and management. 2019.
7. Cilloniz C, Liapikou A, Ceccato A, Torres A. Risk factors for community-acquired pneumonia in adults. *Minerva Pneumol.* 2017;56(3):206-16.
8. Feikin DR, Fu W, Park DE, Shi Q, Higdon MM, Baggett HC, et al. Is higher viral load in the upper respiratory tract associated with severe pneumonia? Findings from the PERCH study. *Clin Infect Dis.* 2017;64(suppl_3):S337-46.
9. He XL, Liao XL, Xie ZC, Han L, Yang XL, Kang Y. Pulmonary infection is an independent risk factor for long-term mortality and quality of life for sepsis patients. *Biomed Res Int.* 2016;2016:4213712.
10. Llinares P, Menéndez R, Mujal A, Navas E, Barberán J. Guidelines for the management of community-acquired pneumonia in the elderly patient. *Rev Esp Quimioter.* 2014;27(1):69-86.
11. Zar HJ, Ferkol TW. The global burden of respiratory disease—impact on child health. *Pediatr Pulmonol.* 2014;49(5):430-4.
12. Wang KW, Chen HJ, Lu K, Liliang PC, Huang CK, Tang PL, et al. Pneumonia in patients with severe head injury: Incidence, risk factors, and outcomes. *J Neurosurg.* 2013;118(2):358-63.
13. Brar NK, Niederman MS. Management of community-acquired pneumonia: A review and update. *Ther Adv Respir Dis.* 2011;5(1):61-78.
14. Almirall J, Bolibar I, Serra-Prat M, Roig J, Carandell E, Agustí M, et al. New evidence of risk factors for community-acquired pneumonia: A population-based study. *Eur Respir J.* 2008;31(6):1274-84.
15. Metlay JP, Waterer GW, Long AC, Anzueto A, Brozek J, Crothers K, et al. Diagnosis and treatment of adults with community-acquired pneumonia. An official clinical practice guideline of the American Thoracic Society and Infectious Diseases Society of America. *Am J Respir Crit Care Med.* 2019;200(7):e45-e67.

16. Lim WS, Baudouin SV, George RC, Hill AT, Jamieson C, Jeune IL, et al. BTS guidelines for the management of community acquired pneumonia in adults: Update 2009. *Thorax*. 2009;64(Suppl 3):iii1-55.
17. Seth S, Ganguly S, Satpathy SK. Risk factors responsible for lower respiratory tract infections in children aged under five: A hospital based study. *Int J Contemp Pediatr*. 2020;7(7):1578-83.
18. Brown JD, Harnett J, Chambers R, Sato R. The relative burden of community-acquired pneumonia hospitalizations in older adults: A retrospective observational study in the United States. *BMC Geriatr*. 2018;18(1):92.
19. Shi T, Vennard S, Mahdy S, Nair H. Risk factors for RSV associated acute lower respiratory infection poor outcome and mortality in young children: A systematic review and meta-analysis. *J Infect Dis*. 2021.
20. Noorani TA, Kothari J, Kanabar B. A cross sectional study on risk factors associated with acute lower respiratory tract infection among children aged 6 to 60 months. *Int J Contemp Pediatr*. 2020;7(11):2219.
21. Hosker HS, Jones GM, Hawkey P. Fortnightly review: Management of community acquired lower respiratory tract infection. *BMJ*. 1994;308(6930):701-5.
22. Yathiraj KM, Gudur SP. Strategic approaches to managing respiratory tract infection risk factors: A comprehensive review. *Asian J Hosp Pharm*. 2024;4(1):1-8.